



IPM PACKAGE NO. 23



INTEGRATED PEST MANAGEMENT PACKAGE

FOR OKRA



Government of India
Ministry of Agriculture
Department of Agriculture & Cooperation
Directorate of Plant Protection, Quarantine & Storage
N.H. IV, Faridabad - 121 001
May, 2001



Government of India
Ministry of Agriculture
(Department of Agriculture & Cooperation)

DIRECTORATE OF PLANT PROTECTION, QUARANTINE & STORAGE
NH IV, FARIDABAD - 121 001 (Haryana)


DR. V. RAGUNATHAN
Plant Protection Adviser
To the Government of India

FORWARD

Integrated Pest management (IPM) approach has been globally accepted for achieving sustainability in agriculture. It has become more relevant due to a number of advantages like safety to environment, pesticide-free food commodities, low input based Crop Production Programme etc. Though IPM approach has been taken up since 1981, its impact has not been felt until 1994. Human Resource Development has helped to sensitise extension functionaries and farmers about the usefulness of IPM.

For successful implementation of IPM, the scattered information on various components of this eco-friendly approach forms basic necessity. In this direction, initial attempts were made in 1992 to harmonise the IPM Package of Practices of various crops. Subsequently concerted efforts were made in 1998, 2001 and 2002 to update and develop IPM package of practices for agricultural and horticultural crops. Presently, IPM package of practices for 51 crops have been finalised to help the extension workers and farmers to manage the pests/ diseases and to minimise the over use/ misuse of chemical pesticides. Efforts have been made to incorporate the relevant available technical input provided by the scientists of ICAR Institutes/ SAUs and State Departments of Agriculture. However, suggestions for further improvement in future publication/ revision will be of immense help. Hopefully, these IPM Package of Practices will be useful for the Researchers, Plant Protection Workers and Farmers alike.

April 1, 2002


(V. Raguathan)

P R E F A C E

In order to minimise the indiscriminate and injudicious use of chemical pesticides, INTEGRATED PEST MANAGEMENT (IPM) has been enshrined as cardinal principle of Plant Protection in the overall Crop Protection Programme under the National Agricultural Policy of the Govt. of India. IPM is an eco-friendly approach for managing pest and disease problems encompassing available methods and techniques of pest control such as cultural, mechanical, biological and chemical in a compatible and scientific manner. The greater emphasis has been given on biological control including use of biopesticides.

With a view to provide technical knowledge to the extension functionaries and farmers in the States, first National Workshop on IPM for harmonisation of Package of Practices was organized at National Plant Protection Training Institute (NPPTI), Hyderabad during June 29-30, 1992. Subsequently workshops were organized from April 15-17, 1998 and Nov. 5-6, 1998 at Directorate of Plant Protection, Quarantine & Storage, Faridabad and IPM package of practices for 20 crops were evolved on rice, cotton, vegetables, pulses, and oilseeds. In this series, two National Workshops on IPM have been conducted at NPPTI, Hyderabad and Dte. of PPQ&S, Faridabad during May 14-17, 2001 and Feb. 20-22, 2002 respectively to update 20 available IPM Packages and develop 31 new IPM Packages specially for Horticultural crops. In these workshops, 51 IPM Package of Practices for cereal crops (Rice, Wheat, Maize, Sorghum, Millets), commercial crops (Cotton, Sugarcane, Tobacco, Tea), pulse crops (Pigeonpea, Gram, Black gram/Green gram, Pea, Rajma), oilseeds (Groundnut, Soybean, Rapeseed/Mustard, Sesame, Safflower, Castor, Sunflower, Oilpalm), vegetables (Potato, Onion, Tomato, Brinjal, Okra, Chillies, Cruciferous vegetables, Leguminous vegetables, Cucurbitaceous vegetables), fruit crops (Citrus, Banana, Apple, Mango, Guava, Grapes, Pineapple, Sapota, Pomegranate, Litchi), spice and plantation crops (Small Cardamom, Large Cardamom, Black Pepper, Ginger, Coriander, Cumin, Fennel, Coconut, Cashew and Arecanut) have been finalised.

IPM technology manages the pest population in such a manner that economic loss is avoided and adverse side effects of chemical pesticides are minimized. The IPM packages encompasses various management strategies for containing the pest and disease problems. Pest monitoring is also one of the important component of IPM to take proper decision to manage any pest problem. It can be done through Agro-Ecosystem Analysis (AESA), field scouting, light, pheromone, sticky/yellow pan traps. The economic threshold levels (ETL) of important pests and diseases are also given in the packages to take appropriate control measures when pest population crosses ETL.

These IPM packages developed with the technical inputs from experts from Indian Council of Agriculture Research, State Agricultural Universities, Central Directorate of Plant Protection, Pesticide Industries and State Departments of Agriculture/Horticulture will provide technical backup in the management of pests, diseases, weeds, nematodes and rodents in the Indian Agriculture and Horticulture. These will also be useful in reducing the pesticide residues in exportable agricultural commodities and would also help in the management of pests/diseases/weeds/nematodes which may get inadvertently introduced in the country.

IPM Package of Practices for Agricultural and Horticultural crops will be helpful to minimize the ill effects of chemical pesticides to promote the IPM for sustainable production. These packages will be useful for the researchers, extension workers and farmers alike who are engaged in the agricultural practices.



(A.D. Pawar)
Director (IPM)

April 1, 2002

ACKNOWLEDGEMENTS

The IPM Package of Practices for Okra crop was discussed and finalised in the National Workshop on IPM held at National Plant Protection Training Institute (NPPTI), Hyderabad during May 14-17, 2001. The technical input received from the following experts is thankfully acknowledged.

- I. Chairman, Technical Session : Dr. A.D. Pawar, Director (IPM), Dte of PPQS, Faridabad
- II. Coordinator, Technical Session : Dr. M.P. Misra, Dy. Director (Ent.), Dte. of PPQS, Faridabad
- III. Co-chairman, Technical Session : Dr. R.B.L. Bhaskar, Jt. Director (Bio.) Dte of PPQS, Faridabad
- IV. Expert input :
 1. Dr. B.J. Divakar, Sr. Entomologist, NPPTI, Hyderabad
 2. Dr. N.S. Rao, Sr. Scientist (Ent), PDBC, ICAR, Bangalore
 3. Dr. U. Sreedhar, Sr. Scientist CTRI (ICAR), Rajahmundry
 4. Dr. C.M. Srivastava, PPO(PP), Central IPM Centre, Gorakhpur
 5. Dr. K.S. Kapoor, PPO (E), Cnetral IPM Centre, Jalandhar
 6. Sh. M.V. Srinivasa, PPO(E), Central IPM Centre, Bangalore
 7. Dr. D. Karungo, Jt. Director, CIL, Dte of PPQS, Faridabad
 8. Dr. Brajendra Singh, Entomologist, Dte of PPQS, Faridabad
 9. Dr. B.S. Phogat, Agronomist (H), Dte of PPQS, Faridabad
 10. Dr. Jasvir Singh, AD(E), Dte of PPQS, Faridabad
- V. Technical input:
 1. Sh. S.P. Singh, APPO, IPM Div., Dte of PPQS, Faridabad.
 2. Sh. K.S. Sharma, SSA, IPM Div., Dte of PPQS, Faridabad.
 3. Sh. Yogesh Kumar, SSA, IPM Div., Dte of PPQS, Faridabad.
 4. Sh. R.S. Tomer, SSA, IPM Div., Dte of PPQS, Faridabad.
 5. Sh. Laxmi Chand, SSA, IPM Div., Dte of PPQS, Faridabad.
 6. Mohd. Abrar Alam, Stenographer, IPM Div., Dte of PPQS, Faridabad.
 7. Sh. N.K. Mishra, LDC, IPM Div., Dte of PPQS, Faridabad.

IPM PACKAGE FOR OKRA/BHINDI

Contents

SUBJECT	Page No.
Foreward	i
Preface	ii
Acknowledgements	iii
I. MAJOR PESTS	
A. Pests of National Significance	1
B. Pests of Regional Significance	1
II. PEST MONITORING	
A. Agro-eco-system Analysis (AESAs)	2
B. Field scouting	2
C. Pheromones	3
D. Yellow water pan/sticky traps	3
E. Root Knot Nematode	3
F. Economic threshold levels (ETL)	3
III. IPM STRATEGIES FOR OKRA/BHINDI	
A. Cultural Practices	4
B. Mechanical control	4
C. Biological control	4
D. Biopesticides	5
E. Botanical pesticides	5
F. Chemical control	5
IV. OKRA CROP STAGE-WISE IPM PRACTICES	6-7
V. DO'S AND DON'TS IN OKRA IPM	8-10
VI. SAFETY PARAMETERS IN PESTICIDES USAGE	11-13
ANNEXURES - I & II	14-21

IPM PACKAGE FOR OKRA/BHINDI

I. MAJOR PESTS

A. Pests of National Significance

I Insect Pests

- 1.1 Jassids
- 1.2 Fruit borer
- 1.3 Aphid
- 1.4 White fly
- 1.5 American bollworm

2. Diseases

- 2.1 Damping off
- 2.2 Yellow vein mosaic virus
- 2.3 Powdery mildew
- 2.4 Leaf Spot

B. Pests of Regional Significance

1. Insect Pests

- 1.1 Red cotton bug
- 1.2 Green leaf hopper
- 1.3 Spider mites
- 1.4 Grey weevils
- 1.5 Stem fly

2. Major Weeds

- 2.1 Lamb square (*Chenopodium album*)
- 2.2 Pimpernel (*Anagallis arvensis*)
- 2.3 Sweet clover (*Melilotus* spp.)
- 2.4 Fumitory (*Fumaria indica*)
- 2.5 Corn spray (*Spurgula arvensis*)
- 2.6 Blue grass (*Poa annua*)
- 2.7 Canary grass (*Phalaris minor*)
- 2.8 Rabbit foot grass (*Dactyloctenium aegyptium*)

3. Nematodes

- 3.1 Root knot nematode
- 3.2 Reniform nematode

4. Diseases

- 4.1 Wilt
- 4.2 Pod spot
- 4.3 Collor rot

II. PEST MONITORING

A. Agro Eco System Analysis (AESA)

AESA is an approach, which can be gainfully employed by extension functionaries and farmers to analyse field situations with regard to pests, defenders, soil conditions, plant health, the influence of climatic factors and their interrelationship for growing healthy crop. Such a critical analysis of the field situations will help in taking appropriate decision on management practices. The basic components of AESA are:-

1. Plant health at different stages.
2. Built - in-compensation abilities of the plants.
3. Pest and defender population dynamics.
4. Soil conditions.
5. Climatic factors.
6. Farmers' past experience.

The details of the AESA are given in Annexure - I.

B. Field scouting

AESA requires skill and so only the trained farmers can undertake their exercise. However, other farmers also can do field scouting in their own fields at regular intervals to monitor the major pest situation. Simple field scouting on pest situation by the farmers helps to minimize pesticides usage to a large extent.

C. Pheromones

Pheromones traps with lures are commercially available for pests like *Helicoverpa armigera*, *Spodoptera litura*, *Earias* sp. Install 5 traps per ha with lures for each pest. Traps should be installed in the field in such a way the position of lure is always 6 to 12 inches above the crop canopy. Replace the lures once in 15 to 25 days depending upon the weather conditions. The trapped moths should be collected and killed daily. ETL for *Helicoverpa armigera* is 8 to 10 moths per day per trap.

D. Yellow water pan/sticky traps

Set up yellow pan/sticky traps for monitoring whitefly, thrips etc. @ 10 traps per ha. Locally available empty yellow Palmolive tins coated with grease/vaseline/castor oil on outer surface may also be used as yellow pan trap.

E. Root knot nematode

Root knot nematode causes galls and reniform nematode causes "dirty roots". Their presence can be deducted by using "Trypan Blue" stain, which turns nematode egg sacs dark blue in colour, whereas the roots remain uncoloured.

F. Economic threshold levels (ETL)

The Economic threshold level (ETL) is an attempt to improve decision making practices by using partial economic analysis on the impact of the control practice such as spraying a pesticide. At the ETL the benefit of spraying is equal to the losses caused by the insects in the field. The farmers are advised to take up the appropriate measures, whenever the incidence crosses ETL. The ETL for some major pests are listed below.

PEST	ETL
Leaf hopper	2.5 nymphs/leaf
White fly	4 adults/leaf
Shoot borer	1 infected plant/meter row
Spider mite	2 mites/leaf

III. IPM STRATEGIES FOR OKRA / BHINDI

A. Cultural Practices

1. Field sanitation and removing alternate hosts (Wild/Malvaceous) helps the crop free from pests.
2. Crop seedling should be planted timely in well prepared field at recommended spacing. Use balanced dose of fertilizer for obtaining optimum plant population and healthy crop stand, which would be capable of competing with weeds at initial stages of crops growth.
3. Crop rotation with non-host crops of root knot & reniform nematodes eg. Wheat, Sorghum. Inter-cropping with marigold, onion, garlic is also useful.
4. Avoid excess use of water for disease management.
5. Grow less susceptible varieties Punjab Padmini, Arka Anamica, IC 7194, IC-13999 for yellow vein mosaic virus.

B. Mechanical Control

- (a) Destruction of infected fruits.
- (b) Installation of pheromone traps for *Earias* and *Helocoverpa* @ 5 per ha.
- (c) Yellow pan/sticky traps @ 10 per ha. for monitoring white flies.
- (d) Crop should be maintained weed free for 4-6 weeks after planting by resorting to timely hand hoeing/hand weedings.

C. Biological Control

1. Conservation

- 1.1 Conserve parasites like *Erythmelus empoascae*, *Stethynium empoascae*, *Anagrus empoascae* which are active against *Amrasca bigutella*; parasites like *Microbracon lefroyi*, *Rogas testaceus*, *Raligarensis*. *Bracon hebetor*, *B.greeni* and *Tricogramma sp.* on *Earias* are found active.
- 1.2 *Aphelinus flavipes* parasitizes on *Aphis gossypii* and *Eurytoma sp.* parasitizes on *M.hibisci*.
- 1.3 In addition to these parasites, general predators like Coccinellids, Syrphids, Spiders, Carabids, Staphylinids, Dragonfly, Damselfly, predatory miridbugs, predatory pentatomids nabid bug, reduvid bug, Anthocoridbug. Geocorid bug, predatory mite, predatory thrips also actively suppress the pest population. Since these natural enemies are highly prone for pesticides, avoiding unnecessary sprays are the best way to conserve them.

2. Augmentation

1. Release of *Trichogramma priteosum*/*T.chilonis* @ 50,000 / ha. against *Helicoverpa armigera* and *Earias* sp.
2. Release *Chrysoperla carnea* @ 2 grubs per plant 2 times at weekly interval against whitefly and other soft bodied insects.
3. Spray HaNPV 250 LE/ha. for 2 times at a 15 days interval in the evening hours.

D. Biopesticides

1. Seed treatment with *Trichoderma viride* @ 4 gm per kg seed.

E. Botanical pesticides

1. Use of 5 per cent NSKE against *Earias* when infestation is noticed during vegetative stage.
2. Application of neem cake @ 200 kg per ha as a basal dose at the time of land preparation for controlling root knot nematode infestation.

F. Chemical control

1. Chemical pesticide should be used on need basis as a last resort, only when pest population intensity cross economic threshold level, the safer pesticides should be applied judiciously.
2. Apply fluchloralin @ 0.9 to 1.20 kg ai/ha as preparing soil incorporation for weed management.
3. Phosalone 500 g ai/ha, Endosulfan 525-650 g ai/ha, Fenvalerate 60-75 g ai/ha are effective against *Earias*.
4. Sulphur 80% WP @ 3.13 kg ai in 750-1000 lit water/ha against mildew.
5. Seed treatment with *carbosulfan (25 ST) @ 30 per cent (w/w) reduces sucking pest.

* Not as per approved usage under Insecticide Act, 1968

IV. OKRA CROP STAGE WISE IPM PRACTICES

Stage	Pest	Practice
1. Pre-sowing	Exposing resting stages of Pests	Deep summer ploughing. Application of more FYM. Apply Neem-cake @ 200 kgs. per ha.
	Weeds	Apply Fluchloralin @ 0.9 to 1.20 kg ai/ha as preplanting soil incorporation.
	Nematode	Application of neem cake @200 kgs per ha
2. Seed & Vegetative Stage	Whitefly and yellow vein mosaic	Roguing of affected plants. Install yellow pan/sticky traps @ 5/ha. Release <i>Chrysoperla carnea</i> @ 2 grubs per plant.
	Weeds	Timely planting, optimum spacing; application of balanced dose of fertiliser.
		Maintain weed free field for 4-6 weeks after planting by hand hoeing/hand weeding.
	Collar Rot	Seed treatment with <i>Trichoderma viride</i> @ 4 g/kg seeds.
	Jassids	Release <i>Chrysoperla carnea</i> @ 2 grubs per plant.
Earias sp.	Remove affected shoots and destroy. Install Pheromone traps @ 5/ha.	

<p>3. Reproductive</p>	<p>Whitefly</p> <p><i>Earias</i> sp.</p> <p>American bollworm</p>	<p>Release of <i>Trichogramma priteosum</i>/<i>T.chilonis</i> @ 50,000/ha. for 6 times at weekly intervals.</p> <p>Spray <i>Bacillus thuringiensis</i> var <i>kurstaki</i> 500 g/ha.</p> <p>Spray 5% NSKE.</p> <p>Spray endosulfan 525-650 g ai/ha phasolone 500g ai/ha fenvalerate 60-75 g a.i./ha.</p> <p>As shown in vegetative stage</p> <p>As shown in vegetative stage.</p> <p>Install pheromone traps @ 5 per ha with replacement of lures once in 15 days.</p> <p>Release <i>Trichogramma brasiliensis</i>/<i>T.priteosum</i>/<i>T.chilonis</i> @ 50,000/ha. for 6 times at weekly interval.</p> <p>Hand collection of larvae of <i>Helicoverpa armigera</i> and spray Ha NPV @ 250 LE per ha (6×10^{12} PIB/ha) at a 15 days interval in the evening hours. Spray 5% NSKE or <i>Bacillus thuringiensis</i> var <i>kurstaki</i> 500 g/ha.</p>
------------------------	---	---

V. DO's AND DON'Ts IN OKRA IPM

Do's	Don'ts
1. Deep ploughing is to be done on bright sunny days during the months of may and June. The field should be kept exposed to sun light atleast for 2-3 weeks.	Do not plank or irrigate the field after ploughing, atleast for 2-3 weeks, to allow desiccation of weeds bulbs and/or rhizomes of perennial weeds.
2. Adopt crop rotation	Avoid growing monocrop.
3. Grow only recommended varieties.	Do not grow varieties not suitable for the season or the region.
4. Sow early in the season.	Avoid late sowing as this may lead to reduce yields and incidence of white grubs and diseases.
5. Always treat the seeds with approved chemicals/bio products for the control of seed borne diseases/pests.	Do not use seeds without seed treatment with biocides/chemicals.
6. Sow in rows at optimum depths under proper moisture conditions for better establishment.	Do not sow seeds beyond 5-7 cm depth.
7. Pre-emergence herbicides should be applied immediately after sowing before the emergence of weeds and crop.	Pre-emergence herbicides should not be applied after emergence of crop or weeds as they cannot control the germinated weeds as well as they may cause phytotoxicity to the crop.
8. Herbicides like fluchloralin should be incorporated into the soil immediately after spraying, to avoid its photo degradation.	Soil incorporation of fluchloralin should not be delayed or avoided for achieving effective weed control.
9. Apply only recommended herbicides at recommended dose, proper time, as appropriate spray solution with flat fan or flat jet nozzels.	Pre-emergence as well as soil incorporated herbicides should not be applied in dry soils. Do not apply herbicides alongwith irrigation water or by mixing with soil, sand or urea.

<p>10. Maintain optimum and healthy crop stand which would be capable of competing with weeds at a critical stage of crop weed competition.</p>	<p>Crops should not be exposed to moisture deficit stress at their critical growth stages.</p>
<p>11. Use NPK fertilisers as per the soil test recommendations.</p>	<p>Avoid imbalanced use of fertilizers.</p>
<p>12. Use micronutrient mixture after sowing as top dressing separately.</p>	<p>Do not mix micronutrients with fertilizers and incorporate into the soil.</p>
<p>13. Conduct AESA weekly in the morning preferably before 9 a.m. Take decision on management practice based on AESA, ETLand P:D ratio only.</p>	<p>Do not apply chemical pesticides on calendar basis.</p>
<p>14. Install pheromone traps at appropriate period.</p>	<p>Do not store the pheromone lures at normal room temperature (keep them in refrigerator).</p>
<p>15. Release parasites only after noticing adult moth catches in the pheromone trap or as per field observation.</p>	<p>Do not apply chemical pesticides within 7 days of release of parasites.</p>
<p>16. Apply Ha NPV, SI-NPV at recommended dose when large number of egg masses and early instar larvae are noticed. Apply NPV only in the evening hours after 5 pm.</p>	<p>Do not use the same sprayer for application of chemical pesticides and biocides.</p>
<p>17. In case of pests which are active during night like <i>Spodoptera</i> spray recommended biocides/chemicals at the time of their appearance in the night.</p>	<p>Do not apply pesticides/biocides when it is not absolutely required.</p>
<p>18. Spray pesticides thoroughly to treat the under surface of the leaves particularly for mites.</p>	<p>Do not spray pesticides at midday since, most of the insects are not active during this period.</p>

19. Apply short persistent pesticides to avoid pesticide residue in the soil and produce.

20. Follow the recommended procedure of trap technology in tomato and Cole crops.

Do not apply pesticides during preceding 7 days before harvest.

Do not apply long persistent pesticide on trap crop, otherwise it may not attract the pests and natural

SAFETY PARAMETERS IN PESTICIDES USAGE

S. No	Name of pesticide	Classification as per Insecticides Rules, 1971	Colour of Toxicity Triangle	WHO classification by hazard	First aid measures	Symptoms of poisoning	Treatment of poisoning	Waiting period (No. of days)
INSECTICIDES								
ORGANOCHLORINE PESTICIDES								
1.	Endosulfan	Highly toxic	Yellow	Class II – Moderately Hazardous	<p>Remove the person from the contaminated environment.</p> <p>In case of (a) Skin contact – Remove all contaminated clothings and immediately wash with lot of water and soap; (b) Eye contamination – Wash the eyes with plenty of cool and clean water; (c) Inhalation – Carry the person to the open fresh air, loosen the clothings around neck and chest, and (d) Ingestion – If the victim is fully conscious, induce vomiting by tickling back of the throat. Do not administer milk, alcohol and fatty substances. In case the person is unconscious make sure the breathing passage is kept clear without any obstruction. Victim's head should be little lowered and face should be turned to one side in the lying down position. In case of breathing difficulty, give mouth to mouth or mouth to nose breathing.</p> <p>Medical aid: Take the patient to the docotr/Primary Health Centre immediately along with the original container, leaflet and label.</p>	Nausea, vomiting, restlessness, tremor, apprehension, convulsions, coma, respiratory failure and death	<ul style="list-style-type: none"> - Gastric lavage with 2-4 L. tap water – Catharsis with 30 gm. (10 oz) sodium sulphate in one cup of water - Barbiturates in appropriate dosages repeated as necessary for restlessness or convulsions - Watch breathing closely, aspirate, oxygen and/or artificial respiration, if needed. - Avoid oils, oil laxatives and epinephrine (Adrenalin) – do not give stimulants. - Give calcium gluconate (10% in 10 ml. Ampules) intravenously every four hours. 	
ORGANOPHOSPHATE PESTICIDES								
2.	Phosalon	Highly toxic	Yellow	Class II – Moderately Hazardous		Mild – anorexia, headache, dizziness, weakness, anxiety, tremors of tongue and eyelids, miosis.	For extreme symptoms of O.P poisoning, injection of atropine (2-4 mg., for adults, 0.5-1.0 mg for children) is recommended.	

							Do not give atropine to a cyanotic patient. Give artificial respiration first then administer atropine.	I - SHUKENNA
FUNGICIDES								
3.	Sulphur	Slightly toxic	Green	Table 5 Unlikely to present acute hazard in normal use			Headache, palpitation, nausea, vomiting, flushed face, irritation of nose, throat eyes and skin etc.,	No specific antidote. Treatment is essentially symptomatic.
OTHERS								
4.	Fluchloralin	Moderately toxic	Blue	Class III Slightly hazardous			Headache, palpitation, nausea, vomiting, flushed face, irritation of nose, throat eyes and skin etc.,	No specific antidote. Treatment is essentially symptomatic
5.	Fenvalerate	Highly toxic	Yellow	Class II Moderately hazardous				
6.	Carbosulfan	Extremely toxic	Red	Class I b Highly hazardous				

AGRO ECOSYSTEM ANALYSIS IN OKRA

IPM is based on ecological interactions between the environment, plants, herbivores and their health. The health of the plant is determined by the environment and the herbivores. The herbivores are balanced by their natural enemies.

Agroecosystem analysis (AESA) is an approach, which can be gainfully employed by group of farmers for decision making in IPM. The basic components of AESA are:-

- (a) Plant health at different stages.
- (b) Compensation abilities of the plant.
- (c) Pest and defender population.
- (d) Soil condition, Irrigation status etc.
- (e) Weather conditions.
- (f) Past experience of the farmer in the situation.
- (g) Other investment opportunities.

The goal of the AESA activity is to analyze the field situation by observation; drawing and discussion. At the end of the activity the group should have made a decision about any action required for the field. The eco-system analysis is done weekly.

Time required - 120 minutes

Materials required-1 note book 1 large size paper, pencils and drawing crayons, polythene bags, plastic vials rubber band and chloroform.

Procedure

Go to the field, leave the border side, select 5th plant in a row for observation. Move 1-2 rows and select the fifth plant from that position for second observation. Likewise select a total of ten plants. Observe each plant on the following parameters and record.

Insect-pest, disease and defenders count : Count the larger pests and beneficial found on different branches and larvae of the plant. Start from the top of the plant and work downwards. Count the egg masses, larvae and adults of defoliators and workout percentage defoliation. Count the flower and fruit feeding insects and assess the percentage of fruits affected by them. Count the branches that are affected by die-back or other shoot disease.

Select three leaves from the sample plant, one taken from the top, one from the middle and one from the bottom of the plant. Pick or turn the leaf and count the number of sucking pests and predatory mites. Also record different leaf spot disease symptoms and count the number of spots. Estimate the percentage of leaf area affected.

Out of ten sample plants, assess the number of plants with virus symptoms. Like wise, for wilting symptoms. Pull wilting plants and observe symptoms on the roots (cut the roots to observe the colour of the vascular tissue).

Out of ten sample plants, note the no. of plants with flowers/fruits to assess the percentage of plants flowering/fruitletting.

Plant parameter - Measure the height of the plant, no. of leaves, width of foliage etc.

Walk through the whole plot to assess any other beneficial, pests or diseases, not observed on the ten sample plants, is occurring; note the uniformity in growth of the plant. Make records of the soil condition, water situation and presence of weeds (observe the different kinds of weeds and severity).

Rats:- Count number of plants affected by rats, also record number of live burrows in the plot.

Weather:- Record last week's weather parameters.

Find a shade to sit as a group. Each group should sit together in a circle with pencils, crayons and data from each of the field activities and the drawing of the AESA of the previous week.

Make a drawing on the large piece of paper for each plot observed. The rules for drawing are as follows:-

Draw the plant with the correct number of branches, leaves, flowers, fruits, write the plant height and number of green and yellow leaves some where. If the plant is healthy, colour the plant green. If disease occurs, draw the disease. If the plant is yellow, colour it yellow. For weeds, draw the approximate density and size of weeds in relation to the size of the vegetable plant. Draw the kinds of weed in the field.

For pest population draw the different insects found in the field on the right side of the plant, write the average number next to the insect. Also write the local name next to the insect. The data can also be summarized in a table on the right side.

For natural enemy populations, draw the insects and spiders as found in the field on the left side of the plant. Write the average number of natural enemies and their local names next to the drawing.

For rats write the average number of fruits/heads attacked.

If the week was mostly sunny draw a sun. If the week was mostly sunny and cloudy together draw a sun but half covered with dark clouds. If the week was cloudy all the day for most of the week put just dark clouds.

If the fertilizer was applied, then draw a picture of a hand throwing N, P or K depending on the type of fertilizer used.

If pesticides were used in the field, show sprays with a nozzle and write the type of chemical coming out of the nozzle. If granules were broadcast, show a hand with the name of pesticide being broadcast.

Now discuss the field situation among the group and arrive at a conclusion for the management practices required for the field. The discussion may be centered around as detailed below:

Steps	What to observe ?	What to ask ?	Actions
1	Weather	Rain/dry? Trends in weather dry/wet season, effects on crop	Do we need to water? Preparation of beds? Disease management.
2	Crop condition	Age of crop, seedling/Veg. Growth/flowering/fruitletting/colour of leaves, leaf spots? Die back? Diseases?	Need fertilizer? Need watering? Fungicides? Special topic?
3	Field condition	Soil structure hard/wet? Weeds present?	Hand weeding? Soil topic
4	Herbivores	What herbivores? Population? Damaging?	Insect zoo Special topic
5	Natural enemies (and neutrals)	What NE? Population? Damaging?	Insect zoo Special topic
6	Activities in neighbours fields	Do neighbours spray? What was sprayed?	Compare with FP Plot
7	Decisions made last week	Was the decision made effective? Do we need to do similar action?	Insect zoo Special topic

2. PIT FALL TRAP

Due to nocturnal behaviour and hiding during day time many insect pests like greasy cutworm, *Agrotis ypsilon* can't be observed on plant in the vegetable fields during day time. Like wise many ground dwelling predators like ground beetles which play major role in regulating the population of lepidopterus pest like *Helicoverpa armigera*, *Spodopetra litura* cannot be assessed properly by visual observations.

The pit fall trap method is effective in these situations. Pit fall trap is a simple plastic container (1 liter capacity) without lid. Container with half filled water with a few drops of teepol on water surface is buried in the field in such a way that the top surface is at level with that of soil surface. Observation should be made 48 to 72 hours after fixing the trap. Ten traps may be required per hectare of vegetable crop.

BASIC PRECAUTIONS IN PESTICIDES USAGE**A. Purchase**

1. Purchase only JUST required quantity e.g. 100, 250, 500 or 1000 or 1000 g/ml for single application in specified area.
2. Do not purchase leaking containers, loose, unsealed or torn bags.
3. Do not purchase pesticides without proper/approved LABELS.

B. Storage

1. Avoid storage of pesticides in house premises.
2. Keep only in original container with intact seal.
3. Do not transfer pesticides to other containers.
4. Never keep them together with food or feed/fodder.
5. Keep away from reach of children and livestock.
6. Do not expose to sun-light or rain water.
7. Do not store weedicides along with other pesticides.

C. Handling

1. Never carry/transport pesticides along with food materials
2. Avoid carrying bulk-pesticides (dusts/granules) on head, shoulders or on the back.

D. Precautions for Preparing Spray Solution

1. Use clean water
2. Always protect your NOSE, EYES, MOUTH, EARS and HANDS.
3. Use hand gloves, face mask and cover your head with cap.
4. Use polythin bags as hand gloves, handkerchiefs or piece of clean cloth as mask and a cap or towel to cover the head (Do not use polythin bag contaminated with pesticides).
5. Read the label on the container before preparing spray solution.
6. Prepare spray solution as per requirement on.
7. Do not mix granules with water.
8. Concentrated pesticides must not fall on hands etc. while opening sealed containers. Do not smell the pesticides.

9. Avoid spilling of pesticide solution while filling the sprayer tank.
10. Do not eat, drink, smoke or chew while preparing solution.
11. The operator should protect his bare feet and hands with polythin bags.

E. Equipment

1. Select right kind of equipment.
2. Do not use leaky, defective equipment.
3. Select right kind of nozzle.
4. Don't blow/clean clogged-nozzle with mouth. Use old tooth-brush tied with the sprayer and clean with water.
5. Do not use same sprayer for weedicide and insecticide.

F. Precautions for applying pesticides

1. Apply only at recommended dose and dilution.
2. Do not apply on hot sunny day or strong windy condition.
3. Do not apply just before the rains and also after the rains.
4. Do not apply against the windy direction.
5. Emulsifiable concentrate formulations should not be used for spraying with battery operated ULV sprayer.
6. Wash the sprayer and buckets etc. with soap water after spraying.
7. Containers, buckets etc. used for mixing pesticides should not be used for domestic purposes.
8. Avoid entry of animals and workers in the fields immediately after the spraying.

G. Disposal

1. Left over spray solution not be drained in ponds or water lines etc. Throw it in barren isolated area, if possible.
2. The used/empty containers should be crushed with a stone/stick and buried deep into soil away from water source.
3. Never re-use empty pesticide container for any purpose.

WARNINGS

- 1. Consult expert of the field of Plant Protection before use of any pesticide, regarding doses and time of application.**
- 2. Don't use pesticides in wrong way.**
- 3. Select pesticides wise fully.**
- 4. Use pesticides judiciously on need based**
- 5. Don't use green leaves as food after spray of pesticides.**