



IPM PACKAGE NO. 37



INTEGRATED PEST MANAGEMENT PACKAGE

FOR

LITCHI



Government of India
Ministry of Agriculture
Department of Agriculture & Cooperation
Directorate of Plant Protection, Quarantine & Storage
N. H. IV, Faridabad - 121 001.

IPM PACKAGE FOR LITCHI

CONTENTS

Subjects	Page No.
Foreward	i
Preface	ii
Acknowledgements	iii
I. MAJOR PESTS	
A. Pests of National significance	1
B. Pest of Regional Significance	1
II. PEST MONITORING	
A. Survey	2
B. Field scouting	2
C. Pest monitoring through traps	2
III. INTEGRATED PEST MANAGEMENT STRATEGIES	
A. Cultural Practices	3
B. Mechanical Practices	3
C. Biological Control Practices	3
D. Botanical pesticides	4
E. Chemical Control Practices	4
IV. SAFETY PARAMETERS	5
 Annexure I	 9



Government of India
Ministry of Agriculture
(Department of Agriculture & Cooperation)

DIRECTORATE OF PLANT PROTECTION, QUARANTINE & STORAGE
NH IV, FARIDABAD – 121 001 (Haryana)


DR.V. RAGUNATHAN
Plant Protection Adviser
To the Government of India

FOR E W A R D

Integrated Pest management (IPM) approach has been globally accepted for achieving sustainability in agriculture. It has become more relevant due to a number of advantages like safety to environment, pesticide-free food commodities, low input based Crop Production Programme etc. Though IPM approach has been taken up since 1981, its impact has not been felt until 1994. Human Resource Development has helped to sensitise extension functionaries and farmers about the usefulness of IPM.

For successful implementation of IPM, the scattered information on various components of this eco-friendly approach forms basic necessity. In this direction, initial attempts were made in 1992 to harmonise the IPM Package of Practices of various crops. Subsequently concerted efforts were made in 1998, 2001 and 2002 to update and develop IPM package of practices for agricultural and horticultural crops. Presently, IPM package of practices for 51 crops have been finalised to help the extension workers and farmers to manage the pests/ diseases and to minimise the over use/ misuse of chemical pesticides. Efforts have been made to incorporate the relevant available technical input provided by the scientists of ICAR Institutes/ SAUs and State Departments of Agriculture. However, suggestions for further improvement in future publication/ revision will be of immense help. Hopefully, these IPM Package of Practices will be useful for the Researchers, Plant Protection Workers and Farmers alike.

April 1, 2002


(V. Raguathan)

P R E F A C E

In order to minimise the indiscriminate and injudicious use of chemical pesticides, INTEGRATED PEST MANAGEMENT (IPM) has been enshrined as cardinal principle of Plant Protection in the overall Crop Protection Programme under the National Agricultural Policy of the Govt. of India. IPM is an eco-friendly approach for managing pest and disease problems encompassing available methods and techniques of pest control such as cultural, mechanical, biological and chemical in a compatible and scientific manner. The greater emphasis has been given on biological control including use of biopesticides.

With a view to provide technical knowledge to the extension functionaries and farmers in the States, first National Workshop on IPM for harmonisation of Package of Practices was organized at National Plant Protection Training Institute (NPPTI), Hyderabad during June 29-30, 1992. Subsequently workshops were organized from April 15-17, 1998 and Nov. 5-6, 1998 at Directorate of Plant Protection, Quarantine & Storage, Faridabad and IPM package of practices for 20 crops were evolved on rice, cotton, vegetables, pulses, and oilseeds. In this series, two National Workshops on IPM have been conducted at NPPTI, Hyderabad and Dte. of PPQ&S, Faridabad during May 14-17, 2001 and Feb. 20-22, 2002 respectively to update 20 available IPM Packages and develop 31 new IPM Packages specially for Horticultural crops. In these workshops, 51 IPM Package of Practices for cereal crops (Rice, Wheat, Maize, Sorghum, Millets), commercial crops (Cotton, Sugarcane, Tobacco, Tea), pulse crops (Pigeonpea, Gram, Black gram/Green gram, Pea, Rajma), oilseeds (Groundnut, Soybean, Rapeseed/Mustard, Sesame, Safflower, Castor, Sunflower, Oilpalm), vegetables (Potato, Onion, Tomato, Brinjal, Okra, Chillies, Cruciferous vegetables, Leguminous vegetables, Cucurbitaceous vegetables), fruit crops (Citrus, Banana, Apple, Mango, Guava, Grapes, Pineapple, Sapota, Pomegranate, Litchi), spice and plantation crops (Small Cardamom, Large Cardamom, Black Pepper, Ginger, Coriander, Cumin, Fennel, Coconut, Cashew and Arecanut) have been finalised.

IPM technology manages the pest population in such a manner that economic loss is avoided and adverse side effects of chemical pesticides are minimized. The IPM packages encompasses various management strategies for containing the pest and disease problems. Pest monitoring is also one of the important component of IPM to take proper decision to manage any pest problem. It can be done through Agro-Ecosystem Analysis (AESA), field scouting, light, pheromone, sticky/yellow pan traps. The economic threshold levels (ETL) of important pests and diseases are also given in the packages to take appropriate control measures when pest population crosses ETL.

These IPM packages developed with the technical inputs from experts from Indian Council of Agriculture Research, State Agricultural Universities, Central Directorate of Plant Protection, Pesticide Industries and State Departments of Agriculture/Horticulture will provide technical backup in the management of pests, diseases, weeds, nematodes and rodents in the Indian Agriculture and Horticulture. These will also be useful in reducing the pesticide residues in exportable agricultural commodities and would also help in the management of pests/diseases/weeds/nematodes which may get inadvertently introduced in the country.

IPM Package of Practices for Agricultural and Horticultural crops will be helpful to minimize the ill effects of chemical pesticides to promote the IPM for sustainable production. These packages will be useful for the researchers, extension workers and farmers alike who are engaged in the agricultural practices.

April 1, 2002



(A.D. Pawar)
Director (IPM)

ACKNOWLEDGEMENTS

The IPM Package of Practices for **Litchi** crop was discussed and finalised in the National Workshop on IPM held at National Plant Protection Training Institute (NPPTI), Hyderabad during May 14-17, 2001. The technical input received from the following experts is thankfully acknowledged.

- I. Chairman, Technical Session : Dr. AD Pawar, Director (IPM), Dte of PPQS, Faridabad
- II. Coordinator, Technical Session : Dr. M.P. Misra, Dy. Director (Ent.), Dte. of PPQS, Faridabad
- III. Co-chairman, Technical Session : Dr. R.P. Shukla, Principal Scientist (Ent.) CISTH (ICAR), Lucknow.
- IV. Expert input : 1. Dr. S.D. Sawant, Sr. Scientist (Ent.), NRC for Grapes (ICAR), Pune.
2. Dr. T. Manoharan, Associate Professor (Ento.), TNAU, Coimbatore.
3. Dr. G. Ramasubha Reddy, APPO, NPPTI, Hyderabad.
4. Dr. J.N. Thakur, AD(E), Central IPM Centre, Solan
5. Sh. N. M. Dev., AD(E), Central IPM Centre, Jalandhar.
6. Dr. R.P. Misra, PPO(E), Central IPM Centre, Lucknow.
7. Sh. O.P. Verma, PPO(PP), Central IPM Centre, Jammu.
8. Dr. V. K. Srivastava, PPO(PP), Central IPM Centre, Nagpur
9. Sh. M.K. Das, APPO, Central IPM Centre, Burdwan.
10. Sh. Ashok Shukla, PPO(PP), Central IPM Centre, Baroda
11. Dr. Sandeep Pandey, PPO(PP), Central IPM Centre, Lucknow.
12. Sh. S.K. Verma, PPO(E), Central IPM Centre, Patna.
13. Sh. M. Parray, SSA, Central IPM Centre, Srinagar (J&K)
14. Sh. Padam Singh, SSA, Central IPM Centre, Dimapur.
15. Dr. D. Kanungo, Jt. Director, CIL, Dte. of PPQ&S, Faridabad
16. Dr. R.B.L. Bhaskar, Jt. Director (Bio.), Dte of PPQS, Faridabad
17. Dr. Brajendra Singh, Entomologist, Dte of PPQS, Faridabad
18. Dr. B.S. Fogat, Agronomist (H), Dte of PPQS, Faridabad
19. Sh. K.S. Gautam, Dy. Director (PP), Dte. of PPQ&S, Faridabad.
20. Sh. S.P. Kulshrestha, Dy. Director (PP), Dte. of PPQ&S, Faridabad.
21. Dr. Jasvir Singh, Asstt. Director (E), Dte. of PPQ&S, Faridabad.
- V. Technical input:
1. Sh. S.P. Singh, APPO, IPM Div., Dte of PPQS, Faridabad.
 2. Sh. K.S. Sharma, SSA, IPM Div., Dte of PPQS, Faridabad.
 3. Sh. Yogesh Kumar, SSA, IPM Div., Dte of PPQS, Faridabad.
 4. Sh. R.S. Tomer, SSA, IPM Div., Dte of PPQS, Faridabad.
 5. Sh. Laxmi Chand, SSA, IPM Div., Dte of PPQS, Faridabad.
 6. Mohd. Abrar Alam, Stenographer, IPM Div., Dte of PPQS, Faridabad.
 7. Sh. N.K. Mishra, LDC, IPM Div., Dte of PPQS, Faridabad.

IPM PACKAGE OF PRACTICES FOR LITCHI

I. MAJOR PESTS:

A. Pests of National Significance

1. Insect Pests

- 1.1 Fruit & Stone borer
- 1.2 Litchi mite
- 1.3 Leaf folder
- 1.4 Mealy bug

2. Diseases

- 2.1 Fruit rot
- 2.2 Powdery mildew

3. Weeds

- 3.1 *Cyperus rotundus*
- 3.2 *Digitaria sanguinalis*
- 3.3 *Setaria glauca*
- 3.4 *Cleome vis cosa*
- 3.5 *Tridox procumbesn*
- 3.6 *Cynodon dactylon*

B. Pests of Regional Significance

1. Insect Pest

- 1.1 Bark eating caterpillar
- 1.2 Leaf Miner
- 1.3 Whitefly
- 1.4 Shoot borer

2. Diseases

2.1 Dieback

3. Weeds

3.1 *Panicum repens*

3.2 *Trainthema monogynae*

II. PEST MONITORING

A. Survey:

The objective of survey through roving surveys is to monitor the initial development of pest and disease in the endemic areas. Therefore, for field scouting farmers should be mobilized to observe the pest and disease occurrence at the intervals as stipulated under different developmental stages. The plant protection measures are required to be taken only when biocontrol potential does not show promise and pest and diseases incidences shows increasing trend.

B. Field Scouting:

Field scouting for pests/diseases and biocontrol fauna/flora by extension agencies and farmers once in a fortnight should be undertaken to assess increasing/decreasing trend in the pest/disease incidence and availability of biocontrol potential. This should be done soon after the appearance of new flush after the fall of old leaves as such stage of the crop having succulent tissues is variable to attack by pests and diseases. The state Departments of Horticulture should make all possible efforts by using different media, mode and publicity to inform the farmers for field scouting in the specific crop area having indication of pest and disease build up.

C. Pest Monitoring through Traps:

1. Through yellow sticky traps: Setup yellow fast coloured sticky traps for monitoring sucking one trap/ 5 trees. Locally available empty yellow palmolive-tin coated with grease / vaseline / castor oil on outer surface may also be used.

2. Through pheromone traps: Certain pests of fruit crops required installation of pheromone traps to monitor initial pest build up and suppression of its increasing population. Sticky pheromone traps may also be used 5-7 traps per ha. for effective monitoring.

III. IPM STRATEGIES

A. Cultural practices

1. Repeatedly ploughing around the tree and rap alkathene bands round the trunk during November –December to prevent climbing mealy bug. Apply Neem Seed Cake after ploughing @ 4kg/tree.
2. Repeated hoeing / weeding around the tree basin.

B. Mechanical control practices

1. After mud plastering 25 cm wide, 400 gauge alkathene (Polythene) sheet should be fastened to the tree trunk with the help of sulti about 30 cm above the ground level to prevent crawlers to climbing of freshly hatched first instar nymphs of mealy bug in the month of November – December.
2. The mite affected leaves and twigs should be cut and burnt in the month of May and October.
3. Cutting and destroying branches 10 cm below from the dried portion and pasting of copper oxychloride for the control of die back disease and powdery mildew.

C. Biological control practices

1. Conservation:

Number of effective parasites, predators and pathogens are very active against pests of Litchi. Spider, *Sumnius renardi*, *Coccinellids*, *Chrysoperla lacciperda* could be conserved by using various conservation method.

D. Botanical pesticides

1. NSKE @ 5% helps in reducing the pest population.

E. Chemical control practices

1. Spray *endosulphan 35% EC @ 2.0 ml/lit of water just before flowering against fruit and stone borer.
2. *Dicofol 10.5% (3.0ml/liter of water) or dimethoate 30%EC should be applied twice against Litchi mite at 10 days interval.
3. Remove all frass and seal the holes with cotton swab soaked in petrol or chloroform or *dichlorvos (0.1%) against bark eating caterpillars.
4. Spray *quinalphos (0.05%) or carbaryl (2g/lit. of water) for control of fruit borer and leaf miner.
5. Alternate the foliar spray with *wetttable sulphur (0.2%) or *triadomorph (0.1%) or dinocap (0.1%) at 15 days interval to control powdery mildew.

*Not as per the approved usage under Insecticide Act, 1968

IV. SAFETY PARAMETERS IN PESTICIDES USAGE

S. No	Name of pesticide	Classification as per Insecticides Rules, 1971	Colour of Toxicity Triangle	WHO classification by hazard	First aid measures	Symptoms of poisoning	Treatment of poisoning	Waiting period (No. of days)
INSECTICIDES								
ORGANOCHLORINE PESTICIDES								
1.	Endosulfan	Highly toxic	Yellow	Class II – Moderately Hazardous	Remove the person from the contaminated environment.	Nausea, vomiting, restlessness, tremor, apprehension, convulsions, coma, respiratory failure and death	<ul style="list-style-type: none"> - Gastric lavage with 2-4 L. tap water – Catharsis with 30 gm. (10 oz) sodium sulphate in one cup of water - Barbiturates in appropriate dosages repeated as necessary for restlessness or convulsions - Watch breathing closely, aspirate, oxygen and/or artificial respiration, if needed. - Avoid oils, oil laxatives and epinephrine (Adrenalin) – do not give stimulants. - Give calcium gluconate (10% in 10 ml. Ampules) intravenously every four hours. 	
2.	Dicofol	Moderately toxic	Blue	Class III – Slightly hazardous	<p>Remove all contaminated clothings and immediately wash with lot of water and soap; (b) Eye contamination – Wash the eyes with plenty of cool and clean water; (c) Inhalation – Carry the person to the open fresh air, loosen the clothings around neck and chest, and (d) Ingestion – If the victim is fully conscious, induce vomiting by tickling back of the throat. Do not administer milk, alcohol and fatty substances. In case the person is unconscious make sure the breathing passage is kept clear without any obstruction. Victim's head should be little lowered and face should be turned to one side in the lying down position. In case of breathing difficulty, give mouth to mouth or mouth to nose breathing.</p> <p>Medical aid: Take the patient to the docotr/Primary Health Centre immediately along with the original container, leaflet and label.</p>			

ORGANOPHOSPHATE PESTICIDES

3.	Quinalphos	Highly toxic	Yellow	Class II - Moderately Hazardous		Mild - anorexia, headache, dizziness, weakness, anxiety, tremors of tongue and eyelids, miosis, impairment of visual acuity.	For extreme symptoms of O.P poisoning, injection of atropine (2-4 mg., for adults, 0.5-1.0 mg for children) is recommended, repeated at 5-10 minute intervals until signs of atropinization occur.
4.	Dichlorvos	Extremely toxic	Bright red	Class I b - Highly hazardous		Moderate- nausea, salivation, lacrimation, abdominal cramp, vomiting, sweating, slow pulse, muscular tremors, miosis. Severe - diarrhoea, pinpoint and non-reactive pupils, respiratory difficulty, pulmonary edema, cyanosis, loss of sphincter control, convulsions, coma and heart block.	Speed is imperative - Atropine injection - 1 to 4 mg. Repeat 2 mg. when toxic symptoms begin to recur (15-16 minute intervals), Excessive salivation - good sign, more atropine needed; - Keep airways open, Aspirate, use oxygen, insert endotracheal tube. Do tracheotomy and give artificial respiration as needed. - For ingestion lavage stomach with 5% sodium bicarbonate, if not vomiting. For skin contact, wash with soap and water (eyes- wash with isotonic saline). Wear rubber gloves while washing contact areas.
5.	Diazinon	Moderately toxic	Yellow	Class III			
6.	Fenitrothion	Highly toxic	Yellow	Class II			In addition to atropine give 2-PAM (2-pyridine aldoxime methiodide). 1 g and 0.25 g for infants

							<p>intravenously at a slow rate over a period of 5 minutes and administer again periodically as indicated. More than one injection may be required.</p> <p>Avoid morphine, theophyllin, aminophyllin, barbiturates of phenothiazines.</p> <p>Do not give atropine to a cyanotic patient. Give artificial respiration first then administer atropine.</p>
--	--	--	--	--	--	--	--

CARBAMATES

5.	Carbaryl	Highly toxic	Yellow	Class II – Moderately hazardous		<p>Constriction of pupils, salivation, profuse sweating, lassitude, muscle incoordination, nausea, vomiting, diarrhoea, epigastric pain, tightness in chest.</p>	<ul style="list-style-type: none"> - Atropine injection 1 to 4 mg. Repeat 2 mg when toxic symptoms begin to recur (15-60 minute intervals). Excessive salivation – good sign, more atropine needed. - Keep airway open. Aspirate, use oxygen, insert endotracheal tube. Do tracheotomy and give artificial respiration as needed. - For ingestion, lavage stomach with 5% sodium bicarbonate, if not vomiting. For skin contact was with soap and water (eyes – wash with isotonic saline). Wear rubber gloves while washing contact
----	----------	--------------	--------	---------------------------------	--	--	---

							<p>area.</p> <ul style="list-style-type: none"> - Oxygen - Morphine, if needed. <p>Avoid theophyllin and aminophyllin or barbiturates.</p> <p>2-PAM and other oximes are not harmful and in fact contra indicated for routine usatge.</p> <p>Do not give atropine to a cyanotic patient. Give artificial respiration first then administer atropine.</p>
--	--	--	--	--	--	--	--

FUNGICIDES

6.	Wettable sulphur	Slightly toxic	Green	Table 5 – Unlikely to present acute hazard in normal use.		Headache, palpitation, nausea, vomiting, flushed face, irritation of nose, throat eyes and skin etc.,	No specific antidote. Treatment is essentially symptomatic.
7.	Tridemorph			Class II – Moderatley toxic			
8.	Dinocap	Moderately toxic	Blue	Class III – Slightly hazardous			

OTHERS

9.	Kelthane					Headache, palpitation, nausea, vomiting, flushed face, irritation of nose, throat eyes and skin etc.,	No specific antidote. Treatment is essentially symptomatic
----	----------	--	--	--	--	---	--

BASIC PRECAUTIONS IN PESTICIDE USAGE

A. Purchase:

1. Purchase only JUST required quantity e.g. 100,250,500 or 1000 g/ml for single application in specified area.
2. Do not purchase leaking containers, loose, unsealed or torn bags.
3. Do not purchase pesticides without proper/ approved LABELS.

B. Storage:

1. Avoid storage of pesticides in the house premises.
2. Keep only in original container with intact seal.
3. Do not transfer pesticides to other container.
4. Never keep them together with food or feed/ fodder.
5. Keep away from the reach of children and livestock.
6. Do not expose to sun-light or rain water.
7. Do not store weedicides along with other pesticides.

C. Handling:

1. Never carry/ transport pesticides along with food materials.
2. Avoid carrying bulk - pesticides (dusts / granules) on head, shoulders or on the back.

D. Precautions for Preparing Spray Solution:

1. Use clean water.
2. Always protect your NOSE, EYES, MOUTH, EARS and HANDS.
3. Use hand gloves, face mask and cover your head with cap.
4. Use polyethylene bags as hand gloves, handkerchiefs or piece of clean cloth as mask and a cap or towel to cover the head (Do not use polyethylene bag contaminated with pesticides).
5. Read the label on the container before preparing spray solution.
6. Prepare spray solution as per requirement.
7. Do not mix granules with water.
8. Concentrated pesticides must not fall on hands etc. while opening sealed containers. Do not smell the sprayer tank.
9. Avoid spilling of pesticide solution while filling the sprayer tank.
10. Do not eat, drink, smoke or chew while preparing solution.
11. The operator should protect his bare feet and hands with polyethylene bags.

E. Equipment:

1. Select right kind of equipment.
2. Do not use leaky, defective equipment.
3. Select right kind of nozzle.
4. Don't blow/clean clogged- nozzle with mouth. Use old tooth- brush tied with the sprayer and clean with water.
5. Do not use some sprayer for weedicide and insecticide.

F. Precautions for applying pesticides:

1. Apply only at recommended dose and dilution.
2. Do not apply on hot sunny day or strong windy condition.
3. Do not apply just before the rains and also after the rains.
4. Do not apply against the wind direction.
5. Emulsifiable concentrate formulations should not be used for spraying with battery operated ULV sprayer.
6. Wash the sprayer and bucket etc with soap water after spraying.
7. Containers, buckets etc. used for mixing pesticides should not be used for domestic purposes.

8. Avoid entry of animals and workers in the fields immediately after the spraying.

G. Disposal:

1. Left over spray solution should not be drained in ponds or water lines etc. Throw it in barren isolated area, if possible.
2. The used/ empty containers should be crushed with a stone / stick and buried deep into soil away from water source.
3. Never re-use empty pesticide container for any purpose.
