

E-mail: directorcpps@tnau.ac.in  
kramaraju60@gmail.com

Phone : 0422-6611237  
Tele - Fax: 0422-6611437  
Mobile: 94420 06138



**TAMIL NADU AGRICULTURAL UNIVERSITY  
CENTRE FOR PLANT PROTECTION STUDIES**

Dr. K.RAMARAJU, Ph.D.  
DIRECTOR i/c

Coimbatore - 641 003  
Tamil Nadu, India

---

**PEST AND DISEASE FORECASTING FOR THE MONTH OF NOVEMBER 2013**

Based on the pest and disease surveillance reports from different districts of Tamil Nadu the following forecast is made.

**Paddy**

In Coimbatore, Erode, Nagapattinam, Sivaganga, Theni, Thiruvarur and Thirunelveli districts, leaf folder and stemborer incidence were found. Hence, farmers are advised to spray neem seed kernel extract 5% (NSKE) or flubendiamide 20 WG 125 g / ha. Set up light traps to monitor the pest population.

Bacterial leaf blight incidence was found in Chingulpet, Kanjeeपुरam and Thiruvarur district. Hence, farmers can take proper plant protection measures and spray copper hydroxide @ 2.5g/litre. There is a possibility for the outbreak of paddy blast due to cloudy weather prevailing in coastal areas hence farmers are advised to spray tricyclozole 0.1 per cent immediately. In delta areas, spray phosalone 35 EC @ 1500 ml/ha for the management of mealy bugs in paddy.

**Cotton**

In Coimbatore, Dharmapuri and Theni sucking pests *viz.*, jassids and thrips were noticed. Hence, the farmers are advised the setup the yellow sticky traps to monitor the pest population. If necessary, farmers can spray NSKE 5%.

**Turmeric**

In Erode district, leaf spot of turmeric was noticed. Rainfall and low temperature prevailing during night favours the incidence and spread of the disease. Hence, farmers are advised to take up spraying of carbendazim 1 g/lit.

## **Sugarcane**

In Erode, Vellore, Cuddalore, Thiruvannamalai, Nagapattinam, Sivaganga, Theni, Thiruvarur and Thirunelveli districts borer pest complex was found below ETL level. Hence, the farmers are advised to release the egg parasitoid *Trichogramma* @ 6cc/acre for 6 releases at the rate of 1 cc/release starting from 45<sup>th</sup> day onwards at 15 days interval.

## **Groundnut**

In Erode, Salem, Virudachalam, Thiruvannamalai and Nagapattinam districts, leaf miner incidence was recorded. In addition, Tikka leaf spot was recorded and for managing this disease, spray carbendazim 1g/litre two times at 10 days interval can be recommended. Farmers are advised to monitor leaf miner adults using light traps and if needed farmers can spray neem seed kernel extract at 5%.

## **Coconut**

In Udumelpet and Pollachi areas leaf blight incidence was noticed. Hence, farmers are advised to remove the affected fronds and root feed propiconazole 2 ml / 100 ml of water per tree.

This is for favour of kind information. Necessary control measures may please be adopted.

**DIRECTOR (CPPS) i/c**