

#### Gramin Krishi Mausam Sewa (GKMS)

# for the Thiruchirapalli District

# Forecast Period: 15<sup>th</sup> JUNE to 19<sup>th</sup> JUNE, 2016

Tamil Nadu Rice Research Institute, Aduthurai- 612 101 Phone: 0435 2472098 E-mail: dirtrri@tnau.ac.in Fax: 0435 2472881



Date: 14.06.16

## Weather for the past four days (10.06.2016–13.06.2016)

Parameters	Past	
	weather	
Rainfall (mm)	0.4	
Maximum temperature (°C)	35 - 37	
Minimum temperature (°C)	26 - 28	
Cloud cover (Octa)	4 – 6	
Relative humidity (%)	40 - 73	
Wind speed (km/ hr)	13 - 24	

### Weather forecast for the next five days (until 08:30 hrs of 19.06.2016)

Parameters	Ensemble FCST					
	Day 1 15.06.2016	Day 2 16.06.2016	Day 3 17.06.2016	Day 4 18.06.2016	Day 5 19.06.2016	
Rainfall (mm)	2	0	0	0	0	
Maximum temperature (°C)	37	37	37	37	37	
Minimum temperature (°C)	27	27	27	27	27	
Cloud cover (Octa)	6	6	6	6	6	
Relative humidity (%)	49 – 59	49 – 59	49 – 59	49 – 59	49 – 59	
Wind speed (kmph)	17	17	17	17	17	
Wind direction	South West					

The sky will be mostly cloudy. A light rainfall 2 mm is expected on 15.06.2016.

Weather based Agro-Advisories

Crop: Rice, Black gram, Sesame, Sugarcane, Cotton

**Stage:** Rice fallow cotton – Flowering stage; Rice: Nursery preparation

- For SRI cultivation, prepare 2.5 cent raised bed nursery for 1 ha planting. Broadcast uniformly 5 kg of pre germinated seeds over it. Sprinkle water in the bed during morning and evening using rose cane. Maintain standing water in the channel around the beds from 6<sup>th</sup> day of sowing. Sprinkle 0.5 % urea solution on 9<sup>th</sup> day after sowing. Stop irrigation two days before pulling out of seedlings. The seedlings will be ready at 14 days after sowing.
- For kuruvai nursery, an area of 20 cents can be allotted for planting 1 ha with the application of FYM or compost @ 1 ton. Before the last puddling, apply 40 kg DAP. For clay soil, 4 kg gypsum and 1 kg DAP/cent can be applied 10 days prior to pulling out of seedlings to avoid root snapping.
- ➤ Occurrence of thrips is being noticed in the Kuruvai paddy nursery due to prevailing hot weather condition. To control spray chloripyriphos 2.5 ml/lit of water.