

M.S. Swaminathan Research Foundation, Kannivadi AgroMet Advisory Bulletin (AAB) for PUDUKOTTAI District precast Period: 16th November 20th November 201

<u>Forecast Period: 16th November – 20th November 2016</u>

ovember 2016 es.in Fax: 044-

Phone no. 044-22541229 Email: rengalakshmi@mssrf.res.in Fax: 044-22541319

Cianificant next much on from 11

Bulletin No: MSSRF/DDL/092(11)/2016

Date: 15.11.2016

Significant past weather from 11-11-2016 to 14-11-2016				
Parameters	Past weather			
Rainfall (mm)	2.2			
Maximum Temperature (°C)	32-33			
Minimum Temperature (°C)	23-26			
Sky condition (Octas)	1-6			
Relative humidity (%)	46-78			
Wind speed (kmph)	10-18			

Ensemble Weather forecast until 08:30 hrs of 20.11.16							
Parameters	Ensemble FCST						
	16.11.16	17.11.16	18.11.16	19.11.16	20.11.16		
	Wednesday	Thursday	Friday	Saturday	Sunday		
Rainfall (mm)	0.5	1	1	1	0		
Maximum Temperature (°C)	33	33	33	33	33		
Minimum Temperature (°C)	24	24	24	24	24		
Sky condition (Octas)	4	4	4	4	4		
Max. Relative humidity (%)	78	78	78	78	78		
Min. Relative humidity (%)	60	60	60	60	60		
Wind speed (kmph)	16	16	16	16	16		
Wind direction	East	East	East	East	East		

SMS Advisory:

Light rain may occur during next four days (16^{th} Nov – 19^{th} Nov) over certain spots

Weather Based Agro-Advisories :

Imp. Crops: Rice and Sugarcane

- To control the brown spot disease in paddy, it is recommended to spray 5gm of pseudomonus per litre of water or 2.5gms of dithane M45 per 10 litres of water.
- Due to this humid condition, incidence of internode borer may affect the sugarcane crops, to control this farmers are advised to tie 3 cc of Trichogramma parasitoids per acre in their field.
- Leaf mite incidence is being noticed in the paddy fields due to prevailing weather condition. To manage this, spray Dicofol @ 500 ml/ acre or wettable sulphur @ 400 g/ acre with 200 liter of water.
- Stem borer incidence is noticed in paddy crop. Hence farmers are advised to spray any one of the following Quinalphos 25 EC1000 ml/ha (or) Phosphamidon 40 SL 600 ml/ha (or) Profenophos 50 EC 1000 ml/ha.
- Due to moderate humid conditions mastitis disease may affect the milch animals. Clean the udder with 1% potassium permanganate solution, before and after milking.