

#### Gramin Krishi Mausam Sewa (GKMS) for Thanjavur District Forecast Period: 07<sup>rd</sup> SEPTEMBER to 11<sup>h</sup> SEPTEMBER, 2016 Tamil Nadu Rice Research Institute, Aduthurai- 612 101



Tamil Nadu Rice Research Institute, Aduthurai- 612 101 Phone: 0435 2472098 E-mail: <u>dirtrri@tnau.ac.in</u> Fax: 0435 2472881

#### Date: 06.09.16

## Weather for the past three days (02.09.2016–05.09.2016)

Parameters	Past	
	weather	
Rainfall (mm)	15.0	
Maximum temperature (°C)	34-35	
Minimum temperature (°C)	26-27	
Cloud cover (Octa)	4-5	
Relative humidity (%)	72-76	
Wind speed (km/ hr)	4-5	

### Weather forecast for the next five days (until 08:30 hrs of 11.09.2016)

Parameters	Ensemble FCST				
	Day 1 07.09.2016	Day 2 08.09.2016	Day 3 09.09.2016	Day 4 10.09.2016	Day 5 11.09.2016
Rainfall (mm)	2	0	0	0	3
Maximum temperature (°C)	35	35	35	34	34
Minimum temperature (°C)	27	27	27	26	26
Cloud cover (Octa)	5	5	5	5	5
Relative humidity (%)	62-72	62-72	62-72	66-74	66-74
Wind speed (kmph)	5	5	5	5	5
Wind direction	North West				

The sky will be mostly cloudy. A light rainfall 2 - 3 mm is expected on next two days [07.92016 & 11.09.2016].

### Weather based Agro-Advisories

# Crop: Rice, Black gram, Sugarcane, Cotton Stage: Kuruvai – Flowering & Maturity stage; Samba: Land preparation and sowing

- Field preparation has to be done for samba season paddy crop by utilizing the last week rainfall.
- For direct sown rice in samba season, sowing by using tractor drawn seed drill, seed rate of 15 kg/acre is required. For manual broadcasting seed rate of 30 kg/acre is required. Seeds should be treated with Pseudomonas fluorescense 10g/kg of seed before sowing. Seed treatment increase seed germination, seedling vigour and controls the seed borne disease.
- For 1 acre Samba crop, apply 200 kg of gypsum, 10 kg of zinc sulphate, 15 kg of urea, 40 kg DAP and 10 kg of potash as basal fertilizers. Apply 100 kg of neem cake before transplanting to avoid insect and nematode attack.
- In direct sown wet seeded rice apply pre-emergence herbicide pretilachlor + safener @500 ml/acre on 3-4 days after sowing to control weeds.