



**Gramin Krishi Mausam Sewa (GKMS)
for the Karur District**

Forecast Period: 26th APRIL to 30th APRIL, 2017
Tamil Nadu Rice Research Institute, Aduthurai- 612 101
Phone: 0435 2472098 E-mail: dirtrri@tnau.ac.in Fax: 0435 2472881



Date: 25.04.2017

Weather for the past four days (21.04.2017–24.04.2017)

Parameters	Past weather
Rainfall (mm)	0.0
Maximum temperature (°C)	40 – 41
Minimum temperature (°C)	28 – 29
Cloud cover (Octa)	4 – 5
Relative humidity (%)	31 – 60
Wind speed (km/ hr)	7 – 12

Weather forecast for the next five days (until 08:30 hrs of 29.04.2017)

Parameters	Ensemble FCST				
	Day 1 26.04.2017	Day 2 27.04.2017	Day 3 28.04.2017	Day 4 29.04.2017	Day 5 30.04.2017
Rainfall (mm)	5	5	0	0	0
Maximum temperature (°C)	41	41	41	40	40
Minimum temperature (°C)	28	28	28	29	29
Cloud cover (Octa)	4	4	4	4	4
Relative humidity (%)	35 – 55	35 – 55	35 – 55	35 – 55	35 – 55
Wind speed (kmph)	8	8	8	10	10
Wind direction	North	North	North	North	North

The sky will be partly cloudy. A light rainfall 5 mm is expected on next two days.

Weather based Agro-Advisories

Crop: Rice and Irrigated blackgram

- Leaf mite incidence is being noticed in the summer paddy nursery due to prevailing hot weather condition. To manage this, spray Dicofol @ 600 ml / acre or wetttable sulphur @ 400 g / acre with 200 litres of water.
- Occurrence of thrips is being noticed in the summer paddy nursery due to prevailing hot weather condition. To control this, spray chloripyriphos 2.5 ml / lit of water.
- In irrigated blackgram, occurrence of white flies is being noticed due to the prevailing hot weather. To control this spray azadirachtin 0.03% (3 ml / litre of water) along with sticking agent in the morning or evening using hand sprayer, If damage is more spray imidacloprid 200 SL @ 40ml / acre with 200 litres of water.
- In irrigated blackgram, application of pre-emergence herbicide pendimethalin @ 1.3 litre / acre is to be done on 3 DAS.

Livestocks

- Poultrys should be vaccinated against Ranikhet disease.
- Sprinkling of water on mulching cows in every two hours between 10 a.m. to 5 p.m. to sustain the milk production.

Director