



**“Gramin Krishi Mausam Sewa” (GKMS) -Chennai**  
(Issued jointly by Tamil Nadu Veterinary and  
Animal Sciences University,  
& India Meteorological Department)



**Weather Advisory Bulletin - Vellore District**  
(Period 14<sup>th</sup> to 18<sup>th</sup> Apr. 2017)

**A. Weather for the past two days (11.04.2017 and 12.04.2017)**

During past two days around Vellore area, the sky was clear to mostly clear without rainfall. Maximum temperature was 40.0-41.0°C and Minimum temperature was 25.0-26.0°C. Wind speed was 4-6 km per hour. Relative humidity was recorded between 36-81%.

**B. Weather forecast for next five days (Until 08.30 hrs of 18.04.2017)**

Parameters	14-Apr-17	15-Apr-17	16-Apr-17	17-Apr-17	18-Apr-17
Cloud Cover	Clear Sky	Clear Sky	Clear Sky	Clear Sky	Clear Sky
<b>Rainfall (mm)</b>	0	0	0	0	0
Wind Speed Kmph	4	4	4	4	4
Wind Direction	East	East	East	South	South
Temperature °C					
Maximum	40	40	40	41	41
Minimum	25	25	25	25	25
Relative Humidity %	35-80	35-80	35-80	35-80	35-80

**Agrometeorological Advisories**

**Livestock:**

1. Feeders should be raised in poultry shed between 10 am to 5 pm; provide mineral mixture and B-complex vitamins in drinking water to avoid heat stroke.
2. During hot part of the day dairy cows should be showered with water to reduce heat stress and to sustain milk production.
3. Due to increase in day temperature animals should not be allowed for grazing during hot part of the day, provide cool drinking water and shade to the animals to reduce heat stress.
4. Farmers are advised to feed 100 to 200 grams of concentrate per day for sheep and goats due to non availability of sufficient pasture in the grazing land.
5. Apply Butox solution (2-3ml in one litter water) externally to remove ecto-parasites in sheep & goats.

**Agriculture**

**Imp. Crops: Rice, Sugarcane and Groundnut**

1. To control shoot and fruit borer in Brinjal, apply Emamectin benzoate 5 SG @ 200 g/ha
2. Two percent of DAP spray (Dissolve 4.5 kg of DAP in 10 litres of water previous day night and the supernatant liquid shall be mixed with 190 litres of water) may be done at the time of flowering and another dose fifteen days after the first spray for rice fallow pulses.
3. To control powdery mildew in rice fallow pulses, spray 5% Neem Seed Kernel Extract or Mancozeb @ 400 g/ac or Wettable Sulphur @ 1 kg/ac during morning or evening hours.