



“Gramin Krishi Mausam Sewa” (GKMS) –Chennai
(Issued jointly by Tamil Nadu Veterinary and
Animal Sciences University,
& India Meteorological Department)



Weather Advisory Bulletin – Villupuram District
(Period 26th to 30th Apr. 2017)

A. Weather for the past four days (21.04.2017 to 24.04.2017)

During past four days around Villupuram district, the sky was clear to mostly clear without rainfall. Maximum temperature was 35.0-38.0°C and Minimum temperature was 28.0-29.0°C. Wind speed was 6-11 km per hour. Relative humidity was recorded between 67-80%.

B. Weather forecast for next five days (Until 08.30 hrs of 30.04.2017)

Parameters	26-Apr-17	27-Apr-17	28-Apr-17	29-Apr-17	30-Apr-17
Cloud Cover	Clear Sky	Clear Sky	Clear Sky	Clear Sky	Clear Sky
Rainfall (mm)	3	3	0	0	0
Wind Speed Kmph	10	10	10	10	10
Wind Direction	South East	South East	South East	East South East	East South East
Temperature °C					
Maximum`	37	37	37	38	38
Minimum	28	28	28	29	29
Relative Humidity %	65-73	65-73	65-73	65-67	65-67

Agrometeorological Advisories

Livestock:

1. Light rainfall may be expected for next two days.
2. Coconut leaves or paddy straws may be spread over the roof of the animal shed and spray water to avoid heat stress to the animals.
3. Feeders should be raised in poultry shed between 10 am to 5 pm; provide mineral mixture and B-complex vitamins in drinking water to avoid heat stroke.
4. During hot part of the day dairy cows should be showered with water to reduce heat stress and to sustain milk production.
5. Due to increase in day temperature animals should not be allowed for grazing during hot part of the day, provide cool drinking water and shade to the animals to reduce heat stress.
6. Farmers are advised to feed 100 to 200 grams of concentrate per day for sheep and goats due to non availability of sufficient pasture in the grazing land.
7. Apply Butox solution (2-3ml in one litter water) externally to remove ecto-parasites in sheep & goats.

Agriculture

Imp. Crops: Rice, Sugarcane and Groundnut

1. To control shoot and fruit borer in Brinjal, apply Emamectin benzoate 5 SG @ 200 g/ha
2. Two percent of DAP spray (Dissolve 4.5 kg of DAP in 10 litres of water previous day night and the supernatant liquid shall be mixed with 190 litres of water) may be done at the time of flowering and another dose fifteen days after the first spray for rice fallow pulses.
3. To control powdery mildew in rice fallow pulses, spray 5% Neem Seed Kernel Extract or Mancozeb @ 400 g/ac or Wettable Sulphur @ 1 kg/ac during morning or evening hours.