



# Weather Advisory Bulletin – Chennai District (Period 17<sup>th</sup> to 21<sup>th</sup> Aug. 2017)

#### A. Weather for the past five days (11.08.2017 to 15.08.2017)

During past five days around Chennai area, the sky was partly cloudy to mostly cloudy with 16.5mm of rainfall. Maximum temperature was 34.0-35.0°C and Minimum temperature was 25.0-27.0°C. Wind speed was 4-13 km per hour. Relative humidity was recorded between 66-83%.

#### B. Weather forecast for next five days (Until 08.30 hrs of 21.08.2017)

| D. Weather forecast for next five days (Chan 00.50 hrs of 21.00.2017) |               |               |               |               |               |
|---|---------------|---------------|---------------|---------------|---------------|
| Parameters  | 17-Aug-17     | 18-Aug-17     | 19-Aug-17     | 20-Aug-17     | 21-Aug-17     |
| Cloud Cover   | Partly Cloudy |
| Rainfall (mm)   | 2             | 2             | 0             | 0             | 0             |
| Wind Speed Kmph   | 7             | 8             | 8             | 7             | 6             |
| Wind direction  | South West    | South West    | West          | West          | West          |
| Temperature °C  |               |               |               |               |               |
| Maximum   | 34            | 34            | 35            | 35            | 36            |
| Minimum   | 26            | 26            | 27            | 27            | 28            |
| Relative Humidity %   | 70-75         | 65-70         | 65-70         | 60-65         | 60-65         |
|   |               |               |               |               |               |

### Livestock:

## Agrometeorological Advisories

- 1. Light rainfall may be expected for next two day.
- 2. To control fluke infestation, animal should not be allowed to graze near ponds and lakes.
- 3. Farmers are advised to wash and clean the udder before milking and not to provide dusty roughage immediately after milking to avoid mastitis.
- 4. Sheep and goat shed remove 3-4 inches of mud floor and replace it with new mud to avoid infection.
- 5. Sheep and goat should be dewormed by using suitable anthelminthic by consulting with local veterinary doctor.
- 6. Backyard poultry farmer advised to vaccinate against ranikhet disease.

## Agriculture (Krishi Vigyan Kendra, Kattupakkam)

## Imp. Crops: Rice, Sugarcane and Groundnut

- 1. To control thrips and other sucking pest in cotton spray *Imidacloprid* 200 SL @ 100 ml/ha (or) *Dimethoate* 30 EC @ 500 ml/ha or *Azadirachtin* 0.03% EC @ 500 ml/ha.
- 2. To control bacterial leaf blight in rice spray Copper hydroxide 77 WP @ 2.5g/ litre or 20% fresh cow dung extract or 5 % Neem Seed Kernel Extract.

Nodal Officer, GKMS and Professor and Head, Department of Livestock Production Management Madras Veterinary College, Chennai – 7