



“Gramin Krishi Mausam Sewa” (GKMS) -Chennai
(Issued jointly by Tamil Nadu Veterinary and
Animal Sciences University,
& India Meteorological Department)



Weather Advisory Bulletin – Chennai District
(Period 25th to 29th Apr. 2018)

A. Weather for the past four days (20.04.2018 to 23.04.2018)

During past four days around Chennai area, the sky was mostly clear without rainfall. Maximum temperature was 35.0-37.0°C and Minimum temperature was 28.0°C. Wind speed was 6-10 km per hour. Relative humidity was recorded between 71-75%.

B. Weather forecast for next five days (Until 08.30 hrs of 29.04.2018)

Parameters	25-Apr-18	26-Apr-18	27-Apr-18	28-Apr-18	29-Apr-18
Cloud Cover	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy
Rainfall (mm)	0	0	0	0	0
Wind Speed Kmph	12	12	10	10	10
Wind direction	South East	South East	South	South	South
Temperature °C					
Maximum	37	37	38	38	38
Minimum	28	28	28	28	28
Relative Humidity %	65-70	65-70	63-70	63-70	63-70

Agrometeorological Advisories

Livestock:

1. Due to increase day temperature farmer's are advised to sprinkle water on the roof of the animal shed to reduce heat stress to the animals.
2. Poultry farmers are advised to cover the water tank by roof and pipe lines with wet gunny bags to provide cool drinking water during hot part of the day.
3. Due to increase in day temperature poultry farmers are advised to raise the feeders between 10 am to 5 pm, Provide mineral mixture and B-complex vitamins in drinking water to avoid heat stroke.
4. During day time farmers are advised to sprinkle water to high yielding dairy cows 2 or 3 time to reduce heat stress.
5. Avoid grazing of animals near ponds and lakes to avoid fluke infestation.
6. Sheep and goat should be dewormed by using suitable anthelmintic by consulting with local veterinary doctor.

Agriculture (Krishi Vigyan Kendra, Kattupakkam)

Imp. Crops: Rice, Sugarcane and Groundnut

1. To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent.
2. Two percent of DAP spray (Dissolve 4.5 kg of DAP in 10 litres of water previous day night and the supernatant liquid shall be mixed with 190 litres of water) may be done at the time of flowering and another dose fifteen days after the first spray for rice fallow pulses.
3. To prevent grain discolouration in rice, spray two gram of Carbendazim + Thiram + Mancozeb (1:1:1) mixture in one litre of water at 50% flowering stage or spray NSKE 5%.

*Nodal Officer, GKMS and Professor and Head,
Department of Livestock Production Management
Madras Veterinary College, Chennai – 7*



“Gramin Krishi Mausam Sewa” (GKMS) -Chennai
(Issued jointly by Tamil Nadu Veterinary and
Animal Sciences University,
& India Meteorological Department)



Weather Advisory Bulletin – Kancheepuram District
(Period 25th to 29th Apr. 2018)

A. Weather for the past four days (20.04.2018 to 23.04.2018)

During past four days around Kancheepuram area, the sky was mostly clear to partly cloudy without rainfall. Maximum temperature was 37.0-39.0°C and Minimum temperature was 27.0-28.0°C. Wind speed was 11-17 km per hour. Relative humidity was recorded between 60-74%.

B. Weather forecast for next five days (Until 08.30 hrs of 29.04.2018)

Parameters	25-Apr-18	26-Apr-18	27-Apr-18	28-Apr-18	29-Apr-18
Cloud Cover	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy
Rainfall (mm)	0	0	0	0	0
Wind Speed Kmph	12	12	12	10	10
Wind Direction	South East	South East	South East	South	South
Temperature °C					
Maximum	39	39	39	39	39
Minimum	28	28	28	28	28
Relative Humidity %	60-70	60-70	60-70	60-65	60-65

Agrometeorological Advisories

Livestock:

1. Due to increase day temperature farmer's are advised to sprinkle water on the roof of the animal shed to reduce heat stress to the animals.
2. Poultry farmers are advised to cover the water tank by roof and pipe lines with wet gunny bags to provide cool drinking water during hot part of the day.
3. Due to increase in day temperature poultry farmers are advised to raise the feeders between 10 am to 5 pm, Provide mineral mixture and B-complex vitamins in drinking water to avoid heat stroke.
4. During day time farmers are advised to sprinkle water to high yielding dairy cows 2 or 3 time to reduce heat stress.
5. Avoid grazing of animals near ponds and lakes to avoid fluke infestation.
6. Sheep and goat should be dewormed by using suitable anthelmintic by consulting with local veterinary doctor.

Agriculture (Krishi Vigyan Kendra, Kattupakkam)

Imp. Crops: Rice, Sugarcane and Groundnut

1. To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent.
2. Two percent of DAP spray (Dissolve 4.5 kg of DAP in 10 litres of water previous day night and the supernatant liquid shall be mixed with 190 litres of water) may be done at the time of flowering and another dose fifteen days after the first spray for rice fallow pulses.
3. To prevent grain discolouration in rice, spray two gram of Carbendazim + Thiram + Mancozeb (1:1:1) mixture in one litre of water at 50% flowering stage or spray NSKE 5%



“Gramin Krishi Mausam Sewa” (GKMS) -Chennai
(Issued jointly by Tamil Nadu Veterinary and
Animal Sciences University,
& India Meteorological Department)



Weather Advisory Bulletin – Tiruvallur District
(Period 25th to 29th Apr. 2018)

A. Weather for the past four days (20.04.2018 to 23.04.2018)

During past four days around Tiruvallur area, the sky was mostly clear without rainfall. Maximum temperature was 35.0-37.0°C and Minimum temperature was 28.0°C. Wind speed was 6-10 km per hour. Relative humidity was recorded between 71-75%.

B. Weather forecast for next five days (Until 08.30 hrs of 29.04.2018)

Parameters	25-Apr-18	26-Apr-18	27-Apr-18	28-Apr-18	29-Apr-18
Cloud Cover	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy
Rainfall (mm)	0	0	0	0	0
Wind Speed Kmph	12	12	10	10	10
Wind Direction	South East	South East	South	South	South
Temperature °C					
Maximum	37	37	38	38	38
Minimum	28	28	28	28	28
Relative Humidity %	65-70	65-70	63-70	63-70	63-70

Agrometeorological Advisories

Livestock:

1. Due to increase day temperature farmer's are advised to sprinkle water on the roof of the animal shed to reduce heat stress to the animals .
2. Poultry farmers are advised to cover the water tank by roof and pipe lines with wet gunny bags to provide cool drinking water during hot part of the day.
3. Due to increase in day temperature poultry farmers are advised to raise the feeders between 10 am to 5 pm, Provide mineral mixture and B-complex vitamins in drinking water to avoid heat stroke.
4. During day time farmers are advised to sprinkle water to high yielding dairy cows 2 or 3 time to reduce heat stress.
5. Avoid grazing of animals near ponds and lakes to avoid fluke infestation.
6. Sheep and goat should be dewormed by using suitable anthelmintic by consulting with local veterinary doctor.

Agriculture (Krishi Vigyan Kendra, Kattupakkam)

Imp. Crops: Rice, Sugarcane and Groundnut

1. To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent.
2. Two percent of DAP spray (Dissolve 4.5 kg of DAP in 10 litres of water previous day night and the supernatant liquid shall be mixed with 190 litres of water) may be done at the time of flowering and another dose fifteen days after the first spray for rice fallow pulses.
3. To prevent grain discolouration in rice, spray two gram of Carbendazim + Thiram + Mancozeb (1:1:1) mixture in one litre of water at 50% flowering stage or spray NSKE 5%



“Gramin Krishi Mausam Sewa” (GKMS) -Chennai
(Issued jointly by Tamil Nadu Veterinary and
Animal Sciences University,
& India Meteorological Department)



Weather Advisory Bulletin - Vellore District
(Period 25th to 29th Apr. 2018)

A. Weather for the past four days (20.04.2018 to 23.04.2018)

During past four days around Vellore area, the sky was clear to partly cloudy without rainfall. Maximum temperature was 39.0-41.0°C and Minimum temperature was 26.0-27.0°C. Wind speed was 4-7 km per hour. Relative humidity was recorded between 29-78%.

B. Weather forecast for next five days (Until 08.30 hrs of 29.04.2018)

Parameters	25-Apr-18	26-Apr-18	27-Apr-18	28-Apr-18	29-Apr-18
Cloud Cover	Mostly Clear Sky	Mostly Clear Sky	Partly Cloudy	Partly Cloudy	Partly Cloudy
Rainfall (mm)	0	0	0	0	0
Wind Speed Kmph	6	6	6	6	6
Wind Direction	East South East	East South East	South	South	South
Temperature °C					
Maximum	39	39	39	39	39
Minimum	28	28	27	27	27
Relative Humidity %	40-75	40-75	40-70	40-70	40-70

Agrometeorological Advisories

Livestock:

1. Due to increase day temperature farmer's are advised to sprinkle water on the roof of the animal shed to reduce heat stress to the animals .
2. Poultry farmers are advised to cover the water tank by roof and pipe lines with wet gunny bags to provide cool drinking water during hot part of the day.
3. Due to increase in day temperature poultry farmers are advised to raise the feeders between 10 am to 5 pm, Provide mineral mixture and B-complex vitamins in drinking water to avoid heat stoke.
4. During day time farmers are advised to sprinkle water to high yielding dairy cows 2 or 3 time to reduce heat stress.
5. Avoid grazing of animals near ponds and lakes to avoid fluke infestation.
6. Sheep and goat should be dewormed by using suitable anthelmintic by consulting with local veterinary doctor.

Agriculture

Imp. Crops: Rice, Sugarcane and Groundnut

1. To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent.
2. Two percent of DAP spray (Dissolve 4.5 kg of DAP in 10 litres of water previous day night and the supernatant liquid shall be mixed with 190 litres of water) may be done at the time of flowering and another dose fifteen days after the first spray for rice fallow pulses.
3. To prevent grain discolouration in rice, spray two gram of Carbendazim + Thiram + Mancozeb (1:1:1) mixture in one litre of water at 50% flowering stage or spray NSKE 5%



“Gramin Krishi Mausam Sewa” (GKMS) -Chennai
(Issued jointly by Tamil Nadu Veterinary and
Animal Sciences University,
& India Meteorological Department)



Weather Advisory Bulletin – Thiruvannamalai District
(Period 25th to 29th Apr. 2018)

A. Weather for the past four days (20.04.2018 to 23.04.2018)

During past four days around Thiruvannamalai district, the sky was clear to partly cloudy without rainfall. Maximum temperature was 39.0-41.0°C and Minimum temperature was 26.0-27.0°C. Wind speed was 4-7 km per hour. Relative humidity was recorded between 29-78%.

B. Weather forecast for next five days (Until 08.30 hrs of 29.04.2018)

Parameters	25-Apr-18	26-Apr-18	27-Apr-18	28-Apr-18	29-Apr-18
Cloud Cover	Mostly Clear Sky	Mostly Clear Sky	Partly Cloudy	Partly Cloudy	Partly Cloudy
Rainfall (mm)	0	1	0	0	0
Wind Speed Kmph	6	6	6	6	6
Wind Direction	East South East	East South East	South	South	South
Temperature °C					
Maximum	39	39	39	39	39
Minimum	28	28	27	27	27
Relative Humidity %	40-75	40-75	40-70	40-70	40-70

Agrometeorological Advisories

Livestock:

1. Due to increase day temperature farmer's are advised to sprinkle water on the roof of the animal shed to reduce heat stress to the animals .
2. Poultry farmers are advised to cover the water tank by roof and pipe lines with wet gunny bags to provide cool drinking water during hot part of the day.
3. Due to increase in day temperature poultry farmers are advised to raise the feeders between 10 am to 5 pm, Provide mineral mixture and B-complex vitamins in drinking water to avoid heat stoke.
4. During day time farmers are advised to sprinkle water to high yielding dairy cows 2 or 3 time to reduce heat stress.
5. Avoid grazing of animals near ponds and lakes to avoid fluke infestation.
6. Sheep and goat should be dewormed by using suitable anthelmintic by consulting with local veterinary doctor.

Agriculture

Imp. Crops: Rice, Sugarcane and Groundnut

1. To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent.
2. Two percent of DAP spray (Dissolve 4.5 kg of DAP in 10 litres of water previous day night and the supernatant liquid shall be mixed with 190 litres of water) may be done at the time of flowering and another dose fifteen days after the first spray for rice fallow pulses.
3. To prevent grain discolouration in rice, spray two gram of Carbendazim + Thiram + Mancozeb (1:1:1) mixture in one litre of water at 50% flowering stage or spray NSKE 5%



“Gramin Krishi Mausam Sewa” (GKMS) -Chennai
(Issued jointly by Tamil Nadu Veterinary and
Animal Sciences University,
& India Meteorological Department)



Weather Advisory Bulletin – Cuddalore District
(Period 25th to 29th Apr. 2018)

A. Weather for the past four days (20.04.2018 to 23.04.2018)

During past four days around Cuddalore district, the sky was clear to partly cloudy without rainfall. Maximum temperature was 35.0-36.0°C and Minimum temperature was 27.0-28.0°C. Wind speed was 5-6 km per hour. Relative humidity was recorded between 70-77%.

B. Weather forecast for next five days (Until 08.30 hrs of 29.04.2018)

Parameters	25-Apr-18	26-Apr-18	27-Apr-18	28-Apr-18	29-Apr-18
Cloud Cover	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy
Rainfall (mm)	0	0	0	0	0
Wind Speed Kmph	10	10	10	10	10
Wind Direction	East South East	East South East	South	South	South
Temperature °C					
Maximum	36	36	37	37	37
Minimum	27	27	28	28	28
Relative Humidity %	65-70	65-70	60-70	60-70	60-70

Agrometeorological Advisories

Livestock:

1. Due to increase day temperature farmer’s are advised to sprinkle water on the roof of the animal shed to reduce heat stress to the animals .
2. Poultry farmers are advised to cover the water tank by roof and pipe lines with wet gunny bags to provide cool drinking water during hot part of the day.
3. Due to increase in day temperature poultry farmers are advised to raise the feeders between 10 am to 5 pm, Provide mineral mixture and B-complex vitamins in drinking water to avoid heat stoke.
4. During day time farmers are advised to sprinkle water to high yielding dairy cows 2 or 3 time to reduce heat stress.
5. Avoid grazing of animals near ponds and lakes to avoid fluke infestation.
6. Sheep and goat should be dewormed by using suitable anthelminthic by consulting with local veterinary doctor.

Agriculture

Imp. Crops: Rice, Sugarcane and Groundnut

1. To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent.
2. Two percent of DAP spray (Dissolve 4.5 kg of DAP in 10 litres of water previous day night and the supernatant liquid shall be mixed with 190 litres of water) may be done at the time of flowering and another dose fifteen days after the first spray for rice fallow pulses.
3. To prevent grain discolouration in rice, spray two gram of Carbendazim + Thiram + Mancozeb (1:1:1) mixture in one litre of water at 50% flowering stage or spray NSKE 5%



“Gramin Krishi Mausam Sewa” (GKMS) –Chennai
(Issued jointly by Tamil Nadu Veterinary and
Animal Sciences University,
& India Meteorological Department)



Weather Advisory Bulletin – Villupuram District
(Period 21st to 25th Apr. 2018)

A. Weather for the past three days (17.04.2018 to 19.04.2018)

During past three days around Villupuram district, the sky was clear to partly cloudy without rainfall. Maximum temperature was 35.0-36.0°C and Minimum temperature was 27.0-28.0°C. Wind speed was 5-6 km per hour. Relative humidity was recorded between 70-77%.

B. Weather forecast for next five days (Until 08.30 hrs of 25.04.2018)

Parameters	25-Apr-18	26-Apr-18	27-Apr-18	28-Apr-18	29-Apr-18
Cloud Cover	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy
Rainfall (mm)	0	0	0	0	0
Wind Speed Kmph	10	10	10	10	10
Wind Direction	East South East	East South East	South	South	South
Temperature °C					
Maximum	36	36	37	37	37
Minimum	27	27	28	28	28
Relative Humidity %	65-70	65-70	60-70	60-70	60-70

Agrometeorological Advisories

Livestock:

1. Due to increase day temperature farmer's are advised to sprinkle water on the roof of the animal shed to reduce heat stress to the animals .
2. Poultry farmers are advised to cover the water tank by roof and pipe lines with wet gunny bags to provide cool drinking water during hot part of the day.
3. Due to increase in day temperature poultry farmers are advised to raise the feeders between 10 am to 5 pm, Provide mineral mixture and B-complex vitamins in drinking water to avoid heat stroke.
4. During day time farmers are advised to sprinkle water to high yielding dairy cows 2 or 3 time to reduce heat stress.
5. Avoid grazing of animals near ponds and lakes to avoid fluke infestation.
6. Sheep and goat should be dewormed by using suitable anthelmintic by consulting with local veterinary doctor.

Agriculture

Imp. Crops: Rice, Sugarcane and Groundnut

1. To control the incidence of sucking pest in the vegetable crops farmer are advised to spray neem seed kernal extract 5 percent.
2. Two percent of DAP spray (Dissolve 4.5 kg of DAP in 10 litres of water previous day night and the supernatant liquid shall be mixed with 190 litres of water) may be done at the time of flowering and another dose fifteen days after the first spray for rice fallow pulses.
3. To prevent grain discolouration in rice, spray two gram of Carbendazim + Thiram + Mancozeb (1:1:1) mixture in one litre of water at 50% flowering stage or spray NSKE 5%