



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **ANNAVASAL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	0.0	10.0	29.0	9.0	3.0
Tmax(°C)	35.0	36.0	36.0	35.0	36.0
Tmin(°C)	27.0	28.0	27.0	28.0	28.0
RH-I(%)	76	76	79	78	71
RH-II(%)	45	44	44	43	42
Wind Speed(kmph)	12.0	8.0	7.0	10.0	8.0
Wind Direction(Degree)	187	236	236	237	270
Cloud Cover(Octa)	5	6	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 18,19 and 20 August. Maximum temperature would be in the range 34.5 °C and 36.7°C and the minimum temperature would be in the range 24.6°C- 26.2°C . Cloudy sky will appear . Wind speed is expected to be 11 to 16 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
CORIANDER	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **ARANTANGI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	0.0	9.0	25.0	13.0	3.0
Tmax(°C)	35.0	36.0	36.0	35.0	36.0
Tmin(°C)	27.0	28.0	28.0	28.0	29.0
RH-I(%)	76	76	78	78	71
RH-II(%)	45	44	44	44	42
Wind Speed(kmph)	12.0	9.0	8.0	10.0	8.0
Wind Direction(Degree)	186	232	222	233	267
Cloud Cover(Octa)	5	6	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 18,19 and 20 August. Maximum temperature would be in the range 36.7°C and 37.8°C and the minimum temperature would be in the range 29.1°C- 30.6°C . Cloudy sky will appear . Wind speed is expected to be 12 to 19 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
CORIANDER	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **ARIMALAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	0.0	7.0	22.0	9.0	2.0
Tmax(°C)	35.0	36.0	36.0	35.0	36.0
Tmin(°C)	27.0	28.0	28.0	28.0	29.0
RH-I(%)	77	76	78	77	71
RH-II(%)	46	45	44	43	42
Wind Speed(kmph)	13.0	8.0	7.0	10.0	7.0
Wind Direction(Degree)	186	233	230	234	269
Cloud Cover(Octa)	5	6	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 18,19 and 20 August. Maximum temperature would be in the range 34.3°C and 36.2°C and the minimum temperature would be in the range 24.4°C- 25.8°C . Cloudy sky will appear . Wind speed is expected to be 12 to 18 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
CORIANDER	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **AVADAIYARKOVIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	0.0	10.0	26.0	17.0	4.0
Tmax(°C)	35.0	36.0	36.0	35.0	36.0
Tmin(°C)	27.0	28.0	28.0	28.0	28.0
RH-I(%)	77	75	78	78	72
RH-II(%)	45	45	44	44	43
Wind Speed(kmph)	12.0	9.0	8.0	10.0	8.0
Wind Direction(Degree)	185	231	211	232	265
Cloud Cover(Octa)	5	6	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 18,19 and 20 August. Maximum temperature would be in the range 35.8 °C and 36.7°C and the minimum temperature would be in the range 28.6°C- 30.1°C . Cloudy sky will appear . Wind speed is expected to be 13 to 21 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
CORIANDER	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **GANDARAVAKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	4.0	7.0	34.0	5.0	0.0
Tmax(°C)	35.0	36.0	36.0	35.0	36.0
Tmin(°C)	27.0	28.0	28.0	28.0	28.0
RH-I(%)	77	79	83	81	72
RH-II(%)	48	45	46	47	43
Wind Speed(kmph)	13.0	10.0	7.0	10.0	10.0
Wind Direction(Degree)	191	211	270	230	262
Cloud Cover(Octa)	5	7	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 18,19 and 20 August. Maximum temperature would be in the range 33.9°C and 36.1°C and the minimum temperature would be in the range 24.2°C- 26°C . Cloudy sky will appear . Wind speed is expected to be 12 to 17 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
CORIANDER	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **KARAMBAKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	2.0	9.0	32.0	8.0	1.0
Tmax(°C)	35.0	36.0	36.0	35.0	36.0
Tmin(°C)	27.0	28.0	27.0	28.0	28.0
RH-I(%)	76	77	81	80	71
RH-II(%)	46	44	44	45	42
Wind Speed(kmph)	12.0	9.0	7.0	10.0	9.0
Wind Direction(Degree)	189	224	249	233	265
Cloud Cover(Octa)	5	7	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 18,19 and 20 August. Maximum temperature would be in the range 36.6°C and 33.8°C and the minimum temperature would be in the range 24.2°C- 26.1°C . Cloudy sky will appear . Wind speed is expected to be 13 to 19 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
CORIANDER	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **KUNNANDARKOIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	1.0	19.0	51.0	11.0	5.0
Tmax(°C)	36.0	36.0	35.0	35.0	36.0
Tmin(°C)	26.0	26.0	26.0	26.0	27.0
RH-I(%)	73	75	82	78	69
RH-II(%)	43	42	43	43	41
Wind Speed(kmph)	10.0	9.0	8.0	10.0	11.0
Wind Direction(Degree)	188	245	254	245	271
Cloud Cover(Octa)	5	7	8	8	8

Weather Summary/Alert:

Chances of light rain on 18,19 and 20 August. Maximum temperature would be in the range 35°C and 36 °C and the minimum temperature would be in the range 26°C- 27°C . Cloudy sky will appear . Wind speed is expected to be 8 to 11 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
JASMINE	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **MANALMELKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	0.0	10.0	27.0	19.0	5.0
Tmax(°C)	35.0	36.0	36.0	35.0	36.0
Tmin(°C)	27.0	28.0	28.0	28.0	28.0
RH-I(%)	77	75	78	79	72
RH-II(%)	46	45	44	45	43
Wind Speed(kmph)	13.0	9.0	8.0	10.0	9.0
Wind Direction(Degree)	186	229	209	232	263
Cloud Cover(Octa)	5	6	8	8	8

Weather Summary/Alert:

Chances of light rain on 18,19 and 20 August. Maximum temperature would be in the range 35°C and 36 °C and the minimum temperature would be in the range 27°C- 28°C . Cloudy sky will appear . Wind speed is expected to be 8 to 13 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
JASMINE	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **PONNAMARAVATI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	0.0	11.0	28.0	15.0	4.0
Tmax(°C)	35.0	36.0	36.0	35.0	36.0
Tmin(°C)	27.0	28.0	27.0	28.0	28.0
RH-I(%)	76	75	78	78	71
RH-II(%)	44	44	43	43	42
Wind Speed(kmph)	12.0	8.0	8.0	10.0	8.0
Wind Direction(Degree)	185	236	215	234	268
Cloud Cover(Octa)	5	6	8	8	8

Weather Summary/Alert:

Chances of light rain on 18,19 and 20 August. Maximum temperature would be in the range 35°C and 36 °C and the minimum temperature would be in the range 27°C- 28°C . Cloudy sky will appear . Wind speed is expected to be 8 to 12 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
JASMINE	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **PUDUKKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	0.0	6.0	19.0	7.0	2.0
Tmax(°C)	35.0	36.0	36.0	35.0	36.0
Tmin(°C)	27.0	28.0	28.0	28.0	29.0
RH-I(%)	77	76	78	77	71
RH-II(%)	46	45	44	43	42
Wind Speed(kmph)	13.0	8.0	7.0	10.0	6.0
Wind Direction(Degree)	186	233	233	234	270
Cloud Cover(Octa)	5	6	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 18,19 and 20 August. Maximum temperature would be in the range 35°C and 36°C and the minimum temperature would be in the range 27°C- 29°C . Cloudy sky will appear . Wind speed is expected to be 7 to 18 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
JASMINE	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **THIRUVARANGULAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	0.0	6.0	19.0	7.0	2.0
Tmax(°C)	35.0	36.0	36.0	35.0	36.0
Tmin(°C)	27.0	28.0	28.0	28.0	29.0
RH-I(%)	77	76	78	77	71
RH-II(%)	46	45	44	43	42
Wind Speed(kmph)	13.0	8.0	7.0	10.0	6.0
Wind Direction(Degree)	186	233	233	234	270
Cloud Cover(Octa)	5	6	8	8	8

Weather Summary/Alert:

Chances of light rain on 18,19 and 20 August. Maximum temperature would be in the range 35°C and 36 °C and the minimum temperature would be in the range 27°C- 29°C . Cloudy sky will appear . Wind speed is expected to be 8 to 13 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
JASMINE	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **TIRUMAYAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	0.0	8.0	23.0	10.0	2.0
Tmax(°C)	35.0	36.0	36.0	35.0	36.0
Tmin(°C)	27.0	28.0	28.0	28.0	29.0
RH-I(%)	77	76	78	77	71
RH-II(%)	45	45	44	43	42
Wind Speed(kmph)	13.0	8.0	7.0	10.0	7.0
Wind Direction(Degree)	186	234	226	234	269
Cloud Cover(Octa)	5	6	8	8	8

Weather Summary/Alert:

Chances of light rain on 18,19 and 20 August. Maximum temperature would be in the range 35°C and 36 °C and the minimum temperature would be in the range 27°C- 29°C . Cloudy sky will appear . Wind speed is expected to be 8 to 13 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
JASMINE	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 16-08-2022

Weather Forecast of **VIRALIMALAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-17	2022-08-18	2022-08-19	2022-08-20	2022-08-21
Rainfall	2.0	20.0	54.0	12.0	6.0
Tmax(°C)	36.0	36.0	35.0	35.0	36.0
Tmin(°C)	26.0	26.0	26.0	26.0	27.0
RH-I(%)	73	74	82	78	69
RH-II(%)	42	42	42	43	41
Wind Speed(kmph)	10.0	9.0	8.0	10.0	12.0
Wind Direction(Degree)	188	247	251	246	271
Cloud Cover(Octa)	5	7	8	8	8

Weather Summary/Alert:

Chances of light rain on 18,19 and 20 August. Maximum temperature would be in the range 35°C and 36 °C and the minimum temperature would be in the range 27°C- 29°C . Cloudy sky will appear . Wind speed is expected to be 8 to 13 kmph.

General Advisory:

To control leaf folder incidence in rice, apply fipronil 2ml/litre.

SMS Advisory:

Farmers are advised to harvest the matured paddy as soon as possible.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
JASMINE	Due to prevailing weather alternaria leaf spot incidence may occur in jasmine. To control, spray foliar application of mancozeb @ 2.5 g/l at monthly interval.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	To control Hadda beetle incidence in Brinjal, spray Acepate @ 2 gm /litre.
CHILLI	With the expected rainfall, chilli can be grown at intervals of 45 cm with intercropping of coriander or small onion between the two rows to control weeds and get extra income.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.