



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **ANNAVASAL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	2.0	4.0	14.0	4.0	3.0
Tmax(°C)	37.0	35.0	36.0	35.0	37.0
Tmin(°C)	28.0	27.0	26.0	27.0	27.0
RH-I(%)	61	66	72	70	67
RH-II(%)	38	44	40	43	37
Wind Speed(kmph)	10.0	10.0	9.0	7.0	9.0
Wind Direction(Degree)	279	279	179	252	280
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of very light rain on 25 to 28 August . Maximum temperature would be in the range 33.3°C and 37.1°C and the minimum temperature would be in between 23.7°C - 28.0°C. lousy sky is expected for next 5 days. Wind speed is expected to be 15 to 22 kmph. xpected to be 11 to 22 kmph.

General Advisory:

Fruit trees and Tree spices.Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
COCONUT	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Since moderate rain is forecasted in the fourth coming week farmers are advised to tether livestock in the inside sheds, to prevent from rain water drenching. The concentrated feed and dry fodder must be stored in water proof god-owns to prevent aflotoxin and mould development.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **ARANTANGI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	3.0	3.0	13.0	4.0	3.0
Tmax(°C)	37.0	35.0	36.0	35.0	37.0
Tmin(°C)	29.0	28.0	26.0	27.0	28.0
RH-I(%)	61	66	72	70	67
RH-II(%)	38	45	41	43	37
Wind Speed(kmph)	10.0	10.0	9.0	7.0	8.0
Wind Direction(Degree)	277	278	170	249	275
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain on 26 August .Maximum temperature would be in the range 35.8°C and 38.3°C and the minimum temperature would be in between 28.1°C - 31.0°C.Cloudy sky is expected for next 5 days. lousy sky is expected for next 5 days. Wind speed is expected to be 09 to 14 kmph.

General Advisory:

Fruit trees and Tree spices.Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
COCONUT	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Since moderate rain is forecasted in the fourth coming week farmers are advised to tether livestock in the inside sheds, to prevent from rain water drenching. The concentrated feed and dry fodder must be stored in water proof god-owns to prevent aflotoxin and mould development.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **ARIMALAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	1.0	3.0	14.0	4.0	3.0
Tmax(°C)	37.0	35.0	36.0	35.0	37.0
Tmin(°C)	29.0	28.0	26.0	27.0	28.0
RH-I(%)	61	66	72	70	66
RH-II(%)	38	44	40	43	37
Wind Speed(kmph)	9.0	9.0	9.0	6.0	8.0
Wind Direction(Degree)	278	280	166	250	279
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain on 25 and 26 August . Maximum temperature would be in the range 32.6°C and 36.1°C and the minimum temperature would be in between 23.8°C - 25.6°C.Cloudy sky is expected for next 5 days. lousy sky is expected for next 5 days. Wind speed is expected to be 09 to 14 kmph.

General Advisory:

Fruit trees and Tree spices.Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
COCONUT	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Since moderate rain is forecasted in the fourth coming week farmers are advised to tether livestock in the inside sheds, to prevent from rain water drenching. The concentrated feed and dry fodder must be stored in water proof god-owns to prevent aflotoxin and mould development.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vinrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **AVADAIYARKOVIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	3.0	3.0	13.0	4.0	3.0
Tmax(°C)	37.0	35.0	36.0	35.0	37.0
Tmin(°C)	29.0	28.0	26.0	27.0	28.0
RH-I(%)	61	66	71	70	67
RH-II(%)	38	45	41	43	38
Wind Speed(kmph)	10.0	10.0	9.0	7.0	8.0
Wind Direction(Degree)	276	277	169	249	271
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain on 26 August .Maximum temperature would be in the range 34.6°C and 37.2°C and the minimum temperature would be in between 27.3°C - 30.5°C. Cloudy sky is expected for next 5 days. lousy sky is expected for next 5 days. Wind speed is expected to be 09 to 13 kmph.

General Advisory:

Fruit trees and Tree spices. Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
COCONUT	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Since moderate rain is forecasted in the fourth coming week farmers are advised to tether livestock in the inside sheds, to prevent from rain water drenching. The concentrated feed and dry fodder must be stored in water proof god-owns to prevent aflotoxin and mould development.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vinrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **GANDARAVAKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	7.0	6.0	14.0	4.0	4.0
Tmax(°C)	37.0	36.0	35.0	35.0	36.0
Tmin(°C)	29.0	28.0	26.0	27.0	28.0
RH-I(%)	66	69	74	72	70
RH-II(%)	39	42	43	46	39
Wind Speed(kmph)	10.0	13.0	9.0	9.0	11.0
Wind Direction(Degree)	282	267	218	235	279
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain on 24,25 and 27 August .Maximum temperature would be in the range 34.1°C and 36.2°C and the minimum temperature would be in between 23.6°C - 24.8°C. Cloudy sky is expected for next 5 days. Cloudy sky is expected for next 5 days. Wind speed is expected to be 12 to 23 kmph.

General Advisory:

Fruit trees and Tree spices. Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
COCONUT	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Since moderate rain is forecasted in the fourth coming week farmers are advised to tether livestock in the inside sheds, to prevent from rain water drenching. The concentrated feed and dry fodder must be stored in water proof god-owns to prevent aflotoxin and mould development.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vinrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **KARAMBAKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	5.0	5.0	14.0	4.0	4.0
Tmax(°C)	37.0	35.0	36.0	35.0	37.0
Tmin(°C)	28.0	27.0	26.0	27.0	27.0
RH-I(%)	64	68	73	71	68
RH-II(%)	38	43	42	44	38
Wind Speed(kmph)	10.0	12.0	9.0	8.0	10.0
Wind Direction(Degree)	280	273	199	244	278
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain on 25 to 28 August . Maximum temperature would be in the range 34.4°C and 36.3°C and the minimum temperature would be in between 24.1°C - 25.3°C. loudy sky is expected for next 5 days. Wind speed is expected to be 12 to 18 kmph.

General Advisory:

Fruit trees and Tree spices.Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
COCONUT	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Since moderate rain is forecasted in the fourth coming week farmers are advised to tether livestock in the inside sheds, to prevent from rain water drenching. The concentrated feed and dry fodder must be stored in water proof god-owns to prevent aflotoxin and mould development.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vinrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **KUNNANDARKOIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	5.0	8.0	13.0	7.0	5.0
Tmax(°C)	37.0	35.0	36.0	35.0	37.0
Tmin(°C)	27.0	26.0	26.0	26.0	26.0
RH-I(%)	63	70	71	69	68
RH-II(%)	38	43	41	43	36
Wind Speed(kmph)	13.0	13.0	10.0	9.0	12.0
Wind Direction(Degree)	279	276	222	260	280
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain in Next five days. Maximum temperature would be in the range 35.0°C and 37.0°C and the minimum temperature would be in between 26.0°C - 27.0°C. loudy sky is expected for next 5 days. Wind speed is expected to be 09 to 13 kmph.

General Advisory:

Fruit trees and Tree spices.Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
COCONUT	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Since moderate rain is forecasted in the fourth coming week farmers are advised to tether livestock in the inside sheds, to prevent from rain water drenching. The concentrated feed and dry fodder must be stored in water proof god-owns to prevent aflotoxin and mould development.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vinrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **MANALMELKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	4.0	3.0	12.0	4.0	4.0
Tmax(°C)	37.0	35.0	35.0	35.0	37.0
Tmin(°C)	29.0	28.0	26.0	27.0	28.0
RH-I(%)	62	67	71	70	67
RH-II(%)	39	46	42	44	38
Wind Speed(kmph)	10.0	10.0	9.0	8.0	9.0
Wind Direction(Degree)	275	276	174	248	269
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain in Next five days. Maximum temperature would be in the range 35.0°C and 37.0°C and the minimum temperature would be in between 26.0°C - 29.0°C. loudy sky is expected for next 5 days. Wind speed is expected to be 09 to 10 kmph.

General Advisory:

Fruit trees and Tree spices.Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
COCONUT	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Since moderate rain is forecasted in the fourth coming week farmers are advised to tether livestock in the inside sheds, to prevent from rain water drenching. The concentrated feed and dry fodder must be stored in water proof god-owns to prevent aflotoxin and mould development.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vinrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **PONNAMARAVATI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	3.0	4.0	13.0	3.0	3.0
Tmax(°C)	37.0	35.0	36.0	35.0	37.0
Tmin(°C)	28.0	27.0	26.0	27.0	27.0
RH-I(%)	61	67	71	69	66
RH-II(%)	38	45	41	43	37
Wind Speed(kmph)	10.0	10.0	9.0	7.0	8.0
Wind Direction(Degree)	277	279	169	252	274
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain in Next five days. Maximum temperature would be in the range 35.0°C and 37.0°C and the minimum temperature would be in between 26.0°C - 28.0°C. loudy sky is expected for next 5 days. Wind speed is expected to be 07 to 10 kmph.

General Advisory:

Fruit trees and Tree spices.Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
COCONUT	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Since moderate rain is forecasted in the fourth coming week farmers are advised to tether livestock in the inside sheds, to prevent from rain water drenching. The concentrated feed and dry fodder must be stored in water proof god-owns to prevent aflotoxin and mould development.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vinrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **PUDUKKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	1.0	3.0	14.0	4.0	3.0
Tmax(°C)	37.0	35.0	36.0	35.0	37.0
Tmin(°C)	29.0	28.0	26.0	27.0	28.0
RH-I(%)	60	65	72	70	66
RH-II(%)	38	44	40	43	37
Wind Speed(kmph)	9.0	9.0	9.0	6.0	7.0
Wind Direction(Degree)	279	281	160	250	281
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain on 25 to 28 August . Maximum temperature would be in the range 35.0°C and 37.0°C and the minimum temperature would be in between 26.0°C - 29.0°C. loudy sky is expected for next 5 days. Wind speed is expected to be 06 to 09 kmph.

General Advisory:

Fruit trees and Tree spices.Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
COCONUT	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Since moderate rain is forecasted in the fourth coming week farmers are advised to tether livestock in the inside sheds, to prevent from rain water drenching. The concentrated feed and dry fodder must be stored in water proof god-owns to prevent aflotoxin and mould development.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vinrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **THIRUVARANGULAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	1.0	3.0	14.0	4.0	3.0
Tmax(°C)	37.0	35.0	36.0	35.0	37.0
Tmin(°C)	29.0	28.0	26.0	27.0	28.0
RH-I(%)	60	65	72	70	66
RH-II(%)	38	44	40	43	37
Wind Speed(kmph)	9.0	9.0	9.0	6.0	7.0
Wind Direction(Degree)	279	281	162	250	281
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain on 25 to 28 August . Maximum temperature would be in the range 35.0°C and 37.0°C and the minimum temperature would be in between 26.0°C - 29.0°C. loudy sky is expected for next 5 days. Wind speed is expected to be 06 to 09 kmph.

General Advisory:

Fruit trees and Tree spices.Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
COCONUT	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Since moderate rain is forecasted in the fourth coming week farmers are advised to tether livestock in the inside sheds, to prevent from rain water drenching. The concentrated feed and dry fodder must be stored in water proof god-owns to prevent aflotoxin and mould development.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vinrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **TIRUMAYAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	1.0	3.0	14.0	4.0	3.0
Tmax(°C)	37.0	35.0	36.0	35.0	37.0
Tmin(°C)	29.0	28.0	26.0	27.0	28.0
RH-I(%)	61	66	72	70	66
RH-II(%)	38	44	40	43	37
Wind Speed(kmph)	9.0	9.0	9.0	7.0	8.0
Wind Direction(Degree)	278	280	165	250	278
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain on 25 to 28 August . Maximum temperature would be in the range 35.0°C and 37.0°C and the minimum temperature would be in between 26.0°C - 29.0°C. loudy sky is expected for next 5 days. Wind speed is expected to be 07 to 09 kmph.

General Advisory:

Fruit trees and Tree spices.Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
COCONUT	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Since moderate rain is forecasted in the fourth coming week farmers are advised to tether livestock in the inside sheds, to prevent from rain water drenching. The concentrated feed and dry fodder must be stored in water proof god-owns to prevent aflotoxin and mould development.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vinrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 23-08-2022

Weather Forecast of **VIRALIMALAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-08-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-24	2022-08-25	2022-08-26	2022-08-27	2022-08-28
Rainfall	5.0	8.0	13.0	7.0	5.0
Tmax(°C)	37.0	35.0	36.0	35.0	37.0
Tmin(°C)	27.0	26.0	25.0	26.0	26.0
RH-I(%)	63	70	70	69	68
RH-II(%)	38	43	41	43	36
Wind Speed(kmph)	13.0	14.0	10.0	9.0	12.0
Wind Direction(Degree)	278	276	224	261	279
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain in Next five days. Maximum temperature would be in the range 35.0°C and 37.0°C and the minimum temperature would be in between 26.0°C - 27.0°C. loudy sky is expected for next 5 days. Wind speed is expected to be 09 to 14 kmph.

General Advisory:

Fruit trees and Tree spices.Strengthening the basins may be taken up for better harvesting of rains.

SMS Advisory:

Due to light to moderate rainfall forecast, farmers are advised to provide adequate drainage facilities to the samba rice nursery to avoid seedling damage.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Since the rainfall is light to moderate the transplanting of rice may be continued with proper drainage. Since the wind speed is more than 12 km/hr, postpone the top dressing of nitrogen to 30 days old transplanted rice.
RICE	To increase the productive tillers, as well as yield of paddy farmers are advised to apply 10 kg of farmyard manure enriched with paddy micro nutrient mixture as basal dose per acre under kharif season.
RICE	Direct sowing in samba field is recommended at the rate of 15 kg/acre for tractor sowing. Before sowing the seeds should be treated with pseudomonas at the rate 10g/kg seed. seed-borne pathogens are destroyed. sowing should be before the monsoon.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Staking has to be provide for the Banana plantation for protecting against wind.
BANANA	The present Weather condition causes boron deficiency. To Rectify this apply borax @ 50g/ per tree along with sand or manure.
BANANA	To increase the soil fertility, conserve the soil moisture and control the weeds, farmers are requested to sow cowpea @ 8 kg / acre in between the rows of banana plant.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Prevailing weather condition the incense of leaf Spot in Brinjal. During rainy and cloudy weather the attack of leaf spot is heavy. To control spray any one of the followings 2 gm Copper oxychloride or 2 ½ gm Zineb per litre of water .

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	அடுத்து வரும் நாட்களில் மழை பெய்ய வாய்ப்புள்ளதால் கால்நடைகளை மழைச்சாரல் நனையாமல் இருக்கும்படி பார்த்துக் கொள்ள வேண்டும் . உலர் தீவனம் மற்றும் அடர் தீவனங்களை தண்ணீர் புகாத சேமிப்பு கிடங்குகளில் பாதுகாத்து வைக்க வேண்டும்.
GOAT	To counteract over occurrence of infection during rainy season pregnant goat (adult goat) must be provided with vitamin and mineral supplementation.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea,keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup perlitre of water. Medicated water should be given to all the birdsfor 5 days in the morning and evening time.