



**Gramin Krishi Mausam Sewa**  
**District Level Agromet Advisory Bulletin**  
 Tamil Nadu Agricultural University,  
 Coimbatore, Tamil Nadu



## Agromet Advisory Bulletin

Date : 26-08-2022

Weather Forecast of District THIRUVALLUR(Tamil Nadu) Issued On : 2022-08-26(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-08-27	2022-08-28	2022-08-29	2022-08-30	2022-08-31
Rainfall(mm)	10.0	10.0	12.0	20.0	18.0
Tmax(°C)	35.0	34.0	34.0	34.0	33.0
Tmin(°C)	26.0	25.0	25.0	24.0	24.0
RH-I(%)	80	80	80	80	80
RH-II(%)	40	40	50	50	50
Wind Speed(kmph)	8.0	8.0	8.0	8.0	6.0
Wind Direction(Degree)	280	300	300	40	40
Cloud Cover(Octa)	8	8	8	8	8

### Weather Summary/Alert:

Moderate rainfall may be expected for forthcoming days; Wind speeds may be range from 6.0-8.0 km/ hour; Wind direction may be in the North East to West North West direction; The maximum temperature may be range from 33.0°C-35.0°C and minimum temperatures may be range from 24.0°C-26.0°C; Morning Relative humidity may be range 80% and Evening Relative humidity may be range from 40-50%.

### General Advisory:

From extended range forecast Above normal rainfall, normal maximum temperature and minimum temperature expected for the period of 31.08.2022 to 06.09.2022 over Tamil Nadu.

### SMS Advisory:

1. Farmers are advice to harvest the mature crop immediately and maintain suitable moisture content for safe storage; 2. Avoid to pesticide/weedicide spraying operation.

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Rice- Nursery- Thrips- Spray Thiamethoxam 25% WG 4g/20 cent in 40 litre of water.
RICE	Rice- Tillering Stage- 1. Leaf folder- Spray Acephate 75 % SP 400 ml (Or) Spray Azadirachtin 0.03% 400ml per acre in 200 litres of water; 2. Stem bore- Spray Azadirachtin 0.03% 400 ml (Or) Flubendiamide 20% WG 50g per acre in 200 litres of water.
JASMINE	Jasmine-Bud worm- Spray Thiocloprid 240 SC @ 1 ml /lit (or) Spray Spinosad 45 SC 0.5ml/lit.

### Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
-------------------------	--------------------------------

<b>Horticulture(Varieties)</b>	<b>Horticulture Specific Advisory</b>
CHILLI	Chilli- Damping Off- Treat the seeds with Trichoderma viride @ 4 g / kg or Pseudomonas fluorescens @ 10 g /kg of seeds treat with 24 hours before sowing.
BRINJAL	Brinjal- Shoot and fruit borer- Spray Azadirachtin 1.0% EC (10000 ppm) 3.0 ml/litre of water of chemical starting from one month after planting at 15 days interval. (or) Spray Emamectin benzoate 5 % SG 4 g/10 litre of water.