

## Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Tamil Nadu Agricultural University



Coimbatore, Tamil Nadu

# **Agromet Advisory Bulletin**

Date: 30-12-2022

Weather Forecast of District THE-NILGIRIS(Tamil Nadu) Issued On: 2022-12-30(Valid Till 08:30 IST of the next 5 days)

| Parameter              | 2022-12-31 | 2023-01-01 | 2023-01-02 | 2023-01-03 | 2023-01-04 |
|------------------------|------------|------------|------------|------------|------------|
| Rainfall(mm)           | 0.0        | 0.0        | 0.0        | 0.0        | 0.0        |
| Tmax(°C)               | 22.0       | 22.0       | 23.0       | 23.0       | 23.0       |
| Tmin(°C)               | 5.0        | 5.0        | 5.0        | 6.0        | 6.0        |
| RH-I(%)                | 90         | 90         | 90         | 80         | 80         |
| RH-II(%)               | 40         | 40         | 40         | 40         | 40         |
| Wind Speed(kmph)       | 6.0        | 4.0        | 4.0        | 6.0        | 6.0        |
| Wind Direction(Degree) | 90         | 50         | 90         | 90         | 70         |
| Cloud Cover(Octa)      | 2          | 1          | 2          | 2          | 2          |

#### **Weather Summary/Alert:**

As per District level forecast issued by the IMD for Nilgiris district, sky will be mainly clear. No rainfall is expected for next five days. Maximum temperature will be around 22-23°C and Minimum temperature will be around 5-6°C. Morning relative humidity will be 80-90% and evening relative humidity will be around 40%. Wind speed will be around 04-06 km per hour.

#### **General Advisory:**

To manage blister blight in tea crop, foliar spray of hexaconazole @ 200 ml + copper oxychloride @ 210 g/ha at 5 days interval or copper oxychloride @ 210 g + propiconazole @ 200 ml/ha at 10 days interval is recommended.

### **SMS Advisory:**

Soil application of Bacillus subtilis (BbV 57) or Pseudomonas fluorescens @ 10 g/vine is recommended for the management of root knot and reniform nematode population in black pepper.

## **Crop Specific Advisory:**

| Crop(Varieties) | Crop Specific Advisory   |  |  |  |
|-----------------|--|--|--|--|
| CARROT          | For carrot crop, soil application of neem cake @ 1 ton/ha during sowing, biofumigation of mustard plants followed by application of neem cake @ 250kg/ha along with Purpureocillium lilacinum @ 5kg/ha or application of P. lilacinum as seed treatment @ 20g/kg of seed followed by soil application @ 2.5kg/ha along with FYM @ 2.5 t/ha is recommended. |  |  |  |