

Agromet Advisory Bulletin for the Perambalur District (From 17th December 2022 to 21st December 2022) Issued jointly by Tamil Nadu Rice Research Institute -Aduthurai and IMD



Bulletin no: 100/2022

Date: 16.12.2022

Weather Summary for the period since the issue of last bulletin no. 99 (13.12.2022) for Tamil Nadu: Perambalur district

Maximum temperature of 30 °C - 31 °C, minimum temperature of 22 °C - 23 °C isolated rainfall were recorded in Perambalur district.

Past week Summary 13.12.2022 to 15.12.2022			Parameters	Ensemble weather forecast valid until 08.30 hrs of 21.12.2022				
Day-1 13/12	Day-2 14/12	Day-3 15/12	Date	Day-1 17/12	Day-2 18/12	Day-3 19/12	Day-4 20/12	Day-5 21/12
9.2	24.1	0.0	Rainfall (mm)	0	3	2	8	6
31	30	30	Max. Temp. (°C)	30	30	29	28	27
23	22	23	Min. Temp. (°C)	21	21	20	20	20
8	6	5	Cloud cover (Octa)	8	7	7	7	7
93-93	95-72	90-62	RH I-II (%)	90-50	90-50	90-60	90-60	90-80
7	11	9	Wind (kmph)	8	8	8	8	8
East	East	East	Wind direction	North East	North East	East	East	East

Weather Summary and forecast for Perambalur district

Weather forecast for next five days (17.12.2022 to 21.12.2022)

According to district level forecast issued by India Meteorological Department for Perambalur district, the sky will be moderately cloudy. Rainfall is expected on 18.12.2022 to 21.12.2022. Maximum temperature is expected to be around 27 °C - 30 °C and minimum temperature is expected to be around 20 °C - 21 °C. Morning relative humidity is expected to be around 90 percent and evening relative humidity is expected to be around 50-80 percent. Average wind speed is expected to be around 8 km per hour the wind direction will be east.

Сгор	Stage	Advisory			
		Postpone / avoid irrigation to all crops.			
Gei	neral	Provide adequate drainage facilities to lands liable to water			
		stagnation.			
		If leaf folder symptoms found (folded leaf, white scrape on leaf)			
	Main field	apply Carbosulfan 6% G 16.7 kg/ha or Cartaphydrochloride 50 %			
		SP 1000 g/ha as soil application or Beauveria bassiana 5 gram / 1			
		litter after rain cessation. Beauveria bassiana is available at Tamil			
		Nadu Rice Research Institute – Aduthurai.			
Rice		The prevailing weather condition is favourable for gall midge			
Rice		infection. It leads to the formation of onion like leaf and prevents			
		panicle formation. To control this apply chloripyriphos 20 EC 500			
		ml/ha or Beauveria bassiana 5 gram / 1 litter after rain cessation.			
		Sheath blight is being noticed in the paddy field due to prevailing			
		weather condition. To control this, spray carbendazim 200g/ac or			
		propiconozole 200 ml/acre in 200 liter water after rain cessation.			
	Cow and	The prevailing night time temperature is favorable for pneumonia			
		formation in cattle. To reduce the risk of pneumonia, cow, goat and			
Cattle	Goat	calves shouldn't be left out in open condition during night. A			
	Gout	suitable night time shelter with enough heating provision must be			
		given.			

SMS advisory

Postpone irrigation to all crops and provide adequate drainage facility to cropped fields wherever essential.

Director TRRI, Aduthurai