



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **ANDANALLUR** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	2.0	2.0	3.0	0.0	0.0
Tmax(°C)	31.0	30.0	30.0	31.0	30.0
Tmin(°C)	23.0	24.0	23.0	21.0	22.0
RH-I(%)	96	93	94	91	77
RH-II(%)	59	66	68	62	47
Wind Speed(kmph)	6.0	5.0	6.0	7.0	11.0
Wind Direction(Degree)	103	98	216	288	297
Cloud Cover(Octa)	6	8	6	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (90.0% & 60.0%), 30.0°C and 23.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 7.0 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **LALGUDI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	1.0	2.0	1.0	0.0	0.0
Tmax(°C)	32.0	31.0	31.0	33.0	32.0
Tmin(°C)	24.0	25.0	24.0	22.0	23.0
RH-I(%)	95	93	93	90	77
RH-II(%)	57	66	66	58	49
Wind Speed(kmph)	8.0	7.0	8.0	8.0	12.0
Wind Direction(Degree)	97	84	241	308	308
Cloud Cover(Octa)	6	8	6	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (90.0% & 60.0%), 30.0°C and 23.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 7.0 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **MANACHANELLUR** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	1.0	2.0	1.0	0.0	0.0
Tmax(°C)	32.0	31.0	31.0	33.0	32.0
Tmin(°C)	24.0	25.0	24.0	22.0	23.0
RH-I(%)	95	93	93	90	77
RH-II(%)	57	66	67	58	49
Wind Speed(kmph)	8.0	7.0	8.0	8.0	12.0
Wind Direction(Degree)	89	76	241	308	308
Cloud Cover(Octa)	6	8	6	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (90.0% & 60.0%), 30.0°C and 23.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 7.0 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **MANAPPARAI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	2.0	3.0	2.0	0.0	0.0
Tmax(°C)	32.0	31.0	31.0	32.0	31.0
Tmin(°C)	24.0	24.0	24.0	22.0	23.0
RH-I(%)	94	92	93	90	77
RH-II(%)	57	65	66	58	48
Wind Speed(kmph)	8.0	6.0	8.0	8.0	12.0
Wind Direction(Degree)	101	112	228	292	305
Cloud Cover(Octa)	6	8	6	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (89.0% & 57.0%), 32.0°C and 24.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 5.0 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **MANIKANDAM** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	2.0	2.0	1.0	0.0	0.0
Tmax(°C)	32.0	31.0	31.0	33.0	32.0
Tmin(°C)	24.0	25.0	24.0	22.0	23.0
RH-I(%)	95	93	93	90	77
RH-II(%)	58	67	67	59	49
Wind Speed(kmph)	8.0	7.0	8.0	8.0	12.0
Wind Direction(Degree)	62	50	234	310	307
Cloud Cover(Octa)	6	8	6	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (90.0% & 54.0%), 32.0°C and 23.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 4.8 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **MARUNGAPURI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	1.0	2.0	2.0	0.0	0.0
Tmax(°C)	32.0	31.0	31.0	33.0	31.0
Tmin(°C)	23.0	24.0	23.0	21.0	22.0
RH-I(%)	94	92	92	90	76
RH-II(%)	55	60	64	51	44
Wind Speed(kmph)	6.0	5.0	6.0	7.0	10.0
Wind Direction(Degree)	134	142	253	270	307
Cloud Cover(Octa)	6	8	7	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (90.0% & 54.0%), 32.0°C and 23.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 4.8 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **MUSIRI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	1.0	2.0	2.0	0.0	0.0
Tmax(°C)	32.0	31.0	31.0	33.0	31.0
Tmin(°C)	23.0	24.0	24.0	21.0	23.0
RH-I(%)	94	92	92	90	77
RH-II(%)	55	62	65	54	45
Wind Speed(kmph)	7.0	6.0	7.0	7.0	11.0
Wind Direction(Degree)	129	129	248	281	308
Cloud Cover(Octa)	6	8	6	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (90.0% & 57.0%), 32.0°C and 24.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 5.0 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **PULLAMBADI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	1.0	2.0	1.0	0.0	0.0
Tmax(°C)	32.0	31.0	31.0	33.0	31.0
Tmin(°C)	24.0	25.0	24.0	22.0	23.0
RH-I(%)	95	93	93	90	77
RH-II(%)	57	66	66	58	49
Wind Speed(kmph)	9.0	7.0	8.0	9.0	12.0
Wind Direction(Degree)	121	105	248	308	308
Cloud Cover(Octa)	6	8	6	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (89.0% & 56.0%), 32.0°C and 24.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 6.2 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **TATTAYANGARPETTAI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	1.0	3.0	1.0	0.0	0.0
Tmax(°C)	32.0	31.0	30.0	33.0	31.0
Tmin(°C)	23.0	24.0	23.0	21.0	22.0
RH-I(%)	93	92	92	90	77
RH-II(%)	55	60	64	50	44
Wind Speed(kmph)	6.0	6.0	7.0	7.0	10.0
Wind Direction(Degree)	163	157	246	264	313
Cloud Cover(Octa)	6	8	7	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (90.0% & 54.0%), 32.0°C and 23.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 4.8 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of THIRUVERAMBUR Block in TIRUCHIRAPPALLI(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	2.0	2.0	1.0	0.0	0.0
Tmax(°C)	32.0	31.0	31.0	33.0	32.0
Tmin(°C)	24.0	25.0	24.0	22.0	23.0
RH-I(%)	95	93	93	90	77
RH-II(%)	58	67	67	59	49
Wind Speed(kmph)	8.0	7.0	8.0	8.0	12.0
Wind Direction(Degree)	68	58	234	309	307
Cloud Cover(Octa)	6	8	6	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (89.0% & 57.0%), 32.0°C and 24.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 5.0 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **THOTTIAM** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	1.0	2.0	2.0	0.0	0.0
Tmax(°C)	32.0	31.0	31.0	33.0	31.0
Tmin(°C)	23.0	24.0	23.0	21.0	22.0
RH-I(%)	93	92	92	90	76
RH-II(%)	54	59	64	51	43
Wind Speed(kmph)	5.0	5.0	6.0	6.0	10.0
Wind Direction(Degree)	136	146	253	266	307
Cloud Cover(Octa)	6	8	7	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (90.0% & 54.0%), 32.0°C and 23.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 4.8 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **TURAIYUR** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	1.0	3.0	1.0	0.0	0.0
Tmax(°C)	32.0	31.0	30.0	32.0	31.0
Tmin(°C)	24.0	25.0	24.0	22.0	23.0
RH-I(%)	94	92	92	90	77
RH-II(%)	55	62	65	53	47
Wind Speed(kmph)	8.0	7.0	8.0	8.0	11.0
Wind Direction(Degree)	203	173	274	295	311
Cloud Cover(Octa)	6	8	7	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (89.0% & 57.0%), 32.0°C and 23.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 6.0 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **UPPILIAPURAM** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	1.0	3.0	1.0	0.0	0.0
Tmax(°C)	32.0	31.0	30.0	33.0	31.0
Tmin(°C)	23.0	24.0	23.0	22.0	23.0
RH-I(%)	94	92	92	90	77
RH-II(%)	55	61	64	51	45
Wind Speed(kmph)	7.0	7.0	7.0	7.0	10.0
Wind Direction(Degree)	197	176	259	274	314
Cloud Cover(Octa)	6	8	7	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (90.0% & 53.0%), 32.0°C and 23.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 6.0 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of **VAIYAMPATTI** Block in **TIRUCHIRAPPALLI**(Tamil Nadu) Issued On :2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall	2.0	2.0	2.0	0.0	0.0
Tmax(°C)	32.0	31.0	31.0	32.0	31.0
Tmin(°C)	23.0	24.0	23.0	21.0	22.0
RH-I(%)	95	92	93	90	77
RH-II(%)	57	65	66	59	47
Wind Speed(kmph)	7.0	6.0	7.0	7.0	12.0
Wind Direction(Degree)	105	112	228	288	301
Cloud Cover(Octa)	6	8	6	5	8

Weather Summary/Alert:

Rainfall is expected for 3 days in a week. Average relative humidity (morning & evening), maximum temperature and minimum temperature may be expected around (90.0% & 56.0%), 32.0°C and 23.0°C respectively. The rainy clouds may appear. Expected average and maximum wind speed respectively is 5.6 kmph and 20.0 kmph. The wind will be blow from North & North-East direction.

General Advisory:

Tamilnadu's average rainfall is likely to be normal during this week (07.12.2022 to 13.12.2022) From 2.93 mm to 4.37 mm.

SMS Advisory:

To know the weather forecast and agro advisory bulletins, download MEGHDOOT app and use it wisely.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to change in climate rice blast disease may occur in paddy crop. Hence, farmers are advised to do foliar spraying of Tricyclazole @ 2 gram / litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the prevailing weather condition the incense of Panama Wilt disease in Banana. Externally the first obvious signs of disease in most varieties are wilting and a light yellow colouring of the lower leaves, most prominent around the margins. They eventually turn a bright yellow colour with dead leaf margins. Splitting of pseudo stem base is a characteristic symptom. Practice proper crop rotation with paddy/sugarcane once or twice followed by banana for 2 to 3 cycles. Grow healthy plants with proper fertilization, irrigation and weed control. Provide good drainage. Apply Pseudomonas fluorescence @ 2 ½ kg/ha along with farmyard manure and neem cake.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.
COW	As the night temperature is decreasing, pneumonia is likely to occur. Therefore, goats, cows especially young calves should not be left in the open. Use heaters in sheds at night.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BIRD	Post mortem observations in the birds brought for diagnosis at Poultry Disease Diagnosis and Surveillance Laboratory, VC & RI, Namakkal revealed death of birds due to Wing rot infection and Necrotic enteritis. Hence, farmers are advised to test feed ingredients for Clostridia, Staphylococcus, and E.coli load and follow appropriate feeding management measures also, to use toxin binders in feed ingredients because of increased mycotoxin.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)(Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	To know the precaution of thunder and lightning, download DAMINI app and use it wisely.
GENERAL ADVICE	Moderate rainfall will be expected in the forthcoming days so, avoid and don't spray any chemicals to crop and better to spray chemicals at the time of no rain. Provide adequate drainage facility.