



# Agromet Advisory Bulletin for the **Tiruvarur** District

(From 05<sup>th</sup> February 2022 to 09<sup>th</sup> February 2022)

Issued jointly by Tamil Nadu Rice Research Institute -  
Aduthurai and IMD



**Bulletin no: 10/2022**

**Date: 04.02.2022**

Weather Summary for the period since the issue of last bulletin no. 09 (01.02.2022) for Tamil Nadu:

Maximum temperature of 31 °C to 32 °C and minimum temperature of 22 °C to 24 °C and were recorded in Tiruvarur district.

## Weather Summary and forecast for Tiruvarur district

Past week Summary 01.02.2022 to 03.02.2022			Parameters	Ensemble weather forecast valid until 08.30 hrs of 09.02.2022				
Day-1 01/02	Day-2 02/02	Day-3 03/02		Date	Day-1 05/02	Day-2 06/02	Day-3 07/02	Day-4 08/02
0	0	0	Rainfall (mm)	0	0	0	0	0
32	31	31	Max. Temp. (°C)	32	32	32	32	32
24	23	22	Mini. Temp. (°C)	23	23	23	23	23
4	4	4	Cloud cover (Octa)	4	4	4	6	8
78-66	78-68	70-64	RH I-II (%)	85-55	85-55	85-55	85-55	85-55
4	5	4	Wind (kmph)	6	8	8	10	10
East South East	East South East	South East	Wind direction	South East	East	East	East	East

## Weather forecast for next five days (05.02.2022 to 09.02.2022)

According to district level forecast issued by India Meteorological Department for Tiruvarur district, the sky will be moderately cloudy. No Rainfall expected next coming five days. Maximum temperature is expected to be around 32 °C and Minimum temperature is expected to be around 23 °C. Morning relative humidity is expected to be around 85 percent and evening relative humidity is expected to be around 55 percent. Average wind speed is expected to be around 6-10 km per hour and the wind direction will be east.

## Agro Advisory

Crop	Stage	Advisory
Rice		By using the current dry weather condition, harvested paddy seeds may be dried to 13% moisture content to prevent pest damage in storage.

Blackgram		Treat the black gram seeds with carbendazim 2g/1kg before sowing, to prevent disease incidence.
Rice		The prevailing weather condition conducive for Leaf Folder and Stemborer (white ear) occurrence in thaladi paddy. To control this, apply cartop hydrochloride 4G @ 6kg with sand or choloriphyriphos 400 ml/ac.
Cattle	Cow	Due to drop in night temperature, the livestock particularly the young ones should not be left in open. A proper night shelter with sufficient heating equipments (lighting bulb) must be provided to prevent the incidence of pneumonia.

### **SMS advisory**

- By using the current dry weather condition, harvested paddy seeds may be dried to 13% moisture content to prevent pest damage in storage.

**Director  
TRRI, Aduthurai**