



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **ANNAVASAL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	38.0	37.0	36.0	37.0	38.0
Tmin(°C)	29.0	29.0	28.0	29.0	29.0
RH-I(%)	56	56	56	57	57
RH-II(%)	34	37	37	35	35
Wind Speed(kmph)	15.0	14.0	15.0	12.0	11.0
Wind Direction(Degree)	243	243	239	252	286
Cloud Cover(Octa)	8	8	7	8	8

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 34.6°C and 37.5°C and the minimum temperature would be in between 25.0°C and 26.6°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 20 to 25 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut. Hence, farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and manage mental practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, where as roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **ARANTANGI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	38.0	37.0	36.0	37.0	38.0
Tmin(°C)	29.0	29.0	28.0	29.0	29.0
RH-I(%)	56	56	56	58	57
RH-II(%)	34	37	37	35	36
Wind Speed(kmph)	16.0	14.0	16.0	13.0	10.0
Wind Direction(Degree)	242	242	238	250	283
Cloud Cover(Octa)	8	8	7	8	8

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 37.7°C and 39.6°C and the minimum temperature would be in between 30.3°C and 31.1°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 09 to 18 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut. Hence, farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and manage mental practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, where as roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **ARIMALAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	38.0	37.0	36.0	37.0	38.0
Tmin(°C)	29.0	29.0	28.0	29.0	29.0
RH-I(%)	55	55	55	57	56
RH-II(%)	33	37	37	35	35
Wind Speed(kmph)	15.0	13.0	15.0	12.0	10.0
Wind Direction(Degree)	242	242	237	250	287
Cloud Cover(Octa)	8	8	7	8	8

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 34.9°C and 37.3°C and the minimum temperature would be in between 25.4°C and 26.7°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 11 to 18 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut. Hence, farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and manage mental practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, where as roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **AVADAIYARKOVIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	38.0	37.0	36.0	37.0	38.0
Tmin(°C)	29.0	29.0	28.0	29.0	29.0
RH-I(%)	56	56	57	58	57
RH-II(%)	34	37	37	35	36
Wind Speed(kmph)	17.0	15.0	16.0	14.0	10.0
Wind Direction(Degree)	242	242	237	249	281
Cloud Cover(Octa)	8	8	7	8	8

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 38.0°C and 37.3°C and the minimum temperature would be in between 29.8°C and 30.4°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 10 to 21 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut.Hence,farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and manage mental practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, where as roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **GANDARAVAKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	2.0
Tmax(°C)	37.0	36.0	36.0	36.0	38.0
Tmin(°C)	29.0	29.0	28.0	29.0	29.0
RH-I(%)	57	56	55	60	60
RH-II(%)	35	37	37	37	35
Wind Speed(kmph)	17.0	16.0	16.0	16.0	15.0
Wind Direction(Degree)	246	248	244	256	281
Cloud Cover(Octa)	8	8	7	7	7

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 35.6°C and 37.7°C and the minimum temperature would be in between 26.2°C and 26.4°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 27 to 33 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut. Hence, farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and manage mental practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, where as roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **KARAMBAKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	37.0	36.0	36.0	36.0	38.0
Tmin(°C)	29.0	29.0	28.0	29.0	29.0
RH-I(%)	57	56	56	59	59
RH-II(%)	34	37	37	36	35
Wind Speed(kmph)	17.0	15.0	16.0	15.0	13.0
Wind Direction(Degree)	245	246	242	254	282
Cloud Cover(Octa)	8	8	7	7	7

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 34.3°C and 37.7°C and the minimum temperature would be in between 25.9°C and 26.7°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 18 to 23 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut. Hence, farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and manage mental practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, where as roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **KUNNANDARKOIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	37.0	36.0	36.0	36.0	38.0
Tmin(°C)	28.0	28.0	27.0	28.0	28.0
RH-I(%)	58	59	58	60	59
RH-II(%)	35	38	37	36	35
Wind Speed(kmph)	18.0	16.0	17.0	15.0	15.0
Wind Direction(Degree)	248	248	247	258	280
Cloud Cover(Octa)	8	8	8	7	7

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 36.0°C and 38.0° and the minimum temperature would be in between 27.0°C and 28.0°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 15 to 17 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut.Hence,farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and management practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, whereas roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **MANALMELKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	37.0	36.0	36.0	37.0	38.0
Tmin(°C)	29.0	29.0	28.0	29.0	29.0
RH-I(%)	57	57	57	59	58
RH-II(%)	34	37	37	35	36
Wind Speed(kmph)	17.0	16.0	17.0	15.0	11.0
Wind Direction(Degree)	242	242	238	248	278
Cloud Cover(Octa)	8	8	7	8	8

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 36.0°C and 38.0° and the minimum temperature would be in between 28.0°C and 29.0°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 11 to 17 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut.Hence,farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and manage mental practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, where as roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **PONNAMARAVATI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	38.0	37.0	36.0	37.0	38.0
Tmin(°C)	29.0	29.0	28.0	29.0	29.0
RH-I(%)	56	56	56	58	57
RH-II(%)	34	37	37	35	36
Wind Speed(kmph)	16.0	15.0	16.0	13.0	10.0
Wind Direction(Degree)	243	242	238	250	283
Cloud Cover(Octa)	8	8	7	8	8

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 36.0°C and 38.0° and the minimum temperature would be in between 28.0°C and 29.0°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 10 to 16 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut.Hence,farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and manage mental practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, where as roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **PUDUKKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	38.0	37.0	36.0	37.0	38.0
Tmin(°C)	29.0	29.0	28.0	29.0	29.0
RH-I(%)	55	55	55	56	56
RH-II(%)	33	37	37	35	35
Wind Speed(kmph)	14.0	12.0	14.0	11.0	9.0
Wind Direction(Degree)	241	241	236	250	289
Cloud Cover(Octa)	8	8	7	8	8

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 36.0°C and 38.0°C and the minimum temperature would be in between 28.0°C and 29.0°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 09 to 14 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut.Hence,farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and manage mental practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, where as roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **THIRUVARANGULAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	38.0	37.0	36.0	37.0	38.0
Tmin(°C)	29.0	29.0	28.0	29.0	29.0
RH-I(%)	55	55	55	56	56
RH-II(%)	33	37	37	35	35
Wind Speed(kmph)	14.0	12.0	14.0	11.0	9.0
Wind Direction(Degree)	241	241	236	250	289
Cloud Cover(Octa)	8	8	7	8	8

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 36.0°C and 38.0° and the minimum temperature would be in between 28.0°C and 29.0°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 09 to 14 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut.Hence,farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and manage mental practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, where as roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **TIRUMAYAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	38.0	37.0	36.0	37.0	38.0
Tmin(°C)	29.0	29.0	28.0	29.0	29.0
RH-I(%)	56	56	56	57	57
RH-II(%)	33	37	37	35	35
Wind Speed(kmph)	15.0	13.0	15.0	12.0	10.0
Wind Direction(Degree)	242	242	237	250	286
Cloud Cover(Octa)	8	8	7	8	8

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 36.0°C and 38.0° and the minimum temperature would be in between 28.0°C and 29.0°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 10 to 15 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut.Hence,farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and manage mental practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, where as roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 12-07-2022

Weather Forecast of **VIRALIMALAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-13	2022-07-14	2022-07-15	2022-07-16	2022-07-17
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	37.0	36.0	36.0	36.0	38.0
Tmin(°C)	28.0	28.0	27.0	28.0	28.0
RH-I(%)	59	59	58	60	59
RH-II(%)	35	38	37	36	35
Wind Speed(kmph)	18.0	17.0	17.0	15.0	15.0
Wind Direction(Degree)	248	249	248	258	279
Cloud Cover(Octa)	8	8	8	7	7

Weather Summary/Alert:

Chances of Dry weather next five days. Maximum temperature would be in the range 36.0°C and 38.0° and the minimum temperature would be in between 27.0°C and 28.0°C . Cloudy sky is expected for next 5 days. Wind speed is expected to be 15 to 18 kmph.

General Advisory:

Before the transplanting of tomato farmers are requested to apply 10 tonne well decomposed farmyard manure as basal dose during the time of last ploughing per acre.

SMS Advisory:

In kuruvai paddy one hand weeding is to be given on 20 to 25 days after transplanting.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
MAIZE	Can cultivate Maize Adipattam Season (July - August) irrigated maize sowing may be done with imidacloprid 70 WS seed treatment @ 10g/kg of seeds.
RICE	The prevailing weather condition favours the stem borer incidence in paddy. Place pheromone traps @ 4 / ac to control stem borer population. Also to control this, spray profenophos at 400 ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for effective control of stemborer.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	Prevailing dry weather is favourable for sucking pest multiplication in tomato crop. Hence monitor the pest load and spray fish oil rosin soap @ 25 g/litre of water.
COCONUT	Prevailing hot weather may be favorable for spiralling whitefly in coconut.Hence,farmers are requested to monitor the incidence and fix yellow sticky traps in 5-6 places/ acre.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
-----------------------	------------------------------

Live Stock(Varieties)	Live Stock Specific Advisory
COW	To get high milk yield during any lactation, the milch animal should be properly fed and necessary care and manage mental practices should be followed. Provide green succulent forage together with leguminous hay or straw to the extent of animal can consume, so that all its maintenance requirements met through forage only. Extra concentrate at the rate of 1 kg should be provided for every 2 to 2.5 litres of milk yield. Salt and mineral supplements should be given to maintain the lactation. Never frighten or excite the animals. Always treat them gently and with kindness. Keep up regularity of feeding. Concentrate mix is fed before milking, where as roughages to be fed after milking.
GOAT	Goat rearing farmers are advised to prevent the goat from Tapeworm, which leads to mortality and weight loss. Goats have to be de-wormed from July to September, if not yet done, contact the nearby veterinarian and provide deworm medicines to your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Farmers are requested to take up planting operations, especially vegetable crops like chilies, brinjal, gourd, banana and perennial crops.
GENERAL ADVICE	During adipattam farmers can soak seed in one percent potassium chloride solution for 6 hours one day before sowing of quality seeds and leave to dry for 16 hours in the shade before sowing. Through this seed hardening treatment the young seedlings get uniform germination capacity and growth and the seedlings emerge and develop into active seedlings.