



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **ANNAVASAL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	1.0	0.0	1.0	1.0
Tmax(°C)	38.0	38.0	38.0	38.0	38.0
Tmin(°C)	29.0	29.0	29.0	29.0	29.0
RH-I(%)	57	60	59	59	59
RH-II(%)	35	37	34	34	34
Wind Speed(kmph)	15.0	13.0	13.0	14.0	14.0
Wind Direction(Degree)	286	285	268	270	270
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of dry weather next five days. Maximum temperature would be in the range 38.0°C and the minimum temperature would be in the range 29.0-°C . Partly cloudy sky will appear . Wind speed is expected to be 13- 15 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
JASMINE	Rose , Jasmine and Tuberose are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconzole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures / pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **ARANTANGI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	1.0	0.0	1.0	1.0
Tmax(°C)	38.0	38.0	38.0	38.0	38.0
Tmin(°C)	30.0	29.0	30.0	30.0	30.0
RH-I(%)	57	60	59	58	58
RH-II(%)	36	37	34	34	34
Wind Speed(kmph)	14.0	13.0	13.0	14.0	14.0
Wind Direction(Degree)	282	283	266	268	268
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of dry weather next five days. Maximum temperature would be in the range 38.0°C and the minimum temperature would be in the range 29.0-°C . Partly cloudy sky will appear . Wind speed is expected to be 13- 14 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
JASMINE	Rose , Jasmine and Tuberose are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconzole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **ARIMALAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	1.0	0.0	1.0	1.0
Tmax(°C)	38.0	38.0	38.0	38.0	38.0
Tmin(°C)	30.0	29.0	30.0	30.0	30.0
RH-I(%)	57	59	59	58	58
RH-II(%)	35	37	34	34	34
Wind Speed(kmph)	14.0	12.0	12.0	13.0	13.0
Wind Direction(Degree)	287	286	267	269	269
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of dry weather next five days. Maximum temperature would be in the range 38.0°C and the minimum temperature would be in the range 29.0°C-30.0°C . Partly cloudy sky will appear . Wind speed is expected to be 12- 14 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
JASMINE	Rose , Jasmine and Tuberose are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconazole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures / pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **AVADAIYARKOVIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	2.0	0.0	0.0	0.0
Tmax(°C)	38.0	38.0	38.0	38.0	38.0
Tmin(°C)	29.0	29.0	29.0	29.0	29.0
RH-I(%)	58	60	59	58	58
RH-II(%)	36	37	34	35	35
Wind Speed(kmph)	14.0	13.0	13.0	14.0	14.0
Wind Direction(Degree)	278	282	264	266	266
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain on 10 July. Maximum temperature would be in the range 38.0°C and the minimum temperature would be in the range 29.0°C . Partly cloudy sky will appear . Wind speed is expected to be 13- 14 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
JASMINE	Rose , Jasmine and Tuberose are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconzole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **GANDARAVAKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	1.0	0.0	3.0	3.0
Tmax(°C)	38.0	38.0	38.0	38.0	38.0
Tmin(°C)	29.0	29.0	29.0	29.0	29.0
RH-I(%)	59	63	61	61	61
RH-II(%)	35	35	33	35	35
Wind Speed(kmph)	20.0	16.0	16.0	18.0	18.0
Wind Direction(Degree)	284	282	273	274	274
Cloud Cover(Octa)	8	7	8	8	8

Weather Summary/Alert:

Chances of light rain on 12 and 13 July. Maximum temperature would be in the range 38.0°C and the minimum temperature would be in the range 29.0°C. Partly cloudy sky will appear. Wind speed is expected to be 16- 20 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
JASMINE	Rose , Jasmine and Tuberose are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconazole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **KARAMBAKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	1.0	0.0	2.0	2.0
Tmax(°C)	38.0	38.0	38.0	38.0	38.0
Tmin(°C)	29.0	29.0	29.0	29.0	29.0
RH-I(%)	58	61	60	60	60
RH-II(%)	35	36	34	35	35
Wind Speed(kmph)	17.0	15.0	15.0	16.0	16.0
Wind Direction(Degree)	283	283	270	271	271
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of very light rain on 12 and 13 July. Maximum temperature would be in the range 38.0°C and the minimum temperature would be in the range 29.0°C . Partly cloudy sky will appear . Wind speed is expected to be 16- 17 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
JASMINE	Rose , Jasmine and Tuberose are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconazole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **KUNNANDARKOIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	1.0	0.0	0.0	0.0
Tmax(°C)	37.0	38.0	38.0	38.0	38.0
Tmin(°C)	28.0	28.0	28.0	28.0	28.0
RH-I(%)	60	61	60	60	60
RH-II(%)	36	36	33	35	35
Wind Speed(kmph)	17.0	16.0	17.0	18.0	18.0
Wind Direction(Degree)	280	281	271	272	272
Cloud Cover(Octa)	8	8	7	8	8

Weather Summary/Alert:

Chances of dry weather next five days. Maximum temperature would be in the range 37.0°C- 38.0°C and the minimum temperature would be in the range 28.0°C Partly cloudy sky will appear . Wind speed is expected to be 16 - 18 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
JASMINE	Rose , Jasmine and Tuberose are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconazole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **MANALMELKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	2.0	1.0	0.0	0.0
Tmax(°C)	38.0	38.0	38.0	38.0	38.0
Tmin(°C)	29.0	29.0	29.0	29.0	29.0
RH-I(%)	58	60	60	58	58
RH-II(%)	36	37	34	35	35
Wind Speed(kmph)	14.0	13.0	14.0	15.0	15.0
Wind Direction(Degree)	275	280	264	265	265
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain on 10 July. Maximum temperature would be in the range 38.0°C and the minimum temperature would be in the range 29.0°C Partly cloudy sky will appear . Wind speed is expected to be 13 - 14 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
JASMINE	Rose , Jasmine and Tuberose are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconazole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures / pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **PONNAMARAVATI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	2.0	0.0	0.0	0.0
Tmax(°C)	38.0	38.0	38.0	38.0	38.0
Tmin(°C)	29.0	29.0	29.0	29.0	29.0
RH-I(%)	58	60	59	58	58
RH-II(%)	36	37	34	34	34
Wind Speed(kmph)	14.0	13.0	13.0	14.0	14.0
Wind Direction(Degree)	280	283	266	268	268
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light rain on 10 July. Maximum temperature would be in the range 38.0°C and the minimum temperature would be in the range 29.0°C Partly cloudy sky will appear . Wind speed is expected to be 13 - 14 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
JASMINE	Rose , Jasmine and Tuberose are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconazole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **PUDUKKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	1.0	0.0	1.0	1.0
Tmax(°C)	38.0	38.0	38.0	38.0	38.0
Tmin(°C)	30.0	29.0	30.0	30.0	30.0
RH-I(%)	56	59	59	58	58
RH-II(%)	35	37	34	34	34
Wind Speed(kmph)	14.0	12.0	11.0	12.0	12.0
Wind Direction(Degree)	290	287	267	269	269
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of dry weather next five days. Maximum temperature would be in the range 38.0°C and the minimum temperature would be in the range 29.0°C - 30.0°C. Partly cloudy sky will appear . Wind speed is expected to be 12 - 14 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
JASMINE	Rose , Jasmine and Tuberose are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconazole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **THIRUVARANGULAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	1.0	0.0	1.0	1.0
Tmax(°C)	38.0	38.0	38.0	38.0	38.0
Tmin(°C)	30.0	29.0	30.0	30.0	30.0
RH-I(%)	56	59	59	58	58
RH-II(%)	35	37	34	34	34
Wind Speed(kmph)	14.0	12.0	11.0	12.0	12.0
Wind Direction(Degree)	289	287	267	269	269
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of dry weather next five days. Maximum temperature would be in the range 38.0°C and the minimum temperature would be in the range 29.0°C - 30.0°C. Partly cloudy sky will appear . Wind speed is expected to be 12 - 14 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
JASMINE	Rose , Jasmine and Tuberose are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconazole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures / pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **TIRUMAYAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	1.0	0.0	1.0	1.0
Tmax(°C)	38.0	38.0	38.0	38.0	38.0
Tmin(°C)	30.0	29.0	30.0	30.0	30.0
RH-I(%)	57	59	59	58	58
RH-II(%)	35	37	34	34	34
Wind Speed(kmph)	14.0	12.0	12.0	13.0	13.0
Wind Direction(Degree)	286	285	266	268	268
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of dry weather next five days. Maximum temperature would be in the range 38.0°C and the minimum temperature would be in the range 29.0°C - 30.0°C. Partly cloudy sky will appear . Wind speed is expected to be 12 - 14 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
JASMINE	Rose , Jasmine and Tuberose are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconazole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 08-07-2022

Weather Forecast of **VIRALIMALAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-07-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-07-09	2022-07-10	2022-07-11	2022-07-12	2022-07-13
Rainfall	0.0	1.0	0.0	0.0	0.0
Tmax(°C)	37.0	38.0	38.0	38.0	38.0
Tmin(°C)	28.0	28.0	28.0	28.0	28.0
RH-I(%)	60	62	60	60	60
RH-II(%)	36	36	33	35	35
Wind Speed(kmph)	17.0	17.0	17.0	18.0	18.0
Wind Direction(Degree)	278	280	271	272	272
Cloud Cover(Octa)	8	8	7	8	8

Weather Summary/Alert:

Chances of dry weather next five days. Maximum temperature would be in the range 38.0°C and the minimum temperature would be in the range 28.0°C. Partly cloudy sky will appear . Wind speed is expected to be 17 - 18 kmph.

General Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS Advisory:

Inclusion of Azolla in feed of livestock and poultry reduces the feed cost as well as improves the production.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JASMINE	Rose , Jasmine and Tuberoses are all affected by the sucking Pest All stages . To control sucking pest spraying of prophenophos @ 2 ml per litre of water is recommended.
MAIZE	Farmers Can cultivate Maize during Adipattam (July - August). TNAU Maize hybrid Co 6 in suitable for this Before sowing Seed treatment with imidacloprid 70 ws at the rate of 10 g / kg seed and treat the seed and then sow.
RICE	Due to prevailing climatic conditions, stem borer is seen kuruvai season Paddy. Spraying 3.03% of Azadirachtin 1000 ml /ha or Carbofuran 3% CG @ 25kg /ha in recommended.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	To avoid damping off disease or soil borne diseases in tomato Trichoderma viride or Pseudomonas fluorescens may be applied @ 2.5 kg per hectare.
BRINJAL	This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Trichoderma viride @ 4 g / kg of seed before sowing to avoid rot in seedlings.

Horticulture(Varieties)	Horticulture Specific Advisory
COCONUT	Due to prevailing climatic conditions there will be possibility of Thanjavur Wilt. The progress of the disease is faster during July to November. Symptoms: Initial symptoms of Thanjavur Wilt start with yellowing and drooping of the outer whorl of leaves. This is followed by exudation of reddish brown liquid through cracks at the base of the trunk and oozing spread upwards. Decaying of tissues at bleeding point and rotting of the basal portion of the stem . Management : Aurefungin –sol @+ copper sulphate @ 1 g dissolved 100 ml water or hexaconzole @ 2 ml with 100 ml of water . applied as root feeding for 3 times at 3 months interval.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Farmers are advised to deworm their goat and sheep to avoid the endoparasites. In goat and sheep, to avoid dehydration loss provide salt lick

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Wind with the speed of 14 kmph is expected for the next five days. Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya, Guava and Moringa, to prevent from lodging. For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support. Postpone the spraying activity