



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **ANNAVASAL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	3.0	5.0	0.0	0.0	0.0
Tmax(°C)	39.0	39.0	39.0	39.0	38.0
Tmin(°C)	29.0	30.0	30.0	29.0	30.0
RH-I(%)	65	63	61	64	63
RH-II(%)	32	34	32	34	38
Wind Speed(kmph)	12.0	15.0	13.0	14.0	11.0
Wind Direction(Degree)	275	272	233	216	246
Cloud Cover(Octa)	7	8	6	5	6

Weather Summary/Alert:

Chances of light rain on 01 June .Maximum temperature would be in the range 37.8°C and 39.4°C and the minimum temperature would be in the range 25.7°C and 27.5°C . Partly cloudy sky will appear . Wind speed is expected to be 11 - 18 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain end. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **ARANTANGI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	2.0	4.0	0.0	0.0	0.0
Tmax(°C)	39.0	39.0	39.0	39.0	38.0
Tmin(°C)	30.0	30.0	30.0	29.0	30.0
RH-I(%)	64	62	61	64	63
RH-II(%)	33	34	32	35	39
Wind Speed(kmph)	12.0	15.0	13.0	14.0	11.0
Wind Direction(Degree)	272	269	231	213	241
Cloud Cover(Octa)	7	8	6	5	6

Weather Summary/Alert:

Chances of light rain on 02 June .Maximum temperature would be in the range 38.1°C and 40.4°C and the minimum temperature would be in the range 31.1°C and 31.9°C . Partly cloudy sky will appear . Wind speed is expected to be 15 - 23 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain end. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **ARIMALAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	2.0	4.0	0.0	0.0	0.0
Tmax(°C)	39.0	39.0	39.0	39.0	38.0
Tmin(°C)	30.0	30.0	30.0	29.0	30.0
RH-I(%)	65	62	60	63	63
RH-II(%)	33	34	32	34	39
Wind Speed(kmph)	11.0	14.0	12.0	14.0	11.0
Wind Direction(Degree)	276	270	230	210	242
Cloud Cover(Octa)	7	8	6	5	6

Weather Summary/Alert:

Chances of drizzle on 02 June. Maximum temperature would be in the range 35.6°C and 38.3°C and the minimum temperature would be in the range 26.4°C and 27.5°C. Partly cloudy sky will appear. Wind speed is expected to be 11 - 19 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain end. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **AVADAIYARKOVIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	2.0	4.0	0.0	0.0	0.0
Tmax(°C)	39.0	39.0	39.0	39.0	38.0
Tmin(°C)	30.0	30.0	30.0	29.0	30.0
RH-I(%)	64	62	60	64	63
RH-II(%)	33	35	32	35	39
Wind Speed(kmph)	12.0	15.0	13.0	15.0	12.0
Wind Direction(Degree)	269	266	231	214	240
Cloud Cover(Octa)	7	8	6	5	6

Weather Summary/Alert:

Chances of light rain on 02 June .Maximum temperature would be in the range 36.8°C and 39.3°C and the minimum temperature would be in the range 30.2°C and 31.1°C . Partly cloudy sky will appear . Wind speed is expected to be 19 - 25 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain end. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **GANDARAVAKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	3.0	5.0	1.0	0.0	0.0
Tmax(°C)	39.0	39.0	39.0	39.0	38.0
Tmin(°C)	29.0	29.0	30.0	29.0	30.0
RH-I(%)	67	66	65	66	66
RH-II(%)	32	35	33	34	39
Wind Speed(kmph)	15.0	18.0	18.0	17.0	13.0
Wind Direction(Degree)	269	275	235	223	241
Cloud Cover(Octa)	7	8	6	4	6

Weather Summary/Alert:

Chances of light rain on 01 June .Maximum temperature would be in the range 37.6°C and 39.3°C and the minimum temperature would be in the range 26.5°C and 28.2°C . Partly cloudy sky will appear . Wind speed is expected to be 17 - 20 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain end. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **KARAMBAKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	3.0	5.0	0.0	0.0	0.0
Tmax(°C)	39.0	39.0	39.0	39.0	38.0
Tmin(°C)	29.0	29.0	30.0	29.0	30.0
RH-I(%)	66	64	63	65	65
RH-II(%)	32	34	32	34	38
Wind Speed(kmph)	14.0	17.0	15.0	16.0	12.0
Wind Direction(Degree)	271	273	235	220	244
Cloud Cover(Octa)	7	8	6	5	6

Weather Summary/Alert:

Chances of drizzle on 01 and 03 June. Maximum temperature would be in the range 36.6°C and 38.0°C and the minimum temperature would be in the range 26.8°C and 27.8°C. Partly cloudy sky will appear. Wind speed is expected to be 15 - 19 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain end. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **KUNNANDARKOIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	3.0	5.0	1.0	0.0	3.0
Tmax(°C)	39.0	39.0	39.0	39.0	38.0
Tmin(°C)	28.0	29.0	28.0	28.0	29.0
RH-I(%)	66	64	63	65	64
RH-II(%)	31	33	31	32	36
Wind Speed(kmph)	16.0	18.0	15.0	14.0	13.0
Wind Direction(Degree)	273	276	244	237	259
Cloud Cover(Octa)	8	7	6	5	6

Weather Summary/Alert:

Chances of light rain on 01, 02 and 03 June. Maximum temperature would be in the range 38.0°C and 39.0°C and the minimum temperature would be in the range 28.0°C and 29.0°C. Partly cloudy sky will appear. Wind speed is expected to be 13 - 18 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain ends. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **MANALMELKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	2.0	4.0	0.0	0.0	0.0
Tmax(°C)	38.0	38.0	39.0	39.0	38.0
Tmin(°C)	29.0	30.0	29.0	29.0	30.0
RH-I(%)	64	63	61	64	64
RH-II(%)	33	35	33	36	39
Wind Speed(kmph)	12.0	15.0	14.0	15.0	13.0
Wind Direction(Degree)	267	265	232	216	239
Cloud Cover(Octa)	7	8	6	5	6

Weather Summary/Alert:

Chances of light rain on 01 and 02 June. Maximum temperature would be in the range 38.0°C and 39.0°C and the minimum temperature would be in the range 29.0°C and 30.0°C. Partly cloudy sky will appear. Wind speed is expected to be 12 - 15 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain end. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **PONNAMARAVATI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	2.0	4.0	0.0	0.0	0.0
Tmax(°C)	39.0	39.0	39.0	39.0	38.0
Tmin(°C)	29.0	30.0	29.0	29.0	30.0
RH-I(%)	64	62	60	63	63
RH-II(%)	32	34	32	34	38
Wind Speed(kmph)	12.0	15.0	13.0	14.0	11.0
Wind Direction(Degree)	272	268	232	215	243
Cloud Cover(Octa)	7	8	6	5	6

Weather Summary/Alert:

Chances of light rain on 01 and 02 June. Maximum temperature would be in the range 38.0°C and 39.0°C and the minimum temperature would be in the range 29.0°C and 30.0°C. Partly cloudy sky will appear. Wind speed is expected to be 11 - 15 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain end. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **PUDUKKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	2.0	5.0	0.0	0.0	0.0
Tmax(°C)	39.0	39.0	39.0	39.0	38.0
Tmin(°C)	30.0	30.0	30.0	29.0	30.0
RH-I(%)	65	62	60	63	63
RH-II(%)	33	34	32	34	39
Wind Speed(kmph)	11.0	14.0	11.0	14.0	10.0
Wind Direction(Degree)	278	271	228	207	241
Cloud Cover(Octa)	7	8	6	5	6

Weather Summary/Alert:

Chances of light rain on 01 and 02 June. Maximum temperature would be in the range 38.0°C and 39.0°C and the minimum temperature would be in the range 29.0°C and 30.0°C. Partly cloudy sky will appear. Wind speed is expected to be 10 - 14 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain end. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **THIRUVARANGULAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	2.0	5.0	0.0	0.0	0.0
Tmax(°C)	39.0	39.0	39.0	39.0	38.0
Tmin(°C)	30.0	30.0	30.0	29.0	30.0
RH-I(%)	65	62	60	63	63
RH-II(%)	33	34	32	34	39
Wind Speed(kmph)	11.0	14.0	11.0	14.0	10.0
Wind Direction(Degree)	278	271	228	208	241
Cloud Cover(Octa)	7	8	6	5	6

Weather Summary/Alert:

Chances of light rain on 01 and 02 June. Maximum temperature would be in the range 38.0°C and 39.0°C and the minimum temperature would be in the range 29.0°C and 30.0°C. Partly cloudy sky will appear. Wind speed is expected to be 11 - 14 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain end. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with <i>Trichoderma viride</i> @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **TIRUMAYAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	2.0	4.0	0.0	0.0	0.0
Tmax(°C)	39.0	39.0	39.0	39.0	38.0
Tmin(°C)	30.0	30.0	30.0	29.0	30.0
RH-I(%)	65	62	60	63	63
RH-II(%)	33	34	32	34	39
Wind Speed(kmph)	11.0	14.0	12.0	14.0	11.0
Wind Direction(Degree)	275	270	230	211	242
Cloud Cover(Octa)	7	8	6	5	6

Weather Summary/Alert:

Chances of light rain on 01 and 02 June. Maximum temperature would be in the range 38.0°C and 39.0°C and the minimum temperature would be in the range 29.0°C and 30.0°C. Partly cloudy sky will appear. Wind speed is expected to be 11 - 14 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain end. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 31-05-2022

Weather Forecast of **VIRALIMALAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-05-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-06-01	2022-06-02	2022-06-03	2022-06-04	2022-06-05
Rainfall	3.0	5.0	1.0	0.0	3.0
Tmax(°C)	39.0	39.0	39.0	39.0	38.0
Tmin(°C)	28.0	29.0	28.0	28.0	29.0
RH-I(%)	65	64	62	65	64
RH-II(%)	31	33	31	32	36
Wind Speed(kmph)	16.0	18.0	16.0	14.0	13.0
Wind Direction(Degree)	272	276	245	239	260
Cloud Cover(Octa)	8	7	6	5	6

Weather Summary/Alert:

Maximum temperature would be in the range 38.0°C and 39.0°C and the minimum temperature would be in the range 28.0°C and 29.0°C . Partly cloudy sky will appear . Wind speed is expected to be 13 - 18 kmph

General Advisory:

As 10 to 15 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.

SMS Advisory:

Preparatory cultivation can be started for planting of paddy and horticultural crops such as vegetables, coconut etc.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Considering the anticipated rainfall nursery land preparation may be done for short duration rice. But postpone the sowing till the rain end. Rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 are suitable for kuruvai season.
RICE	The application of Biofertilizer in Paddy will give good results. Raise Azolla as a dual crop by inoculating 250kg/ha 3 to 5 DAT and then incorporating it during weeding for the wet season crop. And broadcast 10kg of soil-based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
MAIZE	Application of micronutrient will rectify the nutrient deficiency in Maize Cultivation. Farmers who have planted maize recently can apply 5kg of Millet nutrient mixture or 10Kg of ZnSo ₄ /acre. Also apply Farm Yard Manure along with 2kgs of Azospirillum and 2kgs of Phosphobacteria per acre.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	June is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg. The seed treatment must be done 24 hours before sowing of seeds.
COCONUT	Hook out the beetles using a long iron rod and kill them at the time of harvest. Apply 3 naphthalene balls/palm with 100g sand at the crown of the tree once in 45 days to control the Rhinoceros Beetle.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
GOAT	Due to the prevailing weather condition the Enterotoxaemia disease may affect the Goats. The main symptom of the disease causes sudden death in young growing kids. Mucous diarrhea may also seen during death. To prevent this disease, do not feed on young grass, allow the goat for grazing one hour after the sunshine. Vaccinate the animals once in a year before the onset of monsoon.
COW	The greens in pasture returned due to recent summer showers. As these greens may remain for 7 to 10 days, farmers are advised to make use of it for their milch animals and other livestocks.