


## Tamil Nadu Agricultural University

&





Issued jointly by Agro Climate Research Centre, TNAU and IMD

Weather Summary for the period since the issue of last bulletin 049 (21.06.2022) for Tamil Nadu. Maximum temperature ranged from 33°C to 38°C and minimum temperature ranged from 23°C to 27°C, over the State. Kodaikanal and Uthagamandalam recorded maximum temperature ranged from 18°C to 21°C and minimum temperature ranged from 10°C to 11°C. During this period isolated rainfall was recorded in Tamil Nadu.

### **Weather Summary for Erode district**

Past week Summary						Ensemble weather forecast valid until				
(19.06.2022 to 23.06.2022)						08.30 hrs of 29.06.2022				
Day-1 19/06	Day-2 20/06	Day-3 21/06	Day-4 22/06	Day-5 23/06	Date	Day-1 25/06	Day-2 26/06	Day-3 27/06	Day-4 28/06	Day-5 29/06
0.1	0.6	0.3	0.0	0.1	Rainfall	3	4	0	0	3
34	34	34	31	31	Max. Temp. (°C)	34	34	34	34	34
23	23	23	23	23	Mini. Temp. (°C)	23	23	23	23	23
6	6	7	7	7	Cloud cover (Octa)	5	5	4	4	5
76	82	83	79	73	RH morning (%)	70	70	70	70	70
65	60	69	65	71	RH evening (%)	60	60	60	60	60
19	16	21	19	20	Wind (kmph)	8	8	8	8	8
230	230	200	230	230	Wind Direction	230	250	250	250	270

#### Weather forecast for next five days: (25.06.2022 to 29.06.2022)

According to district forecast issued by the India Meteorological Department for Erode district, sky will be mostly cloudy. Light rainfall is expected on 25.06.22, 26.06.22 & 29.06.22. Maximum temperature is expected to be around 34°C. Minimum temperature is expected to be around 23°C. Morning relative humidity is expected to be around 70 per cent and evening relative humidity is expected to be around 60 per cent. Average wind speed is expected to be around 8 km per hour and the wind direction will be from South West direction.

# Agro Advisory

Crop	Stage	Advisory
General		Light rainfall along with wind speed @ 8-16 kmph is expected during this week.
		Irrigation may be postponed to the crops based on soil moisture.
		By utilizing the rainfall planting of tree crops may be initiated in pits dug and cured for a week.
Rice	Sowing	Light rainfall is expected and hence maintain thin film of water to avoid seed displacement in nursery and seedling displacement in recently planted main field.
		Postpone the transplanting of rice, since chances for uprooting of transplanted seedlings due to rain.
		The expected temperature is favourable for sucking pest incidence in the rice nursery field. Hence, monitor for the pest infestation and if needed spray 5% neem seed kernel extract.
Maize	Sowing	By utilizing the rainfall, irrigated maize and sunflower may be sowing as June is best season for sowing.  In non-intercropped Maize crops, apply pre emergence weedicide
		Atrazine @ 400g/acre in 200 liters of water, within 3 days after sowing.
Small millets		Considering the anticipated rainfall sowing of small millets may be taken up in the foothills of western ghats areas.
Fodder Sorghum	Vegetative	By utilizing expected rainfall, topdressing with nitrogen to fodder sorghum is advised.
Gourd	Land preparation	By utilizing the last week rain and anticipated rain during this week, gourds vegetables may be sown in pits.
Banana		Even though average wind speed expected is 8 kmph. However instantaneous wind speed may be exceeding 16 kmph. Hence farmers are requested to provide propping to the banana crop which is more than 8 months old.
Turmeric		Since rainfall is expected, irrigation may be postponed to all irrigated crops, especially turmeric and provide drainage.
Coconut		By utilizing rainfall, apply fertilizer for coconut @ 650 gm urea, 1kg SuPo4, 1kg MOP and 2kg Neem cake along with 50kg FYM.  Provide inward basin in around the coconut trees to harvest rainwater in
		root zone.

Animal	Matitis disease may come to milch animals. Clean the under with 1 % potassium permanganate solution, before and after milking.
	Since Wind speed of 8 km is expected along with rainfall, provide adequate care to the animal sheds.
	Do not allow the animals to graze on newly grown grasses, due to the receipt of last week rain.

## **SMS** advisory

➤ This is the best time for application of fertilizer in a semicircle basin urea @ 650g+1kg super phosphate & KCL along with 50 FYM.

Professor and Head
Principal Nodal Officer (GKMS)
Agro Climate Research Centre