

Agromet Advisory Bulletin for the Perambalur District

(From 01st June 2022 to 05th June 2022)



Date: 31.05.2022

Issued jointly by Tamil Nadu Rice Research Institute - Aduthurai and IMD

Bulletin no: 43/2022

Weather Summary for the period since the issue of last bulletin no. 42 (27.05.2022) for Tamil Nadu:

Maximum temperature of 38 °C to 39 °C and minimum temperature of 25 °C to 28 °C and isolated rainfall were recorded in Perambalur district.

Weather Summary and forecast for Perambalur district

Past week Summary 27.05.2022 to 30.05.2022				Parameters	Ensemble weather forecast valid until 08.30 hrs of 05.06.2022				
Day-1	Day-2	Day-3	Day-4	Date	Day-1	Day-2	Day-3	Day-4	Day-5
27/05	28/05	29/05	30/05		01/06	02/06	03/06	04/06	05/06
0.3	9.3	0	0	Rainfall (mm)	0	0	4	2	0
39	39	38	39	Max. Temp. (°C)	37	38	38	37	39
27	27	28	25	Mini. Temp. (°C)	27	28	27	27	28
5	4	6	5	Cloud cover (Octa)	3	3	5	4	3
66-38	60-38	61-49	62-40	RH I-II (%)	65-45	65-45	65-45	65-45	65-45
19	22	27	12	Wind (kmph)	6	6	6	6	6
West	West	South	South	***	South	West	West	South	South
North	North	East	East	Wind direction	South	South	South	West	West
West	West	Lust	Last		West	West	West	vv est	vvest

Weather forecast for next five days (01.06.2022 to 05.06.2022)

According to district level forecast issued by India Meteorological Department for Perambalur district, the sky will be moderate cloudy. Rainfall is expected on 03.06.2022 and 04.06.2022. Maximum temperature is expected to be around $37\,^{\circ}\text{C}$ to $39\,^{\circ}\text{C}$ and Minimum temperature is expected to be around $27\,^{\circ}\text{C}$ to $28\,^{\circ}\text{C}$. Morning relative humidity is expected to be around $45\,^{\circ}$ percent. Average wind speed is expected to be around $65\,^{\circ}$ km per hour and the wind direction will be south west.

Agro Advisory

Crop	Stage	Advisory	
Rice		For kuruvai nursery, an area of 20 cents can be allotted for planting 1 ha with the application of FYM or compost @ 1 ton. For clay soil, 4 kg gypsum and 1 kg DAP/cent can be applied 10 days	
		prior to pulling out of seedlings to avoid root snapping.	

		The seeds should be treated with carbendazim (2 g/kg of seeds) or Pseudomonas (10 g/kg of seeds) before sowing to prevent seed borne diseases.	
R	ice	Do seed treatment before sowing rice varieties ADT 36, ADT 43, ADT 45 and ADT 47 which are suitable for kuruvai season and take necessary action for preparation of nursery.	
Livestock	Cattle	Highly reflective aluminum roofs may be provided to cattle shed to reduce the heat stress to the cattle underneath.	

SMS advisory

Highly reflective aluminum roofs may be provided to cattle shed to reduce the heat stress to the cattle underneath.

Director TRRI, Aduthurai