



**Gramin Krishi Mausam Sew**  
Experimental Block Level Agromet Advisory Bulletin  
(A Joint Initiative of IMD & ICAR)



## Agromet Advisory Bulletin

Date : 13-05-2022

Weather Forecast of **CHENNAI** Block in **CHENNAI**(Tamil Nadu) Issued On :2022-05-13(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-05-14	2022-05-15	2022-05-16	2022-05-17	2022-05-18
Rainfall	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	36.0	36.0	34.0	33.0	35.0
Tmin(°C)	26.0	25.0	25.0	25.0	25.0
RH-I(%)	73	72	76	76	75
RH-II(%)	52	54	55	55	51
Wind Speed(kmph)	15.0	22.0	20.0	19.0	17.0
Wind Direction(Degree)	216	255	212	218	207
Cloud Cover(Octa)	7	8	8	8	8

### Weather Summary/Alert:

According to district forecast issued by the India Meteorological Department for Chennai district, sky will be mostly clear. Maximum temperature is expected to be around 34.0-35.0°C. Minimum temperature is expected to be around 25.0-27.0°C. Morning relative humidity is expected to be around 80 per cent and evening relative humidity is expected to be around 65 per cent. Wind speed is expected to be around 12 km per hour and the wind direction will be from South South West.

### General Advisory:

From extended range forecast above normal rainfall, maximum temperature and normal minimum temperature expected for the period of 18.05.2022 to 24.05.2022 over Tamil Nadu.

### SMS Advisory:

(14.05.2022-15.05.2022) Heavy rain is likely to occur at Kancheepuram, kallakurichi, Tiruvallur, Villupuram, Tiruvannamalai, Ranipet, Tirupattur, Vellore, Krishnagiri and Cuddalore

### Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Accidental grazing of new shoots of sorghum fodder may lead to HCN poisoning and death, hence farmers are advised to take precautionary measures The drainage system in cattle shed must be open and blockage free with access to easy cleaning to prevent fly breeding
GOAT	2. Small ruminants overfed with excessive grain feed are at the edge of procuring Enterotoxaemia, hence precautionary measures should be taken
SHEEP	3. Ewes which have missed the main breeding season must be flushed by feeding Hedge lucerne or any other leguminous fodder supplemented with vitamin tonics and mineral mixture before mating.
PIG	5. To prevent E.coli infection in nursery piggery shed the floor must be washed with luke warm saline / good irritant free disinfectant atleast weekly once.

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

<b>Poultry(Varieties)</b>	<b>Poultry Specific Advisory</b>
CHICKEN	6. Native poultry reared under integrated method incase infested with lice, must be given medicated bath. A sponge dipped in one ml of butox mixed in one litre of water could be rubbed all over the boy on a bright sunny day. Care must be exercised to avoid eyes ear lobe and mouth.