

Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal - 609 603 Agromet Advisory Bulletin (AAB) for KARAIKAL District Issued jointly by IMD & PAJANCOA & RI Forecast Period:04th to 08th October 2022



Phone No. 04368 – 261182 E-mail: agronaas2012@gmail.com

Fax: 04368-261260

Bulletin No.079 (10)/PJN/GKMS/2022

Date: 03.10.2022

Weather Summary for Karaikal district

Past week Summary (28.09.2022to 02.10.2022)					Weather Parameters	Ensemble weather forecast valid until 08.30 hrs of 08.10.2022				
28/09	29/09	30/09	01/10	02/10		Day-1 04/10	Day-2 05/10	Day-3 06/10	Day-4 07/10	Day-5 08/10
78.50	0	0	0	0	Rainfall (mm)	0	0	4	8	12
34	36	35	32	36	Max. Temp. (° C)	34	34	35	34	33
25	25	26	26	26	Mini. Temp. (°C)	26	27	26	25	24
5	2	4	4	3	Cloud cover (Octa)	6	6	8	8	8
68-97	59-87	70-80	69-80	59-80	RH Even- Morn. (%)	30-70	30-70	40-70	40-80	40-80
4	4	4	4	4	Wind (kmph)	10	8	12	12	8
WSW	WSW	WSW	WSW	WSW	Wind direction	WNW	WNW	WNW	W	NE

Forecast for next five days: 04.10.2022 to 08.10.2022

- Light to moderate rain may occur from 06^{th} to 08^{th} October.
- ▶ Thunderstorm with lightning is likely to occur at isolated places on 04th and 05th October.

During next five days, the sky will be cloudy. The temperature range is expected to be around 24-35°C. Relative humidity range is expected to be around 30-80%. The wind speed is expected to be 08-12 km per hour and the wind will be mostly from west north west direction.

Rainfall (7.39 mm) will be above normal, maximum temperature (31.30°C) and minimum temperature (23.09°C) will be normal from 05th to 11th October.

Agro Advisory

Crop/ Livestock	Stage/ Pariculars	Agro Advisory
Rice		To manage weeds in direct wet seeded rice , apply pyrazosulfuron ethyl @ 200 g/ha at 3-5 days after sowing.
	Weeding	To manage weeds in direct dry seeded rice, apply pendimethalin @ 3.3 l/ha on 3 days after sowing (DAS) followed by bispyripac sodium @ 200 ml/ha as early post emergence (20 DAS) and followed by hand weeding on 40 DAS to manage weeds.
Cattle	Paddy straw enriching	During rainy days, urea enriched paddy straw can be prepared and used for cattle (To enrich 100 kg of paddy straw, dissolve 4 kg urea in 65 litres of water and sprinkle prepared urea solution over the paddy straw and cover it. After 21 days the urea treated paddy straw can be used for feeding).

Vaccination	Before the onset of monsoon, animals may be vaccinated for Foot and Mouth Disease and Hemorrhagic Septicemia.	
Deworming	Grazing animals that are drinking pond water should be dewormed.	

SMS: Light to moderate rain is expected from 06th to 08th October.

DEAN PAJANCOA & RI