



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)

Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of **ANNAVASAL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	2.0	4.0	30.0
Tmax(°C)	38.0	37.0	36.0	36.0	34.0
Tmin(°C)	28.0	29.0	28.0	27.0	25.0
RH-I(%)	63	65	69	75	87
RH-II(%)	34	35	39	43	46
Wind Speed(kmph)	11.0	6.0	7.0	5.0	16.0
Wind Direction(Degree)	268	275	290	284	286
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 06, 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 38°C and the minimum temperature would be in between 25°C and 29°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 5 to 16 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against foot and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of ARANTANGI Block in PUDUKKOTTAI(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	3.0	4.0	28.0
Tmax(°C)	38.0	37.0	36.0	36.0	34.0
Tmin(°C)	28.0	29.0	29.0	28.0	25.0
RH-I(%)	63	65	69	75	87
RH-II(%)	34	35	39	43	47
Wind Speed(kmph)	10.0	6.0	7.0	5.0	17.0
Wind Direction(Degree)	265	270	289	283	280
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 06, 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 38°C and the minimum temperature would be in between 25°C and 29°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 5 to 17 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against foot and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of **ARIMALAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	3.0	4.0	30.0
Tmax(°C)	38.0	37.0	36.0	36.0	34.0
Tmin(°C)	28.0	29.0	29.0	28.0	25.0
RH-I(%)	62	65	69	75	87
RH-II(%)	34	35	39	43	47
Wind Speed(kmph)	10.0	5.0	6.0	5.0	17.0
Wind Direction(Degree)	265	274	292	286	290
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 06, 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 38°C and the minimum temperature would be in between 25°C and 29°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 5 to 17 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against foot and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of AVADAIYARKOVIL Block in PUDUKKOTTAI(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	3.0	5.0	27.0
Tmax(°C)	38.0	37.0	36.0	35.0	34.0
Tmin(°C)	28.0	29.0	29.0	28.0	25.0
RH-I(%)	63	65	69	75	86
RH-II(%)	34	35	40	43	47
Wind Speed(kmph)	10.0	6.0	7.0	5.0	17.0
Wind Direction(Degree)	263	266	289	284	274
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 06, 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 38°C and the minimum temperature would be in between 25°C and 29°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 5 to 17 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against foot and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of **GANDARAVAKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	0.0	3.0	14.0
Tmax(°C)	37.0	37.0	36.0	34.0	33.0
Tmin(°C)	28.0	29.0	28.0	27.0	26.0
RH-I(%)	68	68	71	78	85
RH-II(%)	35	33	40	44	47
Wind Speed(kmph)	13.0	8.0	7.0	4.0	16.0
Wind Direction(Degree)	277	241	261	227	271
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 37°C and the minimum temperature would be in between 26°C and 29°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 4 to 16 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against foot and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of **KARAMBAKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	1.0	4.0	22.0
Tmax(°C)	37.0	37.0	36.0	35.0	34.0
Tmin(°C)	28.0	29.0	28.0	27.0	25.0
RH-I(%)	66	66	70	76	86
RH-II(%)	34	34	39	43	47
Wind Speed(kmph)	12.0	7.0	7.0	5.0	16.0
Wind Direction(Degree)	272	258	276	257	276
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 06, 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 37°C and the minimum temperature would be in between 25°C and 29°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 5 to 17 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against foot and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of **KUNNANDARKOIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	1.0	5.0	30.0
Tmax(°C)	37.0	37.0	35.0	35.0	34.0
Tmin(°C)	26.0	28.0	27.0	26.0	25.0
RH-I(%)	65	66	69	76	88
RH-II(%)	34	34	37	41	45
Wind Speed(kmph)	13.0	8.0	8.0	5.0	15.0
Wind Direction(Degree)	277	273	280	279	271
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 06, 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 37°C and the minimum temperature would be in between 25°C and 28°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 5 to 15 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against foot and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of MANALMELKUDI Block in PUDUKKOTTAI(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	3.0	5.0	25.0
Tmax(°C)	38.0	37.0	36.0	35.0	34.0
Tmin(°C)	28.0	29.0	28.0	27.0	25.0
RH-I(%)	63	65	69	75	86
RH-II(%)	34	35	40	44	48
Wind Speed(kmph)	10.0	6.0	7.0	5.0	17.0
Wind Direction(Degree)	263	261	286	281	268
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 06, 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 38°C and the minimum temperature would be in between 25°C and 29°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 5 to 17 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against foot and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.

Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of **PONNAMARAVATI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	3.0	5.0	30.0
Tmax(°C)	38.0	37.0	36.0	36.0	34.0
Tmin(°C)	28.0	29.0	28.0	27.0	25.0
RH-I(%)	63	65	69	75	87
RH-II(%)	34	34	39	43	47
Wind Speed(kmph)	10.0	6.0	7.0	5.0	16.0
Wind Direction(Degree)	264	273	290	288	277
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 06, 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 38°C and the minimum temperature would be in between 25°C and 29°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 5 to 16 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against foot and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of **PUDUKKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	2.0	4.0	31.0
Tmax(°C)	38.0	37.0	36.0	36.0	34.0
Tmin(°C)	28.0	29.0	29.0	28.0	25.0
RH-I(%)	62	65	69	75	87
RH-II(%)	34	35	39	43	47
Wind Speed(kmph)	10.0	5.0	6.0	5.0	17.0
Wind Direction(Degree)	265	277	295	288	296
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 06, 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 38°C and the minimum temperature would be in between 25°C and 29°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 5 to 17 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.

Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of **THIRUVARANGULAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	2.0	4.0	31.0
Tmax(°C)	38.0	37.0	36.0	36.0	34.0
Tmin(°C)	28.0	29.0	29.0	28.0	25.0
RH-I(%)	62	65	69	75	87
RH-II(%)	34	35	39	43	47
Wind Speed(kmph)	10.0	5.0	6.0	5.0	17.0
Wind Direction(Degree)	265	276	294	287	295
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 06, 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 38°C and the minimum temperature would be in between 25°C and 29°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 5 to 17 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against foot and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of **TIRUMAYAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	3.0	4.0	30.0
Tmax(°C)	38.0	37.0	36.0	36.0	34.0
Tmin(°C)	28.0	29.0	29.0	28.0	25.0
RH-I(%)	62	65	69	75	87
RH-II(%)	34	35	39	43	47
Wind Speed(kmph)	10.0	5.0	6.0	5.0	17.0
Wind Direction(Degree)	265	274	292	287	288
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 06, 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 38°C and the minimum temperature would be in between 25°C and 29°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 5 to 17 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against foot and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 03-10-2022

Weather Forecast of **VIRALIMALAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-10-03(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-04	2022-10-05	2022-10-06	2022-10-07	2022-10-08
Rainfall	0.0	0.0	1.0	5.0	31.0
Tmax(°C)	37.0	37.0	35.0	35.0	34.0
Tmin(°C)	26.0	28.0	27.0	26.0	25.0
RH-I(%)	65	66	69	76	88
RH-II(%)	34	34	37	41	45
Wind Speed(kmph)	13.0	8.0	8.0	5.0	15.0
Wind Direction(Degree)	277	274	280	281	268
Cloud Cover(Octa)	8	8	8	8	8

Weather Summary/Alert:

Chances of light to moderate rain on 06, 07 and 08 th October. Maximum temperature would be in the range 34.0°C and 37°C and the minimum temperature would be in between 25°C and 28°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 5 to 15 kmph.

General Advisory:

Chances of light rainfall for next five days. Proper drainage facilities to be ensured in the crops fields. Due to the rains for past few days there might be deficiency symptoms on leaves. Soil application of Urea 10 kg/ha, Gypsum 8 kg/ha and Neemcake 2 kg/ha can be done.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Foliar spray of the micronutrient is important for the Blackgram crop. Foliar spray of NAA 40mg/litre and Salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later. For rice fallow crops, foliar spray of DAP 20gm/litre once at flowering and another at 15 days later. For irrigated and rain-fed crops, foliar spray of DAP 20gm/litre or Urea 20 gm/litre once at flowering and another at 15 days later. Foliar spray of salicylic acid 100 mg/litre once at pre-flowering and another at 15 days later.
RICE	As the sowing of Samba season is nearing end, direct sowing with pre germinated short duration paddy varieties seeds in puddled soil is recommended. Drainage may be provided in recently transplanted samba rice.

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	There are chances of diarrhoea in the poultry. To protect the other birds from Diarrhoea, keep the diseased birds separately. Medicate the drinking water with 2gm Microsol or Bactrisol + 5ml Vimrol Syrup per litre of water. Medicated water should be given to all the birds for 5 days in the morning and evening time.