



**Gramin Krishi Mausam Sewa**  
**District Level Agromet Advisory Bulletin**  
 Tamil Nadu Agricultural University,  
 Coimbatore, Tamil Nadu



## Agromet Advisory Bulletin

Date : 25-10-2022

Weather Forecast of District PUDUKKOTTAI(Tamil Nadu) Issued On : 2022-10-25(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-10-26	2022-10-27	2022-10-28	2022-10-29	2022-10-30
Rainfall(mm)	2.0	0.0	0.0	0.0	4.0
Tmax(°C)	33.0	34.0	34.0	33.0	32.0
Tmin(°C)	25.0	26.0	26.0	26.0	25.0
RH-I(%)	80	80	80	80	80
RH-II(%)	70	70	70	70	70
Wind Speed(kmph)	10.0	10.0	10.0	10.0	12.0
Wind Direction(Degree)	110	20	340	340	340
Cloud Cover(Octa)	5	3	3	4	6

### Weather Summary/Alert:

Chances of light rain from 26 and 30 th october. Maximum temperature would be in the range 32.0°C and 34.0°C and the minimum temperature would be in between 25.0°C - 26.0°C. Generally cloudy sky is expected for next 5 days. Wind speed is expected to be 10 to 12 kmph.

### General Advisory:

By using this weather condition, root feeding of TNAU coconut tonic @200ml/palm is advised in nut bearing coconut trees.

### SMS Advisory:

As the rains are expected, the farmers have to clean the canals and water bodies and carry out rain water harvesting activities.

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Using the prevailing weather conditions, paddy farmers can raise Azolla as a dual crops by in corporate during wedding for the wet season crop and broadcast 10 kg of soil based powdered BGA flakes at 10 DAT and maintain a thin film of water for multiplication.
RICE	With the prevailing weather condition, in dry seeded Rain fed Paddy, it is time for application of Manures and Fertilizer. Farmers are advised to apply 50kg N, 28kg P, 25kg K per hectare as blanket recommendation and apply N and K in two equal splits as 12.5kg at 20 - 25 and 40 - 45 days after germination.

### Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
-------------------------	--------------------------------

<b>Horticulture(Varieties)</b>	<b>Horticulture Specific Advisory</b>
TOMATO	Tomato- Early blight incidence Spray of hexaconazole 5% SC @ 1ml/l or propiconazole 25% EC @ 1 ml/l at 30 and 50 days after planting.