



TNAU & IMD

Weather Summary for the period since the issue of last bulletin 77 (27.09.2022) for Tamil Nadu. Maximum temperature ranged on 30-37°C and minimum temperature ranged on 22-27°C, over the State. Kodaikanal and Uthagamandalam recorded maximum temperature ranged on 18-21°C and minimum temperature ranged on 10-11°C. During this period isolated distribution with very light to moderate intensity of rainfall was recorded in Tamil Nadu.

Past week weather Summary (25.09.2022 to 29.09.2022)				Weather	Ensemble weather forecast valid until 08:30 hrs of 05.10.2022					
Day 1 25/09	Day 2 26/09	Day 3 27/09	Day 4 28/09	Day 5 29/09	Parameters	Day 1 01/10	Day 2 02/10	Day 3 03/10	Day 4 04/10	Day 5 05/10
0.0	8.2	0.0	0.1	0.9	Rainfall (mm)	0	0	6	6	8
37	38	35	37	37	Max. Temp. (°C)	37	36	36	36	36
24	25	25	25	26	Min. Temp. (°C)	26	26	26	26	26
3	8	3	5	5	Cloud cover (Octa)	2	4	6	8	8
					Relative humidity -					
54	74	57	59	74	Mor. (%)	60	60	60	60	70
					Relative humidity -					
39	56	47	45	46	Eve. (%)	30	40	40	30	40
					Wind speed					
4	4	6	8	6	(kmph)	6	6	6	6	6
90/E	320/NW	90/E	290/ WNW	90/E	Wind direction (dd)	260/W	260/W	270/W	280/W	280/W

Weather Summary for Virudhunagar district

Weather forecast for next five days: 01.10.2022 to 05.10.2022

According to district forecast issued by the India Meteorological Department for Virudhunagar district, sky condition will be partly to generally cloudy sky. Light rainfall is predicted for day 3 (03.09.2022) to day 5 (05.10.2022). The Maximum temperature is expected to be around 36-37°C and Minimum temperature is expected to be around 26°C. Morning relative humidity is expected to be around 60-70 per cent and evening relative humidity is expected to be around 30-40 per cent. 24-hour wind speed is expected to be around 8-10 km per hour and the wind direction will be from West direction.

Agromet Advisory

Crop/ Ecosystem	Stage / Activities	Advisories
Irrigated crops	Water management	Very light to light rainfall received during past week may not be sufficient for crop. Light rainfall is predicted for day 3 (03.10.2022) to day 5 (05.10.2022). So, the farmers are advise to plan the irrigation, only if the crop was under stress.
Irrigated crops	Spraying	As rainfall is predicted, spraying of chemical for sucking pest may be avoided if the pest is below ETL. However, if the pest is found above the ETL spraying may be done.
Cotton	vegetative stage	In order to utilize the receipt of rainfall for the crop growth, farmers are requested to go for weeding.
Blackgram	vegetative stage	 Prevailed dry condition is favorable for Yellow Mosaic Virus. Symptoms: The leaves show irregular yellow and green patches alternating with each other. Then leaves turn completely yellow and became papery. Infected leaves show necrotic symptoms. Diseased plants stunted, mature late produce very few flowers and pods. Pods are reduced in size and shriveled. No. of seeds are also reduced. Management- Place yellow sticky traps 12 nos./ha. Spray methyl demeton 500ml/ha and repeat 15 days interval if it is needed. Tobacco cutworm: Symptoms: 1. Early instar larvae feed gregariously by scraping on the leaf.
		2. Later instar larvae completely deover the leaves causing defoliation. Management : 1. Setup light trap to monitor and kill the attracted the adult moth. 2. Install pheromone trap @ 5 Nos./acre. 3. Removal and destruction of egg masses. 3. Spray Chlorantraniliprole 18.5% SC @ 60ml or Flubendiamide 39.35% SC @40ml in 200litres of water per acre.
Groundnut	Flowering stage	 Red Hairy Caterpillar: Symptoms: 1. Early instar larvae feed voraciously by scraping the under surface of the tender leaves. 2. Grown up larvae feed voraciously on entire foliage, flowers and growing points. They often migrate from field to field in search of food after devastating the foliage in the field where they have hatched. 3. Severely damaged crop gives the appearance of having been grazed by cattle. Management: 1. Collect and destroy the egg masses and voracious early instar larvae. 2. Set up bone fires or light trap up to 11.00 P.M. after receipt of rain to attract and kill the moths. 3. Spray Quinalphos 25% Ec @ 500ml in 300litres of water per acre. 4. Grow cowpea or red gram as an intercrop to attract adult moths to lay more eggs.
Maize	Vegetative satge	The prevailing weather condition is conducive for fall armyworm incidence. To manage this pest, spray chlorantraniliprole 18.5% SC @ 0.4ml/ litre of water or Flubendiamide 480 SC @ 0.4 ml/ litre of water. Use only hand sprayer and direct the spray towards whorl region.
Guava	Fruiting stage	Prevailing weather is conducive for Tea mosquito bug. Spray malathion 2ml/lit water. (or) Thiamethoxam 25WG 40g/acre. Spray 3% neem oil at 3 weeks interval for 4 times. (or) Collect and destroy the damaged plant parts.
Coconut		Due to lack of nutrient, Button shedding is noticed. Root feeding of the TNAU coconut tonic @ 200 ml / palm twice in a year at six months interval, increases number and size of nuts.

Chilli	Nursery preparation	As the rainfall is predicted farmers are advised to utilize the rainfall for nursery sowing after incorporation of neem cake in the soil to protect the seedlings from pest and diseases.
Livestock	Sheep	Haemorrhagic Septicaemia: It is a serious acute disease of cattle and Buffaloes caused by a bacteria. Occurrence in more common in rainy season. Affected animals will have high fever, salivation, respiratory distress, and swelling in the throat and neck region. In most of the cases, affected animals will die within 12 to 24 hrs. Disease transmission occurs through direct contact with the affected animals and ingestion of contaminated feed and water. Treatment: Treating the animals with suitable antibiotics at the early stage can prevent the death of the animals. Control: Annual vaccination should be followed before monsoon. Reducing stress to the animals may prevent the occurrence.

SMS: Rain:03-05.10-6,6&8 mm. Maize - fall armyworm- spray chlorantraniliprole 18.5% SC @ 0.4ml/ litre of water. Use only hand sprayer and direct the spray towards whorl region.

		Tamil Nadu
		n over Westcentral Bay of Bengal off Andhra Pradesh coast persists bove mean sea level.
Next five days	Weather warning	Light to moderate rain at a few places with isolated thunderstorm & lightning is likely to occur over Tamilnadu, Puducherry & Karaikal area on Day 1 (01.10.2022). Heavy rain is likely to occur at one or two places over Salem, Dharmapuri, Krishnagiri, Tirupattur, Vellore, Ranipet, Tiruvallur, Kancheepuram and Tiruvannamalai districts of Tamilnadu on Day 1 (01.10.2022). Thunderstorm & lightning with light to moderate rain is likely to occur at one or two places over Tamilnadu, Puducherry & Karaikal area on Day 2 (02.10.2022). Heavy rain is likely to occur at one or two places over Nilgiris, Coimbatore, Tiruppur, Dindigul and Theni of Tamilnadu on Day 2 (02.10.2022).
		Thunderstorm & lightning with light to moderate rain is likely to occur at one or two places over Tamilnadu, Puducherry & Karaikal area on Day 3 (03.10.2022). Light to moderate rain is likely to occur at one or two places over Tamilnadu, Puducherry & Karaikal area on Day 4 (04.10.2022) & Day 5 (05.10.2022).
Extended range of weather forecast		Week (07 th - 13 th Oct 2022): There is possibility of normal to above normal rainfall over Tamilnadu, Puducherry and Karaikal during the week.

Professor and Head Agricultural Research Station Kovilpatti