



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **ANNAVASAL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	13.0	13.0	8.0	22.0	21.0
Tmax(°C)	33.0	34.0	36.0	36.0	35.0
Tmin(°C)	27.0	27.0	28.0	28.0	28.0
RH-I(%)	79	77	76	78	78
RH-II(%)	53	49	44	47	50
Wind Speed(kmph)	11.0	7.0	7.0	7.0	7.0
Wind Direction(Degree)	216	234	199	203	234
Cloud Cover(Octa)	3	4	3	4	6

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 36°C and the minimum temperature would be in between 27°C - 28.0°C. scattered sky is expected for next 5 days. Wind speed is expected to be 7 to 11 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with *Bacillus subtilis* @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COWPEA/ LOBIA	As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **ARANTANGI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	12.0	13.0	7.0	20.0	21.0
Tmax(°C)	33.0	34.0	36.0	36.0	35.0
Tmin(°C)	27.0	27.0	28.0	28.0	28.0
RH-I(%)	79	77	76	77	77
RH-II(%)	54	49	44	48	50
Wind Speed(kmph)	12.0	8.0	7.0	7.0	7.0
Wind Direction(Degree)	213	229	192	198	229
Cloud Cover(Octa)	3	4	3	4	6

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 36°C and the minimum temperature would be in between 27°C - 28.0°C. scattered sky is expected for next 5 days. Wind speed is expected to be 7 to 12 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with *Bacillus subtilis* @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COWPEA/ LOBIA	As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **ARIMALAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	12.0	14.0	7.0	20.0	21.0
Tmax(°C)	33.0	34.0	36.0	36.0	35.0
Tmin(°C)	27.0	27.0	28.0	28.0	28.0
RH-I(%)	79	78	76	78	78
RH-II(%)	54	50	45	48	51
Wind Speed(kmph)	12.0	7.0	6.0	7.0	6.0
Wind Direction(Degree)	213	228	188	194	226
Cloud Cover(Octa)	3	4	3	4	6

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 36°C and the minimum temperature would be in between 27°C - 28.0°C. scattered sky is expected for next 5 days. Wind speed is expected to be 6 to 12 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Bacillus subtilis @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COWPEA/ LOBIA	As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **AVADAIYARKOVIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	12.0	12.0	6.0	20.0	21.0
Tmax(°C)	33.0	34.0	36.0	36.0	35.0
Tmin(°C)	27.0	27.0	28.0	28.0	28.0
RH-I(%)	79	77	75	77	77
RH-II(%)	54	49	44	47	50
Wind Speed(kmph)	12.0	9.0	7.0	7.0	7.0
Wind Direction(Degree)	213	228	192	199	230
Cloud Cover(Octa)	3	4	3	4	6

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 36°C and the minimum temperature would be in between 27°C - 28.0°C. scattered sky is expected for next 5 days. Wind speed is expected to be 7to 12 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Bacillus subtilis @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COWPEA/ LOBIA	As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **GANDARAVAKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	1.0	10.0	7.0	20.0	21.0
Tmax(°C)	33.0	34.0	36.0	36.0	34.0
Tmin(°C)	27.0	28.0	28.0	28.0	28.0
RH-I(%)	81	78	78	81	80
RH-II(%)	53	49	44	48	51
Wind Speed(kmph)	11.0	10.0	9.0	9.0	7.0
Wind Direction(Degree)	200	228	226	216	236
Cloud Cover(Octa)	4	4	3	4	5

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 36°C and the minimum temperature would be in between 27°C - 28.0°C. scattered sky is expected for next 5 days. Wind speed is expected to be 6 to 12 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Bacillus subtilis @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COWPEA/ LOBIA	As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **KARAMBAKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	8.0	11.0	8.0	22.0	21.0
Tmax(°C)	33.0	34.0	36.0	36.0	35.0
Tmin(°C)	27.0	27.0	28.0	28.0	28.0
RH-I(%)	80	77	77	79	79
RH-II(%)	53	49	44	48	50
Wind Speed(kmph)	11.0	9.0	8.0	8.0	7.0
Wind Direction(Degree)	209	232	213	211	236
Cloud Cover(Octa)	4	4	3	4	6

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 36°C and the minimum temperature would be in between 27°C - 28.0°C. scattered sky is expected for next 5 days. Wind speed is expected to be 7 to 11 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Bacillus subtilis @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COWPEA/ LOBIA	As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **KUNNANDARKOIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	19.0	10.0	13.0	31.0	21.0
Tmax(°C)	33.0	35.0	37.0	36.0	35.0
Tmin(°C)	25.0	27.0	27.0	26.0	27.0
RH-I(%)	80	76	76	78	77
RH-II(%)	50	46	41	45	47
Wind Speed(kmph)	8.0	8.0	8.0	8.0	8.0
Wind Direction(Degree)	226	252	237	234	260
Cloud Cover(Octa)	4	5	4	6	6

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 37°C and the minimum temperature would be in between 25°C - 27°C. scattered sky is expected for next 5 days. Wind speed is expected to be 8 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Bacillus subtilis @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
SORGHUM (JOWAR/GREAT MILLET)	By utilizing rainfall, fodder sorghum or cowpea may be sown under rainfed lands.
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **MANALMELKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	12.0	11.0	6.0	20.0	20.0
Tmax(°C)	33.0	34.0	36.0	36.0	35.0
Tmin(°C)	27.0	27.0	28.0	28.0	28.0
RH-I(%)	79	76	75	77	76
RH-II(%)	54	49	44	48	50
Wind Speed(kmph)	12.0	9.0	7.0	8.0	8.0
Wind Direction(Degree)	212	228	197	203	231
Cloud Cover(Octa)	3	4	3	4	6

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 36°C and the minimum temperature would be in between 27°C - 28°C. scattered sky is expected for next 5 days. Wind speed is expected to be 7 to 12 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Bacillus subtilis @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
SORGHUM (JOWAR/GREAT MILLET)	By utilizing rainfall, fodder sorghum or cowpea may be sown under rainfed lands.
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **PONNAMARAVATI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	14.0	12.0	7.0	22.0	22.0
Tmax(°C)	33.0	34.0	36.0	36.0	35.0
Tmin(°C)	27.0	27.0	28.0	28.0	28.0
RH-I(%)	79	77	75	77	77
RH-II(%)	53	49	44	47	49
Wind Speed(kmph)	11.0	8.0	6.0	7.0	7.0
Wind Direction(Degree)	215	230	190	199	232
Cloud Cover(Octa)	3	4	3	4	6

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 36°C and the minimum temperature would be in between 27°C - 28°C. scattered sky is expected for next 5 days. Wind speed is expected to be 6 to 11 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Bacillus subtilis @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
SORGHUM (JOWAR/GREAT MILLET)	By utilizing rainfall, fodder sorghum or cowpea may be sown under rainfed lands.
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **PUDUKKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	12.0	14.0	7.0	19.0	21.0
Tmax(°C)	33.0	34.0	36.0	36.0	35.0
Tmin(°C)	27.0	27.0	28.0	28.0	28.0
RH-I(%)	79	78	76	78	78
RH-II(%)	54	50	45	48	51
Wind Speed(kmph)	12.0	7.0	6.0	7.0	6.0
Wind Direction(Degree)	212	227	183	189	223
Cloud Cover(Octa)	3	4	3	4	6

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 36°C and the minimum temperature would be in between 27°C - 28.0°C. scattered sky is expected for next 5 days. Wind speed is expected to be 6 to 12 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Bacillus subtilis @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
SORGHUM (JOWAR/GREAT MILLET)	By utilizing rainfall, fodder sorghum or cowpea may be sown under rainfed lands.
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.



Gramin Krishi Mausam Sew
Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **THIRUVARANGULAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	12.0	14.0	7.0	19.0	21.0
Tmax(°C)	33.0	34.0	36.0	36.0	35.0
Tmin(°C)	27.0	27.0	28.0	28.0	28.0
RH-I(%)	79	78	76	78	78
RH-II(%)	54	50	45	48	51
Wind Speed(kmph)	12.0	7.0	6.0	7.0	6.0
Wind Direction(Degree)	212	227	184	190	224
Cloud Cover(Octa)	3	4	3	4	6

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 36°C and the minimum temperature would be in between 27°C - 28°C. scattered sky is expected for next 5 days. Wind speed is expected to be 6 to 12 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Bacillus subtilis @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
SORGHUM (JOWAR/GREAT MILLET)	By utilizing rainfall, fodder sorghum or cowpea may be sown under rainfed lands.
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **TIRUMAYAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	12.0	14.0	7.0	20.0	21.0
Tmax(°C)	33.0	34.0	36.0	36.0	35.0
Tmin(°C)	27.0	27.0	28.0	28.0	28.0
RH-I(%)	79	77	76	78	77
RH-II(%)	54	50	44	48	50
Wind Speed(kmph)	12.0	8.0	6.0	7.0	6.0
Wind Direction(Degree)	213	228	187	194	227
Cloud Cover(Octa)	3	4	3	4	6

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 36°C and the minimum temperature would be in between 27°C - 28°C. scattered sky is expected for next 5 days. Wind speed is expected to be 6 to 12 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Bacillus subtilis @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
SORGHUM (JOWAR/GREAT MILLET)	By utilizing rainfall, fodder sorghum or cowpea may be sown under rainfed lands.
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 02-09-2022

Weather Forecast of **VIRALIMALAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2022-09-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-03	2022-09-04	2022-09-05	2022-09-06	2022-09-07
Rainfall	20.0	9.0	13.0	32.0	21.0
Tmax(°C)	33.0	35.0	37.0	36.0	35.0
Tmin(°C)	25.0	27.0	27.0	26.0	27.0
RH-I(%)	80	76	76	77	77
RH-II(%)	50	46	41	45	47
Wind Speed(kmph)	8.0	8.0	8.0	8.0	9.0
Wind Direction(Degree)	227	253	239	236	262
Cloud Cover(Octa)	4	5	4	6	6

Weather Summary/Alert:

Chances of light to moderate rain from 03 to 07 th September. Maximum temperature would be in the range 33°C and 37°C and the minimum temperature would be in between 26°C - 27°C. scattered sky is expected for next 5 days. Wind speed is expected to be 8 to 9 kmph.

General Advisory:

This is a season for vegetable sowing, hence the farmers are advised to treat the vegetable seeds with Bacillus subtilis @ 4 g / kg of seed before sowing to avoid rot in seedlings.

SMS Advisory:

As regular monsoon rains are expected, small pits can be dug at regular intervals in rainfed areas to store rainwater, thus increasing soil moisture by 10-20 per cent.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
SORGHUM (JOWAR/GREAT MILLET)	By utilizing rainfall, fodder sorghum or cowpea may be sown under rainfed lands.
JASMINE	If jasmine plants are deficient in nutrients due to heavy rains so far, spray 5 g of ferrous sulfate solution per liter once in 3 months to prevent yellowing of leaves, iron deficiency, root rot and root rot and iron deficiency. For root rot, a 0.25% solution of copper oxychloride should be applied to the soil around the plant. The disease can be controlled by cultivating coriander in well drained lands. For nematode infestation, apply 5 g of furadon around the plant and mix with soil and water.
RICE	To control leaf blight and sucking insects, mix 2 ml / l of Azadirachtin in running water and spray through the leaf.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	The incense of Bacterial leaf spot disease in Tomato growing areas. To control this spraying with a combination of copper and organic fungicides in a regular preventative spray program at 5 to 10 day intervals or spraying with Agrimycin-100 (100 ppm) or Copper hydroxide at fortnight intervals is recommended for effective disease control.

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Fodder is prone to mildew due to rains so the fodder should be dried in the sun during non-rainy seasons and then fed to the cattle. Protect folder from rain soaking.