

Agromet Advisory Bulletin for the Thanjavur District (From 17th September 2022 to 21st September 2022) Issued jointly by Tamil Nadu Rice Research Institute Aduthurai and IMD



Date: 16.09.2022

Bulletin no: 74/2022

Weather Summary for the period since the issue of last bulletin no. 73 (13.09.2022) for Tamil Nadu: Thanjavur district

Maximum temperature of 32 °C to 35 °C and minimum temperature of 26 °C to 27 °C were recorded in Thanjavur district.

Weather Summary and forecast for Thanjavur district

Past week Summary 13.09.2022 to 15.09.2022			Parameters	Ensemble weather forecast valid until 08.30 hrs of 21.09.2022				
Day-1	Day-2	Day-3	Date	Day-1	Day-2	Day-3	Day-4	Day-5
13/09	14/09	15/09		17/09	18/09	19/09	20/09	21/09
0	0	0	Rainfall (mm)	0	0	2	2	0
35	32	34	Max. Temp. (°C)	35	35	35	35	35
27	27	26	Min. Temp. (°C)	27	27	27	27	27
4	6	5	Cloud cover	4	4	4	5	4
73-68	76-66	73-63	RH I-II (%)	70-60	70-60	70-60	70-60	70-60
5	4	4	Wind (kmph)	8	8	8	12	12
West North West	West North West	West North West	Wind direction	South	South West	West South West	West	West

Weather forecast for next five days (17.09.2022 to 21.09.2022)

According to district level forecast issued by India Meteorological Department for Thanjavur district, the sky will be moderate cloudy. Minimal rainfall is expected on 19.09.2022 and 20.09.2022. Maximum temperature is expected to be around 35 °C and Minimum temperature is expected to be around 27 °C. Morning relative humidity is expected to be around 70 percent and evening relative humidity is expected to be around 8-12 km per hour and the wind direction will be west.

Agro Advisory

Crop	Stage	Advisory			
Gei	neral	Apply 5 tons of FYM per acre during land preparation for rice planting.			
General		Farmers can utilize the mobile apps viz., (TNAU AAS, Meghdoot and DAMINI), for more updated weather information			
Rice	Nursery	For rice nursery, applications of 2 kg DAP/cent is recommended. In rice nursery, spray Azadiraction 0.03% (3 ml/litre of water) with sticking agent Khadi Gopura Bar Soap @ 3g/litre to prevent thrips infection on 10 days after sowing.			
Cattle	Cow	This situation is favorable for the development of Aflatoxin infection in the Corn kernels and Paddy Straw, which may affect the health of the cattle. To save them from infection, dry the Corn kernels and Paddy Straw before feeding.			

SMS advisory

Apply 5 tons of FYM per acre during land preparation for rice planting.

Director TRRI, Aduthurai