9/6/22, 4:59 PM District Advisory



Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin

Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu



Agromet Advisory Bulletin

Date: 06-09-2022

Weather Forecast of District THIRUVALLUR(Tamil Nadu) Issued On: 2022-09-06(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-07	2022-09-08	2022-09-09	2022-09-10	2022-09-11
Rainfall(mm)	8.0	6.0	12.0	6.0	4.0
Tmax(°C)	35.0	35.0	34.0	34.0	35.0
Tmin(°C)	26.0	26.0	25.0	25.0	25.0
RH-I(%)	75	75	75	75	75
RH-II(%)	60	60	60	60	60
Wind Speed(kmph)	10.0	10.0	10.0	12.0	14.0
Wind Direction(Degree)	200	200	230	230	230
Cloud Cover(Octa)	5	6	7	5	5

Weather Summary/Alert:

Light to moderate rainfall may be expected for forthcoming days, Wind speeds may be range from 10.0-14.0 km/ hour; Wind direction may be in the South South West to South West direction; The maximum temperature may be range from 34.0°C-35.0°C and minimum temperatures may be range from 25.0°C-26.0°C; Morning Relative humidity may be range 75% and Evening Relative humidity may be range 60%.

General Advisory:

From extended range forecast Below normal rainfall, normal maximum temperature and minimum temperature expected for the period of 11.09.2022 to 17.09.2022 over Tamil Nadu.

SMS Advisory:

Light to moderate rainfall may be expected for forthcoming days, So farmers are advised to provide proper drainage facilities to crop.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory			
RICE	Rice- Transplanting and Tillering stage-1. Stem bore- Spray Azadirachtin 0.03% 400 ml (Or) Flubendiamide 20% WG 50g. (Or); 2.Whorl maggots- Spray Cartap hydrochloride 4% G 7.5-10 kg (or) Spray Fipronil 5% SC 400-600 g per acre in 200 litres of water; 3. Alage growth of paddy field- apply 2 kg of copper sulfate mix with sand.			
SUGARCANE	Mealy bug- Application of imidachloprid 17.8 SL 3 ml in 10 litres of water.			
	Groundnut-leaf miner-Quinalphos 25 EC 400 ml per acre in 200 litres of water and Set up light trap between 8 to 11 pm at ground level.			
$H \Delta \times M H \times H$	Blossom midge- Spray Chlorantraniliprole 18.5 SC @ 0.5 ml/lit (or) Spray Novaluron 10 EC 3ml/lit.			

Horticulture Specific Advisory:

9/6/22, 4:59 PM District Advisory

Horticulture(Varieties)	Horticulture Specific Advisory			
BRINJAL	Fruit borer- Spray Neem Seed Kernel Extract 5 % (or) Spray Dimethoate 30 % EC 7.0 ml/10 l. and Encourage the activity of larval parasitoids: Pristomerus testaceus, Cremastus flavoorbitalis.			