

Provide drainage facilities in the in Paddy and other fields to avoid water stagnation.

Weather Based Agro-Advisories: (Banana, Paddy & Animal Husbandry).

Banana:

The incidence of yellow Sigatoka disease is seen in Banana fields. The symptoms are slight discolouration between the leaf's secondary veins.

Over a while these points develop into pale yellow streaks, brown streaks and elliptic necrotic spots arranged parallel to the secondary veins. This disease spread rapidly.

To control this, add 10gms of Pseudomonas Fluorescence + 10gms of Bacillus Subtilis + 10 ml of Panchakavya + 5gms of Jaggery in one litre of water and spray it during the morning or evening hours.

Crop protection activities should be done only on non-rainy days.

Paddy:

The incense of Brown Spot disease is expected in Paddy. The disease appears first as minute brown dots later becoming cylindrical or oval to circular on leaves. Spots coalesce to form large patches leading to the leaf drying up.

Infection also occurs on the panicle and neck with a brown colour appearance.

To control spray Metominostrobin @ 500ml /acre.

Crop protection activities should be done only on non-rainy days.

Animal Husbandry:

Due to the prevailing cold weather conditions, cattle rearers are advised to allow the cattle to graze only after 2 hours of sunrise.

Cover the cow shed with gunny bags, and keep the floor in dry condition.

Dry fodder and greens should be fed without moisture to avoid fungal infection.