2/17/23, 4:45 PM District Advisory



Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin M.S.Swaminathan Research Foundation (MSSRF)



Taramani, Chennai

Agromet Advisory Bulletin

Date: 17-02-2023

Weather Forecast of District THENI(Tamil Nadu) Issued On: 2023-02-17(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-02-18	2023-02-19	2023-02-20	2023-02-21	2023-02-22
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	34.0	34.0	35.0	35.0	35.0
Tmin(°C)	19.0	20.0	20.0	21.0	21.0
RH-I(%)	80	90	90	80	70
RH-II(%)	30	20	30	20	20
Wind Speed(kmph)	4.0	6.0	4.0	8.0	8.0
Wind Direction(Degree)	20	20	50	20	360
Cloud Cover(Octa)	0	0	0	0	0

Weather Summary/Alert:

For the next five days, Maximum temperature would be in the range of 34°C to 35°C and the minimum temperature would be between 19°C - 21°C. Clear sky will appear for the next five days. Wind expected with the speed of 4 - 8 Kmph from Northeast & North directions.

General Advisory:

Extended range Forecast for next week (17th – 23rd Feb 2023): Rainfall: As per extended range predictions, there is possibility of below normal to near normal rainfall likely over the district during 17th to 23rd Feb 2023. Maximum Temperature: Maximum temperatures is likely to be generally near normal and are likely to be in the range 30°C to 32°C over the plains of the district during next week. 17th to 23rd Feb 2023. Minimum Temperature: Minimum temperatures are likely to be generally near normal and are likely to be in the range 16°C to 20°C over the district during next week 17th to 23rd Feb 2023.

SMS Advisory:

For the next five days, Max temp- 34°C-35°C & Min temp -19°C- 21°C. Clear sky for next 5 days. Wind speed 4-8 kmph from NE & N directions.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory			
	The incidence of Thrips is expected in Grapes. The symptoms of damage are presence of scorch patches on the leaves and fruits, the premature fruit get dried. To control, spray Thiomethoxam @ 50gms per acre.			
	There is incense of Jassids attack in Bhendi. The symptoms are tender leaves become yellow, margin of the leaves start curling downwards and reddening, severe infestation leaves get a bronze which is typical "hopper burn" symptom, leaf get broken, crumble and growth of the crop is retarded. To control spray any one of the following insecticides Imidacloprid 200 SL at 100 ml/ha with spray fluid 500 litres /ha.			

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory	
	Due to the prevailing weather condition there is chance of infection Anthrax in cattle. The symptoms are fever, severe diarrhea and dysentery, respiration becomes rapid and shortly	

2/17/23, 4:45 PM District Advisory

Live Stock(Varieties)	Live Stock Specific Advisory
	before death the mucous membranes appear cyanotic. The nasal discharge appears with blood stain or purulent. As a preventive method immediately contact the local veterinarian
	and vaccinate your animals once in a year.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	In order to overcome the fodder crisis during summer months, farmers are requested to allot 10 per cent of the irrigated land to fodder crops viz., Cumbu Napier Hybrid or any other fodder crops. Since mainly dry weather is expected for the next five days. Maize & Paddy cultivators are advised to plan for harvest.