



Agromet Advisory Bulletin for the **Tiruvarur** District

(From 04th October 2023 to 08th October 2023)

Issued jointly by Tamil Nadu Rice Research Institute -
Aduthurai and IMD



Bulletin no: 79/2023

Date: 03.10.2023

Weather Summary for the period since the issue of last bulletin no. 78 (29.09.2023) for Tamil Nadu: Tiruvarur district

Maximum temperature of 33 - 35 °C, minimum temperature of 25 - 26 °C, isolated rainfall was recorded in Tiruvarur district.

Weather Summary and forecast for Tiruvarur district

Past week Summary 29.09.2023 to 02.10.2023				Parameters	Ensemble weather forecast valid until 08.30 hr of 08.10.2023				
Day-1 29/09	Day-2 30/09	Day-3 01/10	Day-4 02/10		Date	Day-1 04/10	Day-2 05/10	Day-3 06/10	Day-4 07/10
4.0	0.0	6.0	0.0	Rainfall (mm)	0	0	0	2	0
35	33	34	35	Max. Temp. (°C)	35	36	36	35	35
25	25	26	26	Min. Temp. (°C)	26	25	25	25	25
6	6	6	5	Cloud cover (Octa)	3	1	4	6	4
81-63	74-71	78-62	74-65	RH I-II (%)	80-40	80-40	80-40	90-40	80-50
4	4	4	4	Wind (kmph)	18	16	16	16	16
West North West	West North West	West North West	West North West	Wind direction	South South West	South South West	South South West	South South West	South

Weather forecast for next five days (04.10.2023 to 08.10.2023)

According to district level forecast issued by India Meteorological Department for Tiruvarur district, the sky will be clear. Light rainfall is expected on 07.10.2023. Maximum temperature is expected to be around 35 – 36 °C and minimum temperature is expected to be around 25 - 26 °C. Morning relative humidity is expected to be around 80 - 90 percent and evening relative humidity is expected to be around 40 - 50 percent. Average wind speed is expected to be around 16 – 18 km per hour. The wind direction will be south south west.

Agro Advisory

Crop	Stage	Advisory
General		Farmers are requested to download and utilize the agricultural based mobile apps suggested by TNAU (TNAU AAS, Meghdoot and DAMINI).
Rice	Main field	By utilizing dry weather, fertilizers and pesticides can be applied. Apply 32.6 kg Urea and 8.4 kg of MOP should be applied after 35 days after transplanting for 1 acre rice field.
		The prevailing weather condition favors Thrips infestation. To control the Thrips damage in transplanted rice, apply Thiamethxam 25WG @ 40g/ acre.
cotton		Due to prevailing weather condition there are chances of cotton green leaf hopper infestation. To control spray thiamethxam 25 WG 0.6 gram/litre or imidacloprid 17.8 SL 0.6 ml/litre
Cattle	Cow	Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

SMS advisory

Farmers are requested to download and utilize the agricultural based mobile apps suggested by TNAU (TNAU AAS, Meghdoot and DAMINI).

**Director
TRRI, Aduthurai**