



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **ANNAVASAL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	0.0	1.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	24.0	25.0	25.0	25.0	24.0
RH-I(%)	70	70	70	70	70
RH-II(%)	32	41	41	41	41
Wind Speed(kmph)	17	16	14	16	12
Wind Direction(Degree)	209	212	208	211	211
Cloud Cover(Octa)	7	7	6	8	8

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 24.0°C and 25.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 12 to 17 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.
SUGARCANE	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of nitrogen and 112 ½kg of K2O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm).
Live Stock(Varieties)	Live Stock Specific Advisory
Horticulture Specific Advisory: Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the
Live Stock Specific Advisory:	

<p>Live Stock (Varieties)</p>	<p>60th day with another 12 packets (2400gm). Live Stock Specific Advisory</p>
<p>Horticulture Specific Advisory: Horticulture (Varieties) BANANA</p>	<p>Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Maneozeb and spray.</p>
<p>Live Stock Specific Advisory: COW</p>	<p>Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.</p>
<p>GOAT</p>	<p>Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.</p>

Others (Soil / Land Preparation) Specific Advisory:

<p>Others (Soil / Land Preparation) (Varieties)</p>	<p>Others (Soil / Land Preparation) Specific Advisory</p>
<p>GENERAL ADVICE</p>	<p>Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.</p>



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **ARANTANGI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	0.0	1.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	25.0	25.0	25.0	25.0	24.0
RH-I(%)	70	70	70	71	70
RH-II(%)	35	43	43	43	42
Wind Speed(kmph)	18	17	15	16	13
Wind Direction(Degree)	210	210	205	207	207
Cloud Cover(Octa)	7	7	6	8	8

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 24.0°C and 25.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 13 to 18 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.
SUGARCANE	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of nitrogen and 112 ½kg of K2O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm).
Live Stock(Varieties)	Live Stock Specific Advisory
Horticulture Specific Advisory:	
Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the
Live Stock Specific Advisory:	

Live Stock (Varieties)	60th day with another 12 packets (2400gm). Live Stock Specific Advisory
Horticulture Specific Advisory: Horticulture (Varieties) BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
Live Stock Specific Advisory: COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.
GOAT	Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **ARIMALAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	0.0	1.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	24.0	25.0	25.0	25.0	24.0
RH-I(%)	70	70	70	70	70
RH-II(%)	32	41	41	41	41
Wind Speed(kmph)	18	16	14	16	12
Wind Direction(Degree)	205	205	203	204	204
Cloud Cover(Octa)	7	7	6	8	8

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 24.0°C and 25.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 12 to 18 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.
SUGARCANE	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of nitrogen and 112 ½kg of K2O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm).
Live Stock(Varieties)	Live Stock Specific Advisory
Horticulture Specific Advisory:	
Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the
Live Stock Specific Advisory:	

<p>Live Stock (Varieties)</p>	<p>60th day with another 12 packets (2400gm). Live Stock Specific Advisory</p>
<p>Horticulture Specific Advisory: Horticulture (Varieties) BANANA</p>	<p>Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.</p>
<p>Live Stock Specific Advisory: COW</p>	<p>Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.</p>
<p>GOAT</p>	<p>Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.</p>

Others (Soil / Land Preparation) Specific Advisory:

<p>Others (Soil / Land Preparation) (Varieties)</p>	<p>Others (Soil / Land Preparation) Specific Advisory</p>
<p>GENERAL ADVICE</p>	<p>Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.</p>



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **AVADAIYARKOVIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	0.0	1.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	25.0	25.0	25.0	25.0	24.0
RH-I(%)	70	70	70	72	70
RH-II(%)	37	44	44	44	42
Wind Speed(kmph)	18	17	16	16	14
Wind Direction(Degree)	214	213	205	207	207
Cloud Cover(Octa)	6	7	6	8	7

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 24.0°C and 25.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 14 to 18 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.
SUGARCANE	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of nitrogen and 112 ½kg of K2O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm).
Live Stock(Varieties)	Live Stock Specific Advisory
Horticulture Specific Advisory:	
Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the
Live Stock Specific Advisory:	

Live Stock (Varieties)	60th day with another 12 packets (2400gm). Live Stock Specific Advisory
Horticulture Specific Advisory: Horticulture (Varieties) BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
Live Stock Specific Advisory: COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.
GOAT	Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **GANDARAVAKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	2.0	3.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	26.0	26.0	26.0	25.0	25.0
RH-I(%)	70	70	70	70	70
RH-II(%)	48	49	49	49	49
Wind Speed(kmph)	19	19	18	18	16
Wind Direction(Degree)	204	206	204	205	205
Cloud Cover(Octa)	7	7	7	8	8

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 25.0°C and 26.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 16 to 19 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.
SUGARCANE	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of nitrogen and 112 ½kg of K2O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm).
Live Stock(Varieties)	Live Stock Specific Advisory
Horticulture Specific Advisory:	
Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the
Live Stock Specific Advisory:	

Live Stock (Varieties)	60th day with another 12 packets (2400gm). Live Stock Specific Advisory
Horticulture Specific Advisory: Horticulture (Varieties) BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
Live Stock Specific Advisory: COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.
GOAT	Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **KARAMBAKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	0.0	2.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	25.0	25.0	25.0	25.0	24.0
RH-I(%)	70	70	70	70	70
RH-II(%)	40	45	45	45	45
Wind Speed(kmph)	18	17	16	17	15
Wind Direction(Degree)	209	211	207	210	210
Cloud Cover(Octa)	7	7	7	8	8

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 24.0°C and 25.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 15 to 18 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.
SUGARCANE	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of nitrogen and 112 ½kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm).
Live Stock(Varieties)	Live Stock Specific Advisory
Horticulture Specific Advisory:	
Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the
Live Stock Specific Advisory:	

Live Stock (Varieties)	60th day with another 12 packets (2400gm). Live Stock Specific Advisory
Horticulture Specific Advisory Horticulture (Varieties) BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
Live Stock Specific Advisory COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.
GOAT	Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **KUNNANDARKOIL** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	0.0	2.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	24.0	25.0	25.0	25.0	24.0
RH-I(%)	70	70	70	70	70
RH-II(%)	33	42	42	42	41
Wind Speed(kmph)	16	16	15	14	13
Wind Direction(Degree)	227	237	226	236	236
Cloud Cover(Octa)	7	7	6	8	8

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 24.0°C and 25.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 13 to 16 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
Live Stock(Varieties)	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm).
BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the

Live Stock (Varieties) BANANA	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm). Live Stock Specific Advisory
COW BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Maneozeb and spray.
COW Live Stock Specific Advisory:	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.
GOAT	Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **MANALMELKUDI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	0.0	2.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	25.0	25.0	25.0	25.0	24.0
RH-I(%)	70	70	70	72	70
RH-II(%)	39	45	45	45	43
Wind Speed(kmph)	18	18	16	16	15
Wind Direction(Degree)	216	214	206	208	208
Cloud Cover(Octa)	6	7	7	8	7

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 24.0°C and 25.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 15 to 18 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
Live Stock(Varieties)	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm).
BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the
BANANA	
COW	

Live Stock (Varieties) BANANA	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm). Live Stock Specific Advisory
COW BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Maneozeb and spray.
COW Live Stock Specific Advisory:	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.
GOAT	Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **PONNAMARAVATI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	0.0	1.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	25.0	25.0	25.0	25.0	24.0
RH-I(%)	70	70	70	72	70
RH-II(%)	35	43	43	43	41
Wind Speed(kmph)	17	17	15	15	13
Wind Direction(Degree)	216	217	208	211	211
Cloud Cover(Octa)	7	7	6	8	8

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 24.0°C and 25.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 13 to 17 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
Live Stock(Varieties)	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm).
BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the
BANANA	
COW	

Live Stock (Varieties) BANANA	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm). Live Stock Specific Advisory
COW BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Maneozeb and spray.
COW Live Stock Specific Advisory:	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.
GOAT	Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **PUDUKKOTTAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	0.0	1.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	24.0	25.0	25.0	25.0	24.0
RH-I(%)	70	70	70	70	70
RH-II(%)	30	40	40	40	40
Wind Speed(kmph)	18	16	14	16	12
Wind Direction(Degree)	200	200	200	200	200
Cloud Cover(Octa)	7	7	6	8	8

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 24.0°C and 25.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 12 to 18 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
Live Stock(Varieties)	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm).
BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the

Live Stock (Varieties) BANANA	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm). Live Stock Specific Advisory
COW BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Maneozeb and spray.
COW Live Stock Specific Advisory:	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.
GOAT	Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **THIRUVARANGULAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	0.0	1.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	24.0	25.0	25.0	25.0	24.0
RH-I(%)	70	70	70	70	70
RH-II(%)	30	40	40	40	40
Wind Speed(kmph)	18	16	14	16	12
Wind Direction(Degree)	201	201	201	201	201
Cloud Cover(Octa)	7	7	6	8	8

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 24.0°C and 25.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 12 to 18 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
Live Stock(Varieties)	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Live Stock Specific Advisory
BANANA	Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm).
COW	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
BANANA	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the
COW	

Live Stock (Varieties) BANANA	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm). Live Stock Specific Advisory
COW BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Maneozeb and spray.
COW Live Stock Specific Advisory:	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.
GOAT	Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **TIRUMAYAM** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	0.0	1.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	24.0	25.0	25.0	25.0	24.0
RH-I(%)	70	70	70	71	70
RH-II(%)	32	42	42	42	41
Wind Speed(kmph)	18	16	15	16	13
Wind Direction(Degree)	206	207	203	204	204
Cloud Cover(Octa)	7	7	6	8	8

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 24.0°C and 25.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 13 to 18 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
Live Stock(Varieties)	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Live Stock Specific Advisory
BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Repeat the same on the 60th day with another 12 packets (2400 gm). If necessary, mix 2 g of Mancozeb and spray.
COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the
BANANA	
COW	

Live Stock (Varieties) BANANA	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of K ₂ O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm). Live Stock Specific Advisory
COW BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Maneozeb and spray.
COW Live Stock Specific Advisory:	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.
GOAT	Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.



Gramin Krishi Mausam Sew

Experimental Block Level Agromet Advisory Bulletin
(A Joint Initiative of IMD & ICAR)



Agromet Advisory Bulletin

Date : 29-09-2023

Weather Forecast of **VIRALIMALAI** Block in **PUDUKKOTTAI**(Tamil Nadu) Issued On :2023-09-29(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-30	2023-10-01	2023-10-02	2023-10-03	2023-10-04
Rainfall	0.0	0.0	0.0	0.0	2.0
Tmax(°C)	35.0	35.0	36.0	35.0	35.0
Tmin(°C)	24.0	25.0	25.0	25.0	24.0
RH-I(%)	70	70	70	71	70
RH-II(%)	33	42	42	42	41
Wind Speed(kmph)	16	16	15	14	13
Wind Direction(Degree)	230	241	228	239	239
Cloud Cover(Octa)	7	7	6	8	8

Weather Summary/Alert:

Chances of Dry weather Next five days. Maximum temperature would be in the range 35.0°C and 36.0°C and the minimum temperature would be in between 24.0°C and 25.0°C. Cloudy sky is expected for next 5 days. Wind speed is expected to be 13 to 16 kmph.

General Advisory:

Farmers are advised to fumigate the goat shed With Notchi leaves to prevent fly attack.

SMS Advisory:

Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
BLACK GRAM	Prevailing present weather condition the incense of Yellow mosaic in pulses. To manage this disease, farmers are advised to set up yellow sticky trap @ 5 No. / acre to monitor the movements of the vector whitefly and if the vector population is noticed farmers are advised to spray Imidacloprid 2ml per litre of water to manage the vector white fly.
RICE	The incidence of false smut may be noticed in rice. The damage symptoms are individual rice grain transformed into a mass of yellow fruiting bodies and infected grain has greenish smut balls with a velvety appearance. If symptoms are observed in field, spray Propiconazole 25 EC @ 200 ml/ac or Copper hydroxide 77% WP 500 g/ac with 200 litre of water.
SUGARCANE	The fertilizer application process can be done for the Sugarcane crop on non-rainy days. Apply 225 kg of nitrogen and 112 ½kg of K2O/ha in three equal splits at 30, 60 and 90 days. Mix 12 packets (2400g/ha) of Azospirillum with 25kg of FYM and 25kg soil and apply near the clumps on the 30th day of planting. Repeat the same on the 60th day with another 12 packets (2400gm).
Live Stock(Varieties)	Live Stock Specific Advisory
Horticulture Specific Advisory:	
Horticulture(Varieties)	Horticulture Specific Advisory
BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the
Live Stock Specific Advisory:	

Live Stock (Varieties)	60th day with another 12 packets (2400gm). Live Stock Specific Advisory
Horticulture Specific Advisory: Horticulture (Varieties) BANANA	Due to the current rains, green fodder is available in large quantities so it is likely that cattle will consume more fodder. Due to this, the risk of diarrhea and acidosis is high. Therefore farmers should reduce greenery to their livestock and give more dry fodder. Due to high clouds and humidity in the air, cigarette butts are prone to gray rot. If necessary, mix 2 g of Mancozeb and spray.
Live Stock Specific Advisory: COW	Once in a month, periodical de-worming is essential for Calves. Check the cow dung often, so that we can detect the worm infection if present. Farmers are advised to consult the nearby veterinarian for deworming. Periodical de-worming reduces the worm infection at calves and also induces the growth, weight & puberty.
GOAT	Goat rearing farmers are advised to prevent the goat from mortality, as prevention against Haemorrhagic Septicemia, goats have to be vaccinated once a year before the onset of the rainy season, contact the nearby veterinarian and vaccinate your goats.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation) (Varieties)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	Nitrogen fertilizers should be mixed with neem paste or mixed with tar so that the water and crop nutrient absorption is less as the clouds and air humidity are high for the last few days. Thus the utilization capacity of the nutrients will be higher.