

Pandit Jawaharlal Nehru College of Agriculture and<br/>Research Institute, Karaikal - 609 603Agromet Advisory Bulletin (AAB) for KARAIKAL District<br/>Issued jointly by IMD & PAJANCOA & RI<br/>Forecast Period: 21st to 25th August 2024



Phone No. 04368 – 261182 E-mail: agronaas2012@gmail.com

Fax: 04368-261260

Bulletin No. 067 (08)/PJN/GKMS/2024

Date: 20.08.2024

## Weather Summary for Karaikal district

	Past w	eek Sum	mary		Weather	Ensemble weather forecast valid until				
	(15.08.20)	24 to 19.0	08.2024)		Parameters	08.30 hrs of 25.08.2024				
15/08	16/08	17/08	18/08	19/08		Day-1 21/08	Day-2 22/08	Day-3 23/08	Day-4 24/08	Day-5 25/08
0	0	0	7.2	8.5	Rainfall (mm)	8	4	0	0	0
35	35	35	35	35	Max. Temp. (° C)	33	32	33	33	34
27	27	27	27	27	Mini. Temp. (° C)	26	25	25	26	26
4	3	4	1	4	Cloud cover (Octa)	8	8	5	4	6
57-91	59-85	69-86	69-86	67-94	RH Even- Morn. (%)	60-80	50-70	50-70	50-70	42-60
4	5	6	3	5	Wind (kmph)	30	32	30	26	22
ESE	SW	SE	SW	SW	Wind direction	SSW	SSW	SSW	SSW	SW

## Forecast for next five days: 21.08.2024 to 25.08.2024

Light rain is expected on  $21^{st}$  and  $22^{nd}$  August.

During next five days the sky will be cloudy. The temperature range is expected to be around 25-34 °C. Relative humidity range is expected to be around 42-80%. The wind speed is expected to be 22-32 km per hour and the wind will be mostly from south south west direction.

Rainfall (7.7 mm) will be above normal, maximum temperature (31.5°C) and minimum temperature (23.5°C) will be normal from  $25^{\text{th}}$  to  $31^{\text{st}}$  August.

Crop/livestock	Stage/Particulars	Agro Advisory
Cotton	Flowering to picking stage	To manage mealy bug in cotton, spray Thiamethoxam 25 WG @ 0.2 g/l or Buprofezin 25 SC @ 2 ml/l. To manage white fly in cotton, spray Azadirachtin 0.03 (300 ppm) @ 5-10 ml/l or Thiamethoxam 25 WG 0.4 g/l.
Brinjal	Bearing stage	To manage to bacco cutworm in brinjal, spray Bacillus thuring iensis 2 g/l or Novaluron 10% EC 1 ml/l or flubendamide 0.2 - 0.3 ml/l
Cattle	Babesiasis	The prevailing weather conditions are favorable for the Babesiasis disease in cattle. High fever, dark coloured

## Agro Advisory

urine, less intake of feed and less milk yield are the symptoms of Babesiasis disease. If these symptoms appear in the cattle, contact nearby dispensary. As a preventive measure, apply Butox @ 2 ml/l of water on
cattle and also spray it on the nearby areas of the cattle
shed to control ectoparasites

SMS: Light rain is expected for next few days. Hence, postpone irrigation to crops.

DEAN PAJANCOA & RI