



## Agromet Advisory Bulletin for the **Thiruvavarur** District

(From 03<sup>rd</sup> August 2024 to 07<sup>th</sup> August 2024)

Issued jointly by Tamil Nadu Rice Research Institute -  
Aduthurai and IMD



**Bulletin no: 36/2024**

**Date: 02.08.2024**

Weather Summary for the period since the issue of last bulletin no. 35 (30.07.2024) for Tamil Nadu: Thiruvavarur district

Maximum temperature of 35 - 39 °C, minimum temperature of 29 - 30 °C, and was recorded in Thiruvavarur district.

### **Weather forecast for next five days (03.08.2024 to 07.08.2024)**

Past week Summary 30.07.2024 to 01.08.2024			Parameters	Ensemble weather forecast valid until 08.30 hr of 07.08.2024				
Day-1 30/07	Day-2 31/07	Day-3 01/08		Date	Day-1 03/08	Day-2 04/08	Day-3 05/08	Day-4 06/08
0	0	0	Rainfall (mm)	2	2	4	2	5
35	38	39	Max. Temp. (°C)	39	39	38	38	38
29	30	29	Min. Temp. (°C)	30	30	29	29	29
5	5	5	Cloud cover (Octa)	5	6	6	7	8
69-51	70-52	72-57	RH I-II (%)	60-40	60-40	60-50	70-40	70-40
5	6	6	Wind (kmph)	14	12	12	14	14
West South West	West South West	West South West	Wind direction	South	South West	West South West	East South East	East South East

### **Weather Summary and forecast for Thiruvavarur district**

According to district level forecast issued by India Meteorological Department for Thiruvavarur district, the sky will be clear. Light drizzling expected on 03.08.2024 to 07.08.2024. Maximum temperature is expected to be around 38 - 39 °C and minimum temperature is expected to be around 29 - 30 °C. Morning relative humidity is expected to be around 60 - 70 percent and evening relative humidity is expected to be around 40 - 50 percent. Average wind speed is expected to be around 12 - 14 km per hour. The wind direction will be east south east.

### **Agro Advisory**

Crop	Stage	Advisory
------	-------	----------

General		Farmers are requested to download and utilize the agricultural based mobile apps suggested by TNAU (TNAU AAS, Meghdoot and DAMINI).
Rice	Main field	The prevailing weather condition favorable for gall midge infection in rice. It leads to the formation of onion like leaf and prevents the panicle formation. To control this apply chloripyriphos 20 EC 500 ml/ ha.
Rice	Main field	At active tillering stage of Kuruvai paddy, top dressing can be done with 35 kg urea and 10 kg MOP/acre by keeping thin film of water in the field.
Banana		Provide support with wooden poles to avoid lodging of tree due to wind.
Cattle	Cow	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.

### **SMS advisory**

Owing to moderate wind, more moisture loss will be expected and hence, more irrigation should be given.

**Director  
TRRI, Aduthurai**