

District:

Bulletin No:

**Dindigul** 

MSSRF/ DGL /001/ 2024

## AgroMet Advisory Bulletin (AAB)

Jointly released by

Regional Meteorology Centre, Chennai – India Meteorology Department,
Reddiarchathram Seed Growers Association &

M. S. Swaminathan Research Foundation, Kannivadi

Forecast Period:

Release Date :

Phone no.: 044 - 22541229 E mail: amfu.kanniwadi@gmail.com Fax: 044 - 22541319



07/01/2024

Significant past weather	From	29/12/202	to	01/01/202	
Parameters		Past Weather			
Rainfall (mm)		0.0			
Maximum Temperature (°C)		31-32			
Minimum Temperature (°C)			23		
Sky Condition (Octas)			4-6		
Relative Humidity (%)			58-83		
Wind Speed (kmph)			5-9		

Ensemble Weather Forecast until

08:30Hrs of

07-01-2024

03/01/2024

02/01/2024

Parameters	03-01- 2024	04-01- 2024	05-01- 2024	06-01- 2024	07-01- 2024
	Wednesday	Thursday	Friday	Saturday	Sunday
Rainfall (mm)	3	4	6	6	4
Maximum Temperature (°C)	32	31	31	29	29
Minimum Temperature (°C)	23	22	21	21	20
Sky Condition (Octa)	6	7	4	7	8
Max. Relative Humidity (%)	90	90	90	90	90
Min. Relative Humidity (%)	70	70	70	70	70
Wind Speed (Kmph)	8	6	6	6	8
Wind Direction	NE	E	E	NE	E

SMS Advisory Chances of light rain during 3, 4, 5, 6 & 7 Jan. Maximum temperature would be in the range of 29°C to 32°C and the minimum temperature would be between 20°C-23°C. Generally cloudy sky will appear for the next five days. Wind expected with the speed of 6-8 Kmph from, Northeast & East directions.

## General Advisory:

Since very less amount of rainfall is forecasted for next 5 days, Maize & Cotton cultivators are advised to plan for harvest.

Since cloudy humid weather is prevailing, farmers are advised to use the yellow sticky trap @ 10-15/acre to monitor the incidence of sucking pests and Blue sticky trap @ 5-10/acre to monitor the aphids and leaf miner.

Weather Based Agro-Advisories: (Guava, Beans & Animal Husbandry). Guava: The incidence of Fruit Flies is noticed in Guava. The symptoms of damage appear on fruits. The fruit fly adults and maggots attack the semi-ripened fruits, ovipositor punctures are found on fruits, maggots destroy the pulp and a bad smell comes out of it.

To control this, set up a pheromone trap. Fill a 5-litre bucket with 3 litres water + 2 dry fish + 50gm Jaggery + 2 well-rotten Bananas + 1kg castor cake + 2 pieces of Pineapple and fix it as 12 Nos per Acre.

## Beans:

There are chances for the incidence of Powdery mildew in Beans crop. The symptom of damage is small cottony spots appearing on the infected leaves. Defoliation of severely infected plants occurs before maturity.

To prevent this, spray Pseudomonas @ 60gms + Trichoderma Viride @ 60gms per 10litres of water along with 10ml of Sticky agent, 2 times in 10 days interval.

During severe infection, spray Tebuconazole @ 100ml per acre with sticky agent @ 1ml per 1 litre of water.

## **Animal Husbandry:**

As the night temperature is very low, keep the cattle safely in the sheds.

Tie the jute bags around the shed. Allow the cattle to graze 2 hours after sunrise.