

Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal - 609 603

Agromet Advisory Bulletin (AAB) for KARAIKAL District Issued jointly by IMD & PAJANCOA & RI





Fax: 04368-261260





Date: 09.07.2024

Weather Summary for Karaikal district

| Past week Summary | | | | Weather | Ensemble weather forecast valid until | | | | | |
|----------------------------|-------|-------|-------|------------|---------------------------------------|----------------|----------------|----------------|----------------|----------------|
| (04.07.2024 to 08.07.2024) | | | | Parameters | 08.30 hrs of 14.07.2024 | | | | | |
| 04/07 | 05/07 | 06/07 | 07/07 | 08/07 | | Day-1 10/07 | Day-2 11/07 | Day-3 12/07 | Day-4 13/07 | Day-5 14/07 |
| 0 | 0 | 0 | 0 | 0 | Rainfall (mm) | 3 | 2 | 3 | 13 | 4 |
| 39 | 39 | 39 | 39 | 39 | Max. Temp. (°C) | 35 | 34 | 34 | 33 | 33 |
| 27 | 27 | 26 | 27 | 27 | Mini. Temp. (°C) | 27 | 26 | 26 | 25 | 25 |
| 3 | 3 | 4 | 3 | 4 | Cloud cover (Octa) | 7 | 6 | 6 | 8 | 8 |
| 54-83 | 50-87 | 54-80 | 50-87 | 60-87 | RH Even- Morn. (%) | 50-70 | 50-60 | 50-70 | 60-80 | 50-70 |
| 6 | 5 | 6 | 4 | 6 | Wind (kmph) | 18 | 20 | 18 | 16 | 18 |
| WSW | SW | WSW | SW | SW | Wind direction | SSW | S | S | SW | SW |

Forecast for next five days: 10.07.2024 to 14.07.2024

Light to moderate rain is expected for next five days.

During next five days the sky will be cloudy. The temperature range is expected to be around 25-35°C. Relative humidity range is expected to be around 50-80%. The wind speed is expected to be 16-20 km per hour and the wind will be mostly from south west direction.

Rainfall (2.9 mm), maximum temperature (32.8°C) and minimum temperature (24.1°C) will be normal from 14^{th} to 20^{th} July.

Agro Advisory

| 11610114110 | - 🧸 | | | |
|----------------|----------------------|---|--|--|
| Crop/livestock | Stage/Particulars | Agro Advisory | | |
| _ | Flowering to picking | To manage mealy bug in cotton, spray Thiamethoxam 25 WG @ 0.2 g/l or Buprofezin 25 SC @ 2 ml/l. | | |
| Cotton | stage | To manage white fly in cotton, spray Azadirachtin 0.03 (300 ppm) @ 5-10 ml/l or Thiamethoxam 25 WG 0.4 g/l. | | |
| Brinjal | Bearing stage | To manage tobacco cutworm in brinjal, spray Bacillus thuringiensis 2 g/l or Novaluron 10% EC 1 ml/l or flubendamide 0.2 – 0.3 ml/l | | |
| Cattle | Babesiasis | The prevailing weather conditions are favorable for the Babesiasis disease in cattle. High fever, dark coloured urine, less intake of feed and less milk yield are the symptoms of Babesiasis disease. If these symptoms appear in the cattle, contact nearby dispensary. As a preventive measure, apply Butox @ 2 ml/l of water on | | |

| | cattle and also spray it on the nearby areas of the cattle |
|--|--|
| | shed to control ectoparasites |

SMS: Light to moderate rain is expected for next few days. Hence, postpone irrigation to crops.

DEAN PAJANCOA & RI