

Agromet Advisory Bulletin for Thanjavur District (From 20th July 2024 to 24th July 2024)



Issued jointly by Tamil Nadu Rice Research Institute - Aduthurai and IMD

Bulletin no: 32/2024

Date: 19.07.2024

Weather Summary for the period since the issue of last bulletin no. 31 (16.07.2024) for Tamil Nadu: Thanjavur district

Maximum temperature of 31 - 35 $^{\circ}$ C, minimum temperature of 27 - 28 $^{\circ}$ C, and isolated rainfall was recorded in Thanjavur district.

Weather forecast for next five days (20.07.2024 to 24.07.2024)

Past week Summary 16.07.2024 to 18.07.2024			Parameters	Ensemble weather forecast valid until 08.30 hr of 24.07.2024				
Day-1	Day-2	Day-3		Day-1	Day-2	Day-3	Day-4	Day-5
16/07	17/07	18/07	Date	20/07	21/07	22/07	23/07	24/07
0.5	0.2	1.2	Rainfall (mm)	2	4	5	3	0
33	31	35	Max. Temp. (°C)	34	34	35	35	35
28	28	27	Min. Temp. (°C)	27	27	28	28	28
6	6	6	Cloud cover (Octa)	6	6	5	5	5
76-71	78-64	77-77	RH I-II (%)	78-68	78-68	74-64	74-64	74-64
6	5	6	Wind (kmph)	20	18	22	20	20
West South West	West South West	West South West	Wind direction	West	West South West	West	West	West

Weather Summary and forecast for Thanjavur district

According to district level forecast issued by India Meteorological Department for Thanjavur district, the sky will be clear. Light rainfall expected on 20.07.2024 to 23.07.2024. Maximum temperature is expected to be around 34 - 35 °C and minimum temperature is expected to be around 27 - 28 °C. Morning relative humidity is expected to be around 74 - 78 percent and evening relative humidity is expected to be around 64 - 68 percent. Average wind speed is expected to be around 18 - 22 km per hour. The wind direction will be west.

Agro Advisory

Crop	Stage	Advisory			
		Farmers are requested to download and utilize the agricultural			
		based mobile apps suggested by TNAU (TNAU AAS, Meghdoot and			
Ger	neral	DAMINI).			
		Owing to moderate wind, more moisture loss will be expected and			
		hence, more irrigation should be given.			
		The prevailing weather condition favours the stem borer incidence			
		in paddy. Place pheromone traps @ 4 / ac to control stem borer			
Rice	Main field	population. Also to control this, spray profenophos at 400			
		ml/acre.(or) Apply Cartarp hydrochloride 4 G @ 6 kg/ac for			
		effective control of stemborer.			
	Cow	Present weather is conducive for the initiation of foot and mouth			
Cattle		disease in dairy animals and cattle. All farmers must vaccinate the			
		animals to control the disease.			

SMS advisory

Irrigate crop abundantly at short interval to avoid moisture loss due to wind.

Director TRRI, Aduthurai