

Agromet Advisory Bulletin for the Chennai District Bulletin no. 009/2025 Issued on 31.01.2025 (01.02.2025-05.02.2025)

Issued jointly by



Tamil Nadu Veterinary and Animal Sciences University & India Meteorological Department

Weather Summary for the period since the issue of last bulletin for Chennai.

During past three days around Chennai area, the sky was mostly clear. Maximum temperature was 30.0-31.0°C and Minimum temperature was 23.0°C. Wind speed was 05-06 km per hour. Relative humidity was recorded between 63-89 %.

	t week Sumn 1.2025-30.01.	•	Parameters	Ensemble weather forecast valid until 08.30 hrs of 05.02.2025				
28.01.2025	29.01.2025	30.01.2025	Date	01.02.2025	02.02.2025	03.02.2025	04.02.2025	05.02.2025
0.0	0.0	0.0	Rainfall (mm)	0	2	0	0	0
31	31	30	Max. Temp. (° C)	31	31	31	32	32
23	23	23	Mini. Temp. (° C)	23	23	23	24	24
89	84	76	Maximum RH (%)	76	76	76	84	84
63	67	74	Minimum RH (%)	67	67	67	74	74
Partly cloudy	Partly cloudy	Partly cloudy	Cloud cover (Octa)	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy
North East	North East	East North East	Wind direction (Degree)	South East	South East	East North East	East	East South East
5	6	6	Wind speed Kmp	4	4	14	14	12

Weather Summary for Chennai district

Weather forecast for next five days: (01.02.2025-05.02.2025)

According to district forecast issued by the India Meteorological Department for Chennai district, sky will be mostly clear. Maximum temperature is expected to be around 31.0-32.0°C. Minimum temperature is expected to be around 23.0-24.0°C. Morning relative humidity is expected to be around 76-84 per cent and evening relative humidity is expected to be around 67-74 per cent. Wind speed is expected to be around 04-14 km per hour and the wind direction will be from East South East.

SMS Advisory:

(01.02.2025-05.02.2025) Mist/ Haze conditions are very likely to prevail at isolated pockets of Tamil Nadu during morning hours.

- 1. Till the inclement weather the livestock must not be allowed for grazing in early morning hours.
- 2. Drinking water must be warmed to body temperature so as to avoid drip in body temperature.
- 3. The livestock must be provided with vitamins and mineral supplementation to maintain good health during cold weather.
- 4. Antibiotics, oral rehydration solution, intravenous fluids must be stored in handy during emergency
- 5. Pregnant animals nearing parturition should be housed separately in clean dry shelter and must be protected from clod stress by covering the side walls

- 6. The young calf shed must be adequately bedded to prevent clod stress which is very detrimental to the young stock
- 7. Rabbit may be allowed by mating now due to conductive environment rabbit will kindle more bunnies during these months due to which necessary reproductive management must be followed.
- 8. Newly born piglets must be protected from cold drafts
- 9. Add 5%shell grit in laying hens for optimum egg production.

Rice:

Due to prevailing weather conditions leaf folder and stem borer infestation found in paddy crop. To manage them, spray carbosulfan 400 ml/ac or cartap hydrochloride 600 g/ac during morning or evening hours.

The forecasted weather may cause stem borer and leaf folder infestation in rice crop. The farmers are advised to monitor the field, if symptoms are observed it's advised to spray Chlorantraniliprole 18.5% 150 ml/ha.

Groundnut:

Due to prevailing weather conditions, there are chances of tobacco caterpillar in groundnut. To control install pheromone trap 5 nos/ac. Spray flubendiamide 20 WG 120 gram/ac or Quinalphos 20 AF @ 500ml/ac.

Blackgram:

Farmers are advised to treat the seeds with Trichoderma viride @ 4g/kg of seed (or) Pseudomonas fluorescens @ 10 g/kg seed before sowing.

Brinjal:

Application of 1.5% foliar spray of PPFM at the flowering stage in Brinjal, will prevent the premature falling of flowers and increase the numbers and quality of fruits.

Banana:

Prevailing weather condition is favorable for sigatoka leaf spot infection in banana. To control spray carbendazim 1.0 g/l or propiconazole 1.0 ml/l or mancozeb 2.5 g/l and sticking agent 1.0 ml/l at 10-15 day intervals.

Prevailing weather condition is favorable for panama wilt and nematode infection in banana. To control apply <u>Bacillus subtilis</u> 1.0 kg/ha + Paecilomyceslilacinus 1.0 kg/ha at 2, 4 and 6th months of planting.

.



Agromet Advisory Bulletin for the Tiruvallur District Bulletin no. 009/2025 Issued on 31.01.2025 (01.02.2025-05.02.2025)

Issued jointly by



Tamil Nadu Veterinary and Animal Sciences University & India Meteorological Department

Weather Summary for the period since the issue of last bulletin for Tiruvallur.

During past three days around Tiruvallur area, the sky was mostly clear. Maximum temperature was 30.0-31.0°C and Minimum temperature was 23.0°C. Wind speed was 05-06 km per hour. Relative humidity was recorded between 63-89 %.

Past week Summary (28.01.2025-30.01.2025)		Parameters	Ensemble weather forecast valid until 08.30 hrs of 05.02.2025					
28.01.2025	29.01.2025	30.01.2025	Date	01.02.2025	02.02.2025	03.02.2025	04.02.2025	05.02.2025
0.0	0.0	0.0	Rainfall (mm)	0	10	0	0	0
31	31	30	Max. Temp. (° C)	31	31	31	32	32
23	23	23	Mini. Temp. (° C)	23	23	23	24	24
89	84	76	Maximum RH (%)	76	76	76	84	84
63	67	74	Minimum RH (%)	67	67	67	74	74
Partly cloudy	Partly cloudy	Partly cloudy	Cloud cover (Octa)	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy
North East	North East	East North East	Wind direction (Degree)	South East	South East	East North East	East	East South East
5	6	6	Wind speed Kmp	4	4	14	14	12

Weather Summary for Tiruvallur district

Weather forecast for next five days: (01.02.2025-05.02.2025)

According to district forecast issued by the India Meteorological Department for Tiruvallur district, sky will be mostly clear. Maximum temperature is expected to be around 31.0-32.0°C. Minimum temperature is expected to be around 23.0-24.0°C. Morning relative humidity is expected to be around 76-84 per cent and evening relative humidity is expected to be around 67-74 per cent. Wind speed is expected to be around 04-14 km per hour and the wind direction will be from East South East.

SMS Advisory:

(01.02.2025-05.02.2025) Mist/ Haze conditions are very likely to prevail at isolated pockets of Tamil Nadu during morning hours.

- 1. Till the inclement weather the livestock must not be allowed for grazing in early morning hours.
- 2. Drinking water must be warmed to body temperature so as to avoid drip in body temperature.
- 3. The livestock must be provided with vitamins and mineral supplementation to maintain good health during cold weather.
- 4. Antibiotics, oral rehydration solution, intravenous fluids must be stored in handy during emergency
- 5. Pregnant animals nearing parturition should be housed separately in clean dry shelter and must be protected from clod stress by covering the side walls
- 6. The young calf shed must be adequately bedded to prevent clod stress which is very detrimental to the young stock

- 7. Rabbit may be allowed by mating now due to conductive environment rabbit will kindle more bunnies during these months due to which necessary reproductive management must be followed
- 8. Newly born piglets must be protected from cold drafts
- 9. Add 5%shell grit in laying hens for optimum egg production.

Rice:

Due to prevailing weather conditions leaf folder and stem borer infestation found in paddy crop. To manage them, spray carbosulfan 400 ml/ac or cartap hydrochloride 600 g/ac during morning or evening hours.

The forecasted weather may cause stem borer and leaf folder infestation in rice crop. The farmers are advised to monitor the field, if symptoms are observed it's advised to spray Chlorantraniliprole 18.5% 150 ml/ha.

Groundnut:

Due to prevailing weather conditions, there are chances of tobacco caterpillar in groundnut. To control install pheromone trap 5 nos/ac. Spray flubendiamide 20 WG 120 gram/ac or Quinalphos 20 AF @ 500ml/ac.

Blackgram:

Farmers are advised to treat the seeds with Trichoderma viride @ 4g/kg of seed (or) Pseudomonas fluorescens @ 10 g/kg seed before sowing.

Brinjal:

Application of 1.5% foliar spray of PPFM at the flowering stage in Brinjal, will prevent the premature falling of flowers and increase the numbers and quality of fruits.

Banana:

Prevailing weather condition is favorable for sigatoka leaf spot infection in banana. To control spray carbendazim 1.0 g/l or propiconazole 1.0 ml/l or mancozeb 2.5 g/l and sticking agent 1.0 ml/l at 10-15 day intervals.

Prevailing weather condition is favorable for panama wilt and nematode infection in banana. To control apply <u>Bacillus subtilis</u> 1.0 kg/ha + Paecilomyceslilacinus 1.0 kg/ha at 2, 4 and 6th months of planting.



Agromet Advisory Bulletin for the Thiruvannamalai District Bulletin no. 009/2025 Issued on 31.01.2025 (01.02.2025-05.02.2025)

Issued jointly by



Tamil Nadu Veterinary and Animal Sciences University & India Meteorological Department

Weather Summary for the period since the issue of last bulletin for TVMalai.

During past three days around TVMalai area, the sky was mostly clear. Maximum temperature was 30.0-31.0°C and Minimum temperature was 18.0-19.0°C. Wind speed was 03-04 km per hour. Relative humidity was recorded between 64-91 %.

Past week Summary (28.01.2025-30.01.2025)			Parameters	Ensemble weather forecast valid until 08.30 hrs of 05.02.2025				
28.01.2025	29.01.2025	30.01.2025	Date	01.02.2025	02.02.2025	03.02.2025	04.02.2025	05.02.2025
0.0	0.0	0.0	Rainfall (mm)	6	10	0	0	0
31	30	31	Max. Temp. (° C)	30	30	30	31	31
18	19	19	Mini. Temp. (° C)	21	21	21	23	23
87	91	87	Maximum RH (%)	87	87	87	91	91
64	64	64	Minimum RH (%)	64	64	64	70	70
Partly cloudy	Partly cloudy	Partly cloudy	Cloud cover (Octa)	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy
East	East	East South East	Wind direction (Degree)	East	East South East	East	East	East South East
4	3	4	Wind speed Kmp	4	4	12	12	14

Weather Summary for TVMalai district

Weather forecast for next five days: (01.02.2025-05.02.2025)

According to district forecast issued by the India Meteorological Department for TVMalai district, sky will be mostly clear. Maximum temperature is expected to be around 30.0-31.0°C. Minimum temperature is expected to be around 21.0-23.0°C. Morning relative humidity is expected to be around 87-91 per cent and evening relative humidity is expected to be around 64-70 per cent. Wind speed is expected to be around 04-14 km per hour and the wind direction will be from East South East.

SMS Advisory:

(01.02.2025-05.02.2025) Mist/ Haze conditions are very likely to prevail at isolated pockets of Tamil Nadu during morning hours.

- 1. Till the inclement weather the livestock must not be allowed for grazing in early morning hours.
- 2. Drinking water must be warmed to body temperature so as to avoid drip in body temperature.
- 3. The livestock must be provided with vitamins and mineral supplementation to maintain good health during cold weather.
- 4. Antibiotics, oral rehydration solution, intravenous fluids must be stored in handy during emergency
- 5. Pregnant animals nearing parturition should be housed separately in clean dry shelter and must be protected from clod stress by covering the side walls
- 6. The young calf shed must be adequately bedded to prevent clod stress which is very detrimental to the young stock

- 7. Rabbit may be allowed by mating now due to conductive environment rabbit will kindle more bunnies during these months due to which necessary reproductive management must be followed
- 8. Newly born piglets must be protected from cold drafts
- 9. Add 5%shell grit in laying hens for optimum egg production.

Rice:

Due to prevailing weather conditions leaf folder and stem borer infestation found in paddy crop. To manage them, spray carbosulfan 400 ml/ac or cartap hydrochloride 600 g/ac during morning or evening hours.

The forecasted weather may cause stem borer and leaf folder infestation in rice crop. The farmers are advised to monitor the field, if symptoms are observed it's advised to spray Chlorantraniliprole 18.5% 150 ml/ha.

Groundnut:

Due to prevailing weather conditions, there are chances of tobacco caterpillar in groundnut. To control install pheromone trap 5 nos/ac. Spray flubendiamide 20 WG 120 gram/ac or Quinalphos 20 AF @ 500ml/ac.

Blackgram:

Farmers are advised to treat the seeds with Trichoderma viride @ 4g/kg of seed (or) Pseudomonas fluorescens @ 10 g/kg seed before sowing.

Brinjal:

Application of 1.5% foliar spray of PPFM at the flowering stage in Brinjal, will prevent the premature falling of flowers and increase the numbers and quality of fruits.

Banana:

Prevailing weather condition is favorable for sigatoka leaf spot infection in banana. To control spray carbendazim 1.0 g/l or propiconazole 1.0 ml/l or mancozeb 2.5 g/l and sticking agent 1.0 ml/l at 10-15 day intervals.

Prevailing weather condition is favorable for panama wilt and nematode infection in banana. To control apply <u>Bacillus subtilis</u> 1.0 kg/ha + Paecilomyceslilacinus 1.0 kg/ha at 2, 4 and 6th months of planting.



Agromet Advisory Bulletin for the Vellore District Bulletin no. 009/2025 Issued on 31.01.2025 (01.02.2025-05.02.2025)

Issued jointly by



Tamil Nadu Veterinary and Animal Sciences University & India Meteorological Department

Weather Summary for the period since the issue of last bulletin for Vellore.

During past three days around Vellore area, the sky was mostly clear. Maximum temperature was 30.0-31.0°C and Minimum temperature was 18.0-19.0°C. Wind speed was 03-04 km per hour. Relative humidity was recorded between 64-91 %.

	t week Sumn 1.2025-30.01.	•	Parameters	Ensemble weather forecast valid until 08.30 hrs of 05.02.2025				
28.01.2025	29.01.2025	30.01.2025	Date	01.02.2025	02.02.2025	03.02.2025	04.02.2025	05.02.2025
0.0	0.0	0.0	Rainfall (mm)	0	12	0	0	0
31	30	31	Max. Temp.	30	30	30	31	31
18	19	19	Mini. Temp. (° C)	21	21	21	23	23
87	91	87	Maximum RH (%)	87	87	87	91	91
64	64	64	Minimum RH (%)	64	64	64	70	70
Partly cloudy	Partly cloudy	Partly cloudy	Cloud cover (Octa)	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy
East	East	East South East	Wind direction (Degree)	East	East South East	East	East	East South East
4	3	4	Wind speed Kmp	4	4	12	12	14

Weather Summary for Vellore district

Weather forecast for next five days: (01.02.2025-05.02.2025)

According to district forecast issued by the India Meteorological Department for Vellore district, sky will be mostly clear. Maximum temperature is expected to be around 30.0-31.0°C. Minimum temperature is expected to be around 21.0-23.0°C. Morning relative humidity is expected to be around 87-91 per cent and evening relative humidity is expected to be around 64-70 per cent. Wind speed is expected to be around 04-14 km per hour and the wind direction will be from East South East.

SMS Advisory:

(01.02.2025-05.02.2025) Mist/ Haze conditions are very likely to prevail at isolated pockets of Tamil Nadu during morning hours.

- 1. Till the inclement weather the livestock must not be allowed for grazing in early morning hours.
- 2. Drinking water must be warmed to body temperature so as to avoid drip in body temperature.
- 3. The livestock must be provided with vitamins and mineral supplementation to maintain good health during cold weather.
- 4. Antibiotics, oral rehydration solution, intravenous fluids must be stored in handy during emergency
- 5. Pregnant animals nearing parturition should be housed separately in clean dry shelter and must be protected from clod stress by covering the side walls
- 6. The young calf shed must be adequately bedded to prevent clod stress which is very detrimental to the young stock

- 7. Rabbit may be allowed by mating now due to conductive environment rabbit will kindle more bunnies during these months due to which necessary reproductive management must be followed
- 8. Newly born piglets must be protected from cold drafts
- 9. Add 5%shell grit in laying hens for optimum egg production.

Rice:

Due to prevailing weather conditions leaf folder and stem borer infestation found in paddy crop. To manage them, spray carbosulfan 400 ml/ac or cartap hydrochloride 600 g/ac during morning or evening hours.

The forecasted weather may cause stem borer and leaf folder infestation in rice crop. The farmers are advised to monitor the field, if symptoms are observed it's advised to spray Chlorantraniliprole 18.5% 150 ml/ha.

Groundnut:

Due to prevailing weather conditions, there are chances of tobacco caterpillar in groundnut. To control install pheromone trap 5 nos/ac. Spray flubendiamide 20 WG 120 gram/ac or Quinalphos 20 AF @ 500ml/ac.

Blackgram:

Farmers are advised to treat the seeds with Trichoderma viride @ 4g/kg of seed (or) Pseudomonas fluorescens @ 10 g/kg seed before sowing.

Brinjal:

Application of 1.5% foliar spray of PPFM at the flowering stage in Brinjal, will prevent the premature falling of flowers and increase the numbers and quality of fruits.

Banana:

Prevailing weather condition is favorable for sigatoka leaf spot infection in banana. To control spray carbendazim 1.0 g/l or propiconazole 1.0 ml/l or mancozeb 2.5 g/l and sticking agent 1.0 ml/l at 10-15 day intervals.

Prevailing weather condition is favorable for panama wilt and nematode infection in banana. To control apply <u>Bacillus subtilis</u> 1.0 kg/ha + Paecilomyceslilacinus 1.0 kg/ha at 2, 4 and 6th months of planting.

.



Agromet Advisory Bulletin for the Villupuram District Bulletin no. 009/2025 Issued on 31.01.2025 (01.02.2025-05.02.2025)

Issued jointly by



Tamil Nadu Veterinary and Animal Sciences University & India Meteorological Department

Weather Summary for the period since the issue of last bulletin for Villupuram.

During past three days around Villupuram area, the sky was mostly clear. Maximum temperature was 30.0°C and Minimum temperature was 20.0-22.0°C. Wind speed was 03-05 km per hour. Relative humidity was recorded between 67-95 %.

	t week Sumn 1.2025-30.01.	•	Parameters Ensemble weather forecast valid until 08.30 05.02.2025					rs of
28.01.2025	29.01.2025	30.01.2025	Date	01.02.2025	02.02.2025	03.02.2025	04.02.2025	05.02.2025
0.0	0.0	0.0	Rainfall (mm)	10	8	0	0	0
30	30	30	Max. Temp. (° C)	30	30	30	31	31
20	21	22	Mini. Temp. (° C)	22	22	22	23	23
95	90	91	Maximum RH (%)	90	90	90	92	92
67	68	77	Minimum RH (%)	68	68	68	77	77
Partly cloudy	Partly cloudy	Partly cloudy	Cloud cover (Octa)	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy
East North East	North North East	North East	Wind direction (Degree)	East North East	East North East	North East	East	East South East
4	3	5	Wind speed Kmp	10	10	12	12	12

Weather Summary for Villupuram district

Weather forecast for next five days: (01.02.2025-05.02.2025)

According to district forecast issued by the India Meteorological Department for Villupuram district, sky will be mostly clear. Maximum temperature is expected to be around 30.0-31.0°C. Minimum temperature is expected to be around 22.0-23.0°C. Morning relative humidity is expected to be around 90-92 per cent and evening relative humidity is expected to be around 68-77 per cent. Wind speed is expected to be around 10-12 km per hour and the wind direction will be from East South East.

SMS Advisory:

(01.02.2025-05.02.2025) Mist/ Haze conditions are very likely to prevail at isolated pockets of Tamil Nadu during morning hours.

- 1. Till the inclement weather the livestock must not be allowed for grazing in early morning hours.
- 2. Drinking water must be warmed to body temperature so as to avoid drip in body temperature.
- 3. The livestock must be provided with vitamins and mineral supplementation to maintain good health during cold weather.
- 4. Antibiotics, oral rehydration solution, intravenous fluids must be stored in handy during emergency
- 5. Pregnant animals nearing parturition should be housed separately in clean dry shelter and must be protected from clod stress by covering the side walls
- 6. The young calf shed must be adequately bedded to prevent clod stress which is very detrimental to the young stock

- 7. Rabbit may be allowed by mating now due to conductive environment rabbit will kindle more bunnies during these months due to which necessary reproductive management must be followed
- 8. Newly born piglets must be protected from cold drafts
- 9. Add 5%shell grit in laying hens for optimum egg production.

Rice:

Due to prevailing weather conditions leaf folder and stem borer infestation found in paddy crop. To manage them, spray carbosulfan 400 ml/ac or cartap hydrochloride 600 g/ac during morning or evening hours.

The forecasted weather may cause stem borer and leaf folder infestation in rice crop. The farmers are advised to monitor the field, if symptoms are observed it's advised to spray Chlorantraniliprole 18.5% 150 ml/ha.

Groundnut:

Due to prevailing weather conditions, there are chances of tobacco caterpillar in groundnut. To control install pheromone trap 5 nos/ac. Spray flubendiamide 20 WG 120 gram/ac or Quinalphos 20 AF @ 500ml/ac.

Blackgram:

Farmers are advised to treat the seeds with Trichoderma viride @ 4g/kg of seed (or) Pseudomonas fluorescens @ 10 g/kg seed before sowing.

Brinjal:

Application of 1.5% foliar spray of PPFM at the flowering stage in Brinjal, will prevent the premature falling of flowers and increase the numbers and quality of fruits.

Banana:

Prevailing weather condition is favorable for sigatoka leaf spot infection in banana. To control spray carbendazim 1.0 g/l or propiconazole 1.0 ml/l or mancozeb 2.5 g/l and sticking agent 1.0 ml/l at 10-15 day intervals.

Prevailing weather condition is favorable for panama wilt and nematode infection in banana. To control apply <u>Bacillus subtilis</u> 1.0 kg/ha + Paecilomyceslilacinus 1.0 kg/ha at 2, 4 and 6th months of planting.



Agromet Advisory Bulletin for the Cuddalore District Bulletin no. 009/2025 Issued on 31.01.2025 (01.02.2025-05.02.2025)

Issued jointly by



Tamil Nadu Veterinary and Animal Sciences University & India Meteorological Department

Weather Summary for the period since the issue of last bulletin for Cuddalore.

During past three days around Cuddalore area, the sky was mostly clear. Maximum temperature was 30.0°C and Minimum temperature was 20.0-22.0°C. Wind speed was 03-05 km per hour. Relative humidity was recorded between 67-95 %.

	t week Sumn 1.2025-30.01.	•	Parameters	Ensemble weather forecast valid until 08.30 hrs of 05.02.2025				
28.01.2025	29.01.2025	30.01.2025	Date	01.02.2025	02.02.2025	03.02.2025	04.02.2025	05.02.2025
0.0	0.0	0.0	Rainfall (mm)	12	10	0	0	10
30	30	30	Max. Temp. (° C)	30	30	30	31	31
20	21	22	Mini. Temp. (° C)	22	22	22	23	23
95	90	91	Maximum RH (%)	90	90	90	92	92
67	68	77	Minimum RH (%)	68	68	68	77	77
Partly cloudy	Partly cloudy	Partly cloudy	Cloud cover (Octa)	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy
East North East	North North East	North East	Wind direction (Degree)	East North East	East North East	North East	East	East South East
4	3	5	Wind speed Kmp	10	10	12	12	12

Weather Summary for Cuddalore district

Weather forecast for next five days: (01.02.2025-05.02.2025)

According to district forecast issued by the India Meteorological Department for Cuddalore district, sky will be mostly clear. Maximum temperature is expected to be around 30.0-31.0°C. Minimum temperature is expected to be around 22.0-23.0°C. Morning relative humidity is expected to be around 90-92 per cent and evening relative humidity is expected to be around 68-77 per cent. Wind speed is expected to be around 10-12 km per hour and the wind direction will be from East South East.

SMS Advisory:

(01.02.2025-05.02.2025) Mist/ Haze conditions are very likely to prevail at isolated pockets of Tamil Nadu during morning hours.

- 1. Till the inclement weather the livestock must not be allowed for grazing in early morning hours.
- 2. Drinking water must be warmed to body temperature so as to avoid drip in body temperature.
- 3. The livestock must be provided with vitamins and mineral supplementation to maintain good health during cold weather.
- 4. Antibiotics, oral rehydration solution, intravenous fluids must be stored in handy during emergency
- 5. Pregnant animals nearing parturition should be housed separately in clean dry shelter and must be protected from clod stress by covering the side walls
- 6. The young calf shed must be adequately bedded to prevent clod stress which is very detrimental to the young stock

- 7. Rabbit may be allowed by mating now due to conductive environment rabbit will kindle more bunnies during these months due to which necessary reproductive management must be followed
- 8. Newly born piglets must be protected from cold drafts
- 9. Add 5%shell grit in laying hens for optimum egg production.

Rice:

Due to prevailing weather conditions leaf folder and stem borer infestation found in paddy crop. To manage them, spray carbosulfan 400 ml/ac or cartap hydrochloride 600 g/ac during morning or evening hours.

The forecasted weather may cause stem borer and leaf folder infestation in rice crop. The farmers are advised to monitor the field, if symptoms are observed it's advised to spray Chlorantraniliprole 18.5% 150 ml/ha.

Groundnut:

Due to prevailing weather conditions, there are chances of tobacco caterpillar in groundnut. To control install pheromone trap 5 nos/ac. Spray flubendiamide 20 WG 120 gram/ac or Quinalphos 20 AF @ 500ml/ac.

Blackgram:

Farmers are advised to treat the seeds with Trichoderma viride @ 4g/kg of seed (or) Pseudomonas fluorescens @ 10 g/kg seed before sowing.

Brinjal:

Application of 1.5% foliar spray of PPFM at the flowering stage in Brinjal, will prevent the premature falling of flowers and increase the numbers and quality of fruits.

Banana:

Prevailing weather condition is favorable for sigatoka leaf spot infection in banana. To control spray carbendazim 1.0 g/l or propiconazole 1.0 ml/l or mancozeb 2.5 g/l and sticking agent 1.0 ml/l at 10-15 day intervals.

Prevailing weather condition is favorable for panama wilt and nematode infection in banana. To control apply <u>Bacillus subtilis</u> 1.0 kg/ha + Paecilomyceslilacinus 1.0 kg/ha at 2, 4 and 6th months of planting.



Agromet Advisory Bulletin for the Kancheepuram District Bulletin no. 009/2025 Issued on 31.01.2025 (01.02.2025-05.02.2025)

Issued jointly by



Tamil Nadu Veterinary and Animal Sciences University & India Meteorological Department

Weather Summary for the period since the issue of last bulletin for Kancheepuram.

During past three days around Kancheepuram area, the sky was mostly clear. Maximum temperature was 30.0-31.0°C and Minimum temperature was 22.0-23.0°C. Wind speed was 05-08 km per hour. Relative humidity was recorded between 69-91 %.

Past week Summary (28.01.2025-30.01.2025)		Parameters	Ensemble weather forecast valid until 08.30 hrs of 05.02.2025				ars of	
28.01.2025	29.01.2025	30.01.2025	Date	01.02.2025	02.02.2025	03.02.2025	04.02.2025	05.02.2025
0.0	0.0	0.0	Rainfall (mm)	0	10	0	0	0
31	31	30	Max. Temp. (° C)	31	31	31	32	32
22	22	23	Mini. Temp. (° C)	23	23	23	24	24
91	88	77	Maximum RH (%)	77	77	77	86	86
69	69	69	Minimum RH (%)	62	62	62	69	69
Partly cloudy	Partly cloudy	Partly cloudy	Cloud cover (Octa)	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy
East	North North East	East	Wind direction (Degree)	East South East	South East	North East	East	East South East
6	8	5	Wind speed Kmp	4	6	12	14	12

Weather Summary for Kancheepuram district

Weather forecast for next five days: (01.02.2025-05.02.2025)

According to district forecast issued by the India Meteorological Department for Kancheepuram district, sky will be mostly clear. Maximum temperature is expected to be around 31.0-32.0°C. Minimum temperature is expected to be around 23.0-24.0°C. Morning relative humidity is expected to be around 77-86 per cent and evening relative humidity is expected to be around 62-69 per cent. Wind speed is expected to be around 04-14 km per hour and the wind direction will be from East South East.

SMS Advisory:

(01.02.2025-05.02.2025) Mist/ Haze conditions are very likely to prevail at isolated pockets of Tamil Nadu during morning hours.

- 1. Till the inclement weather the livestock must not be allowed for grazing in early morning hours.
- 2. Drinking water must be warmed to body temperature so as to avoid drip in body temperature.
- 3. The livestock must be provided with vitamins and mineral supplementation to maintain good health during cold weather.
- 4. Antibiotics, oral rehydration solution, intravenous fluids must be stored in handy during emergency
- 5. Pregnant animals nearing parturition should be housed separately in clean dry shelter and must be protected from clod stress by covering the side walls

- 6. The young calf shed must be adequately bedded to prevent clod stress which is very detrimental to the young stock
- 7. Rabbit may be allowed by mating now due to conductive environment rabbit will kindle more bunnies during these months due to which necessary reproductive management must be followed.
- 8. Newly born piglets must be protected from cold drafts
- 9. Add 5%shell grit in laying hens for optimum egg production.

Rice:

Due to prevailing weather conditions leaf folder and stem borer infestation found in paddy crop. To manage them, spray carbosulfan 400 ml/ac or cartap hydrochloride 600 g/ac during morning or evening hours.

The forecasted weather may cause stem borer and leaf folder infestation in rice crop. The farmers are advised to monitor the field, if symptoms are observed it's advised to spray Chlorantraniliprole 18.5% 150 ml/ha.

Groundnut:

Due to prevailing weather conditions, there are chances of tobacco caterpillar in groundnut. To control install pheromone trap 5 nos/ac. Spray flubendiamide 20 WG 120 gram/ac or Quinalphos 20 AF @ 500ml/ac.

Blackgram:

Farmers are advised to treat the seeds with Trichoderma viride @ 4g/kg of seed (or) Pseudomonas fluorescens @ 10 g/kg seed before sowing.

Brinjal:

Application of 1.5% foliar spray of PPFM at the flowering stage in Brinjal, will prevent the premature falling of flowers and increase the numbers and quality of fruits.

Banana:

Prevailing weather condition is favorable for sigatoka leaf spot infection in banana. To control spray carbendazim 1.0 g/l or propiconazole 1.0 ml/l or mancozeb 2.5 g/l and sticking agent 1.0 ml/l at 10-15 day intervals.

Prevailing weather condition is favorable for panama wilt and nematode infection in banana. To control apply <u>Bacillus subtilis</u> 1.0 kg/ha + Paecilomyceslilacinus 1.0 kg/ha at 2, 4 and 6th months of planting.