

# Precision Production Technology of Marigold

## Introduction:

African marigold (*Tagetes erecta* L.) is an important traditional flower crop under cultivation throughout India. It is extensively used in religious and social functions in different forms. Marigold is widely grown in gardens and pots for display purpose. It has great economic potential in loose flower which find industrial application in preparation of natural dyes and essential oils. It is used as mosquito and nematode repellents. Nowadays Xanthophyll pigments gained economic interest as a feed additive for poultry industry to improve the pigmentation of broiler skin and the egg yolk. Besides, lutein is also used in preparation of eye ailments.

Precision production technology for cultivation of African Marigold for the L3 hybrid was standardized at Tamil Nadu Agricultural University through the ICAR-NAIP project.

## Propagation

Marigold is mainly propagated by seeds.

## Soil and climate

Marigold can be successfully cultivated on a wide variety of soils. Soil that is deep, fertile, friable having good water holding capacity, well drained and with soil pH of 7.5 is ideal for Marigold cultivation. Saline and alkaline soils are not suitable for Marigold cultivation.

It requires continuous warm climate, and extreme hot and cool condition are not good for its growth. It grows well in all seasons. The optimum temperature required for its growth is 15 to 21°C.

## Comparison of precision production system and conventional production system in Marigold.

Precision production system	Conventional production system
<b>Field preparation:</b> <ul style="list-style-type: none"><li>❖ Main field is ploughed using four different implements viz., chisel, disc, cultivator and rotovator.</li></ul>	<b>Field preparation:</b> <ul style="list-style-type: none"><li>❖ Main field is ploughed using three different implements viz., disc, cultivator and rotovator.</li></ul>
<ul style="list-style-type: none"><li>❖ Azospirillum and phospobacteria each @ 2 kg/ha is applied before the last ploughing.</li></ul>	-Nil-

❖ Raised beds (4 feet width x 94 feet) length are formed	❖ Ridges and furrows are formed
<b>Nursery preparation:</b> ❖ The seedlings are raised in protrays ❖ Protrays (54 x 27cm) with 98 cells of 3.5cm diameter and 8mm thickness is ideal	<b>Nursery preparation:</b> ❖ Seeds are sown in raised beds
<b>Seed rate:</b> 200g/ha	<b>Seed rate:</b> 275g/ha
<b>Seedling treatment:</b> Seedlings are dipped in <i>Pseudomonas flourescens</i> @ 0.5 %	<b>Seedling treatment:</b> -Nil-
<b>Transplanting:</b> Seedlings are transplanted within 18 – 20 days	<b>Transplanting:</b> ❖ Seedlings are transplanted after 20 days
<b>Spacing:</b> 90 x 22.5cm	<b>Spacing:</b> ❖ 60 x 45cm
<b>Irrigation:</b> Drip Irrigation (once in 2 days)	<b>Irrigation:</b> ❖ Flood irrigation (once in a week)
<b>Nipping:</b> Carried out 20 days after transplanting	<b>Nipping:</b> ❖ Carried out 20 days after transplanting
<b>Fertilizer application:</b> ❖ 75% of the total recommended Dose of NPK (67.5:67.5:56.25 kg/ha) is applied through drip irrigation	<b>Fertilizer application:</b> ❖ 100% of the total recommended dose of NPK (90:90:75kg/ha) is applied
❖ Water soluble fertilizers viz., Urea Polyfeed(19:19:19) and Potassium nitrate are used	❖ Di Ammonium Phosphate, Murate of Potash are applied through Surface placement method
<b>Micronutrient application:</b> ❖ Foliar application of $FeSO_4$ @ 0.5% and $ZnSO_4$ @ 0.5% at 30 and 45 days after transplanting	<b>Micronutrient application:</b> -Nil-
<b>Biostimulant application:</b> Foliar application of humic acid @ 0.2% and sea weed extract 0.25% at 30 and 45 days after transplanting	<b>Biostimulant application:</b> -Nil-
<b>Integrated pest and disease management:</b> ❖ Application of Spinosad @ 0.75ml/litre for managing the flower borer( <i>Helicoverpa</i>	<b>Integrated pest and disease management:</b> ❖ Indiscriminate use of pesticides

<i>armigera</i> ) ❖ Soil application of <i>Pseudomonas flourescens</i> @ 2.5kg/ha followed by foliar application of <i>Pseudomonas flourescens</i> @ 0.5 % for leaf spot	
<b>Yield:</b> ❖ 32.5 tonnes/ha	<b>Yield:</b> ❖ 25 tonnes/ha
<b>Xanthophyll content:</b> ❖ 1.99g/kg of flowers	<b>Xanthophyll content:</b> ❖ 1.42 g/kg of flowers

Source:

Department of Floriculture and Landscaping  
Tamil Nadu Agricultural University  
Coimbatore