



**Pandit Jawaharlal Nehru College of Agriculture and
Research Institute, Karaikal - 609 603**

Agromet Advisory Bulletin (AAB) for KARAİKAL District
Issued jointly by IMD & PAJANCOA & RI

Forecast Period: 10th to 14th August 2024

Phone No. 04368 – 261182 E-mail: agronaas2012@gmail.com

Fax: 04368-261260



Bulletin No. 064 (08)/PJN/GKMS/2024

Date: 09.08.2024

Weather Summary for Karaikal district

Past week Summary (04.08.2024 to 08.08.2024)					Weather Parameters	Ensemble weather forecast valid until 08.30 hrs of 14.08.2024				
04/08	05/08	06/08	07/08	08/08		Day-1 10/08	Day-2 11/08	Day-3 12/08	Day-4 13/08	Day-5 14/08
15.0	11.0	5	11.2	60.4	Rainfall (mm)	6	8	6	12	5
38	37	37	37	36	Max. Temp. (°C)	34	34	33	32	30
25	24	25	24	24	Mini. Temp. (°C)	24	23	23	22	21
4	5	4	4	4	Cloud cover (Octa)	6	8	8	8	6
44-84	69-92	55-92	60-89	61-90	RH Even- Morn. (%)	50-70	50-80	50-80	50-80	50-80
5	4	2	3	4	Wind (kmph)	18	18	14	16	22
SW	NNE	WNW	SW	WSW	Wind direction	SSW	SSW	S	SSW	S

Forecast for next five days: 10.08.2024 to 14.08.2024

Light rain is expected from 10th to 14th August.

During next five days the sky will be cloudy. The temperature range is expected to be around 21-34°C. Relative humidity range is expected to be around 50-80%. The wind speed is expected to be 14-22 km per hour and the wind will be mostly from south south west direction.

Rainfall (6.1 mm) will be above normal, maximum temperature (32.6°C) and minimum temperature (24.1°C) will be normal from 14th to 20th August.

Agro Advisory

Crop/livestock	Stage/Particulars	Agro Advisory
Cotton	Flowering to picking stage	To manage mealy bug in cotton, spray Thiamethoxam 25 WG @ 0.2 g/l or Buprofezin 25 SC @ 2 ml/l.
		To manage white fly in cotton, spray Azadirachtin 0.03 (300 ppm) @ 5-10 ml/l or Thiamethoxam 25 WG 0.4 g/l.
Brinjal	Bearing stage	To manage tobacco cutworm in brinjal, spray Bacillus thuringiensis 2 g/l or Novaluron 10% EC 1 ml/l or flubendamide 0.2 – 0.3 ml/l
Cattle	Babesiasis	The prevailing weather conditions are favorable for the Babesiasis disease in cattle. High fever, dark coloured

		urine, less intake of feed and less milk yield are the symptoms of Babesiasis disease. If these symptoms appear in the cattle, contact nearby dispensary. As a preventive measure, apply Butox @ 2 ml/l of water on cattle and also spray it on the nearby areas of the cattle shed to control ectoparasites
--	--	--

SMS: Light rain is expected for next five days. Hence, postpone irrigation to crops.

**DEAN
PAJANCOA & RI**

