



Agromet Advisory Bulletin for **Thanjavur** District

(From 03rd August 2024 to 07th August 2024)

Issued jointly by Tamil Nadu Rice Research Institute -
Aduthurai and IMD



Bulletin no: 36/2024

Date: 02.08.2024

Weather Summary for the period since the issue of last bulletin no. 35 (30.07.2024) for Tamil Nadu: Thanjavur district

Maximum temperature of 35 - 39 °C, minimum temperature of 29 - 30 °C, and was recorded in Thanjavur district.

Weather forecast for next five days (03.08.2024 to 07.08.2024)

Past week Summary 30.07.2024 to 01.08.2024			Parameters	Ensemble weather forecast valid until 08.30 hr of 07.08.2024				
Day-1 30/07	Day-2 31/07	Day-3 01/08	Date	Day-1 03/08	Day-2 04/08	Day-3 05/08	Day-4 06/08	Day-5 07/08
0	0	0	Rainfall (mm)	1	1	2	2	5
35	38	39	Max. Temp. (°C)	39	39	38	38	38
29	30	29	Min. Temp. (°C)	30	30	29	29	29
5	5	5	Cloud cover (Octa)	5	6	6	7	8
69-51	70-52	72-57	RH I-II (%)	60-40	60-40	60-40	70-40	70-40
5	6	6	Wind (kmph)	12	12	12	12	12
West South West	West South West	West South West	Wind direction	West South West	West	West	East South East	South South East

Weather Summary and forecast for Thanjavur district

According to district level forecast issued by India Meteorological Department for Thanjavur district, the sky will be clear. Light drizzling expected on 03.08.2024 to 07.08.2024. Maximum temperature is expected to be around 38 - 39 °C and minimum temperature is expected to be around 29 - 30 °C. Morning relative humidity is expected to be around 60 - 70 percent and evening relative humidity is expected to be around 40 percent. Average wind speed is expected to be around 12 km per hour. The wind direction will be west.

Agro Advisory

Crop	Stage	Advisory
General		Farmers are requested to download and utilize the agricultural based mobile apps suggested by TNAU (TNAU AAS, Meghdoot and DAMINI).
Rice	Main field	The prevailing weather condition favorable for gall midge infection in rice. It leads to the formation of onion like leaf and prevents the panicle formation. To control this apply chloripyriphos 20 EC 500 ml/ ha.
Rice	Main field	At active tillering stage of Kuruvai paddy, top dressing can be done with 35 kg urea and 10 kg MOP/acre by keeping thin film of water in the field.
Banana		Provide support with wooden poles to avoid lodging of tree due to wind.
Cattle	Cow	Farmers are advised to apply Butox solution (2 ml in 1 litre water) externally to remove ecto - parasites in sheep and goats.

SMS advisory

Owing to moderate wind, more moisture loss will be expected and hence, more irrigation should be given.

**Director
TRRI, Aduthurai**