



## Agromet Advisory Bulletin for the **Nagaipattinam** District

(From 19<sup>th</sup> June 2024 to 23<sup>rd</sup> June 2024)

Issued jointly by Tamil Nadu Rice Research Institute -  
Aduthurai and IMD



**Bulletin no: 23/2024**

**Date: 18.06.2024**

Weather Summary for the period since the issue of last bulletin no. 22 (14.06.2024) for Tamil Nadu: Nagaipattinam district

Maximum temperature of 35 - 38 °C, minimum temperature of 26 °C, and was recorded in Nagaipattinam district.

**Weather forecast for next five days (19.06.2024 to 23.06.2024)**

Past week Summary 14.06.2024 to 17.06.2024				Parameters	Ensemble weather forecast valid until 08.30 hr of 23.06.2024				
Day-1 14/06	Day-2 15/06	Day-3 16/06	Day-4 17/06		Date	Day-1 19/06	Day-2 20/06	Day-3 21/06	Day-4 22/06
0	0	0	0	<b>Rainfall (mm)</b>	2	3	3	3	2
35	37	38	38	<b>Max. Temp. (°C)</b>	36	36	35	35	35
26	26	26	26	<b>Min. Temp. (°C)</b>	27	27	26	26	26
7	6	6	3	<b>Cloud cover (Octa)</b>	8	7	7	6	6
77-50	78-67	61-70	80-64	<b>RH I-II (%)</b>	70-50	70-50	70-50	60-50	70-50
5	4	7	6	<b>Wind (kmph)</b>	24	22	22	20	18
West	South West	South	South	<b>Wind direction</b>	South South West	South South West	South South West	South South West	South West

### **Weather Summary and forecast for Nagaipattinam district**

According to district level forecast issued by India Meteorological Department for Nagaipattinam district, the sky will be clear. Light drizzling expected on 19.06.2024 to 23.06.2024. Maximum temperature is expected to be around 35 - 36 °C and minimum temperature is expected to be around 26 - 27 °C. Morning relative humidity is expected to be around 50 percent and evening relative humidity is expected to be around 50 percent. Average wind speed is expected to be around 18 - 24 km per hour. The wind direction will be south south west.

### **Agro Advisory**

Crop	Stage	Advisory
General		Farmers are requested to download and utilize the agricultural based mobile apps suggested by TNAU (TNAU AAS, Meghdoot and DAMINI).

Rice	Nursery	Roots of 18 to 22 days of old young seedlings need to be dipped in biofertilizers slurry (1 kg or 125 ml of each <i>Azospirillum</i> and phosphobacteria ) for 15 to 30 minutes before transplanting.
Cotton	Vegetative	Prevailing weather condition favours shedding of buds and squares in cotton. To prevent this spray 40 ppm NAA at 60 and 90 days after sowing.
Sugarcane	Vegetative	Considering weather conditions, detrashing and propping of five months old sugarcane is recommended.
Cattle	Cow	Farmers are advised provide sufficient mineral mixture to nourish the cattle health.

### **SMS advisory**

Farmers are requested to download and utilize the agricultural based mobile apps suggested by TNAU (TNAU AAS, Meghdoot and DAMINI).

**Director  
TRRI, Aduthurai**