



राज्य समग्र कृषि-मौसम परामर्श बुलेटिन (तमिलनाडु, पुडुचेरी और कारईकाल के लिए) State Composite Agromet Advisory Bulletin (for Tamil Nadu, Puducherry, and Karaikal)

Bulletin No: 036/2026, Issued on 05.05.2026

(अवधि: 06 मई से 10 मई 2026, For the period: 06th May to 10th May 2026)



संकलन एवं प्रकाशन / Compiled and issued by:

कृषि-मौसम परामर्श सेवा इकाई (AASU)

भारत मौसम विज्ञान विभाग / India Meteorological Department

प्रादेशिक मौसम विज्ञान केन्द्र चेन्नई / Regional Meteorological Centre (RMC), Chennai

Website: <https://mausam.imd.gov.in/chennai/>

Email: rmcmdsaasu@gmail.com; Contact: +91-44-28246027

**नोट: यह बुलेटिन तमिलनाडु, पुडुचेरी और कारईकाल के अंतर्गत सभी कृषि-मौसम प्रक्षेत्र इकाई द्वारा तैयार की गई कृषि-मौसम परामर्श बुलेटिन का संकलन है।
Note: This bulletin consolidates Agromet advisories prepared by all AMFUs under Tamil Nadu, Puducherry & Karaikal.**

Weather warnings (Met-Subdivision Level)

HEAVY RAINFALL WARNING	
Day 1 (05.05.2026)	Heavy rain is likely to occur at isolated places over Theni, Dindigul, Karur, Tiruchirappalli, Namakkal, Erode and Nilgiris districts of Tamilnadu and Kerala.
Day 2 (06.05.2026)	Heavy rain is likely to occur at isolated places over Ghat areas of Coimbatore district, Nilgiris, Tiruppur, Theni, Tenkasi, Tirunelveli and Kanyakumari districts of Tamilnadu and Kerala.
Day 3 (07.05.2026)	Heavy rain is likely to occur at isolated places over Ghat areas of Coimbatore district, Nilgiris, Tiruppur, Theni, Tenkasi, Erode, Madurai, Virudhunagar, Tirunelveli and Kanyakumari districts of Tamil Nadu, Kerala and South Interior Karnataka.
Day 4 (08.05.2026)	Heavy rain is likely to occur at isolated places over Nilgiris, Coimbatore, Erode, Tiruppur, Dindigul, Theni, Madurai, Sivaganga, Virudhunagar, Tenkasi, Ramanathapuram, Thoothukudi, Tirunelveli, Kanyakumari, Thanjavur, Tiruvarur, Nagapattinam, Mayiladuthurai, Pudukottai districts of Tamilnadu, Karaikal area, Kerala and South Interior Karnataka.
Day 5 (09.05.2026)	Heavy rain is likely to occur at isolated places over Nilgiris, Coimbatore, Erode, Tiruppur, Dindigul, Theni, Madurai, Sivaganga, Virudhunagar, Tenkasi, Ramanathapuram, Thoothukudi, Tirunelveli, Kanyakumari, Thanjavur, Tiruvarur, Nagapattinam, Mayiladuthurai, Pudukottai districts of Tamilnadu, Karaikal area, Kerala and South Interior Karnataka.
Day 6 (10.05.2026)	Heavy rain is likely to occur at isolated places over Nilgiris, Coimbatore, Erode, Tiruppur, Dindigul, Theni, Madurai, Sivaganga, Virudhunagar, Tenkasi, Ramanathapuram, Thoothukudi, Tirunelveli & Kanyakumari districts of Tamilnadu and Kerala.
Day 7 (11.05.2026)	Heavy rain is likely to occur at isolated places over Kerala.

THUNDERSTORM WARNING

Day 1 (06.05.2026) & Day 2 (07.05.2026)	Thunderstorm & lightning with gusty winds speed reaching 40-50 kmph is likely to occur at one or two places over Tamilnadu, Puducherry & Karaikal area, Kerala, South Interior Karnataka, Coastal Andhra Pradesh and Rayalaseema. Thunderstorm & lightning with gusty winds speed reaching 30-40 kmph is likely to occur at one or two places over North Interior Karnataka and Telangana. Thunderstorm & lightning is likely to occur at one or two places over Coastal Karnataka.
Day 3 (08.05.2026) & Day 4 (09.05.2026)	Thunderstorm & lightning with gusty winds speed reaching 40-50 kmph is likely to occur at one or two places over Kerala, South Interior Karnataka, Coastal Andhra Pradesh and Rayalaseema. Thunderstorm & lightning with gusty winds speed reaching 30-40 kmph is likely to occur at one or two places over Tamilnadu, Puducherry & Karaikal area, North Interior Karnataka and Telangana. Thunderstorm & lightning is likely to occur at one or two places over Coastal Karnataka.
Day 5 (10.05.2026)	Thunderstorm & lightning with gusty winds speed reaching 40-50 kmph is likely to occur at one or two places over Tamilnadu, Puducherry & Karaikal area, Kerala, South Interior Karnataka, Coastal Andhra Pradesh and Rayalaseema. Thunderstorm & lightning with gusty winds speed reaching 30-40 kmph is likely to occur at one or two places over North Interior Karnataka and Telangana. Thunderstorm & lightning is likely to occur at one or two places over Coastal Karnataka.

TEMPERATURE WARNING

Day 1 (06.05.2026)	Due to high maximum temperature, combined with high humidity is likely to cause discomfort over Coastal Andhra Pradesh and Rayalaseema.
Day 2 (07.05.2026)	Due to high maximum temperature, combined with high humidity is likely to cause discomfort over Coastal Andhra Pradesh.

Extended Range forecast at Met-Subdivision Level (06th May to 12th May 2026)

Meteorological Sub-Division	Rainfall	Max Temp	Min Temp
Tamilnadu, PDC & KKL	Above Normal	Below Normal	Normal
Kerala & Mahe	Above Normal	Below Normal	Normal
Lakshadweep	Above Normal	Below Normal	Normal
Coastal Karnataka	Above Normal	Below Normal	Above Normal
North Interior Karnataka	Above Normal	Below Normal	Above Normal
South Interior Karnataka	Above Normal	Below Normal	Above Normal
Coastal Andhra Pradesh & Yanam	Below Normal	Below Normal	Normal
Rayalaseema	Above Normal	Below Normal	Normal
Telangana	Below Normal	Below Normal	Normal

Summary of past five days rainfall over the State (30th April to 04th April 2026)

Met Subdivision	Day 1	Day 2	Day 3	Day 4	Day 5
	(30.04.2026)	(01.05.2026)	(02.05.2026)	(03.05.2026)	(04.05.2026)
Tamilnadu, PDC & KKL	ISOL	ISOL	ISOL	SCT	ISOL
Kerala & Mahe	SCT	FWS	ISOL	SCT	ISOL
Lakshadweep	SCT	DRY	DRY	DRY	DRY
Coastal Karnataka	ISOL	ISOL	DRY	DRY	DRY
North Interior Karnataka	ISOL	ISOL	DRY	DRY	ISOL
South Interior Karnataka	SCT	SCT	ISOL	ISOL	ISOL
Coastal Andhra Pradesh & Yanam	ISOL	SCT	SCT	ISOL	SCT
Rayalaseema	ISOL	ISOL	ISOL	DRY	ISOL
Telangana	ISOL	ISOL	ISOL	ISOL	ISOL

ISOL: Isolated *i.e.* Rain at 1 or 2 places

FWS: Fairly Widespread *i.e.* Rain at many places

WS: Widespread *i.e.* Rain at most places

SCT: Scattered *i.e.* Rain at a few places

DRY: No rain

Rainfall forecast for next 5 days

Met Subdivision	Day 1	Day 2	Day 3	Day 4	Day 5
	(06.05.2026)	(07.05.2026)	(08.05.2026)	(09.05.2026)	(10.05.2026)
Tamilnadu, PDC & KKL	ISOL	SCT	SCT	SCT	SCT
Kerala & Mahe	FWS	FWS	FWS	FWS	WS
Lakshadweep	SCT	SCT	SCT	SCT	FWS
Coastal Karnataka	ISOL	ISOL	ISOL	ISOL	ISOL
North Interior Karnataka	ISOL	ISOL	ISOL	ISOL	ISOL
South Interior Karnataka	ISOL	ISOL	SCT	SCT	SCT
Coastal Andhra Pradesh & Yanam	SCT	SCT	SCT	SCT	ISOL
Rayalaseema	ISOL	ISOL	SCT	SCT	ISOL
Telangana	ISOL	SCT	ISOL	ISOL	ISOL

ISOL: Isolated *i.e.* Rain at 1 or 2 places

SCT: Scattered *i.e.* Rain at a few places

FWS: Fairly Widespread *i.e.* Rain at many places

**Agromet Advisories issued by different Agromet Field Units
(AMFUs) of Tamil Nadu and Puducherry**

**CHENNAI, KANCHEEPURAM, VILLUPURAM, TIRUVANNAMALAI,
TIRUVALLUR, VELLORE, CUDDALORE, CHENGALPATTU,
TIRUPATTUR AND RANIPET**

SMS Advisory:

(06.05.2026-10.05.2026) Light to moderate rain at a few places with thunderstorm & lightning with gusty winds speed reaching 40-50 kmph at one or two places is likely to occur over Tamilnadu.

Livestock Advisory:

1. The farmers should avoid allowing the animals to graze in hot parts of the day.
2. Sheep and goat should be dewormed by using suitable anthelmintic by consulting with Local veterinary doctor.
3. Increasing the frequency of feeding from three to four times per day to high yielding crossbreed, dairy cattle would nominally increase milk production and at the same time will reduce the heat stress.
4. The dairy buffaloes are very much prone to summer sub fertility, due to higher ambient air temperature. To prevent this proper sheltering with cooling facility are necessary.
5. White pigs meant for fattening purpose which are fed with high fat containing swill feed would suffer heat load during digestion, during summer season. Hence to prevent this fat containing will feed should be fed during late evening hours.
6. Rabbitaries with asbestos/metal roofs will experience more heat during summer, hence covering the roof with coconut leaves or gunny bags and splashing water over them will reduce the inner micro climate of rabbit shed.
7. Provision of cool wholesome drinking water to livestock and poultry during summer is essential.

Agricultural advisory:

Rice:

Due to prevailing weather condition chances of thrips infestation in paddy field. To control spray thiamethoxam 25% WG 100 g/ha or dimethoate 1.5ml/l.

Due to prevailing weather condition leaf folder and stem borer infestation is found in paddy crop. To manage them, spray carbosulfan 400 ml/ac or cartap hydrochloride 600 g/ac during morning or evening hours.

Harvest the matured paddy crops and store in safe places.

For nursery land preparation, to prevent soil borne diseases and better establishment of seedlings apply Azospirillum 2.5 Kg mixed with 25 kg of soil.

Sugarcane:

Mealy bug and pokkahboeng infection identified in sugarcane crop. Mealy bug affect growth of plants and pokkahboeng cause death of young spindles of sugarcane crop. To control spray imidacloprid 17.8 SL 0.6 ml/l or carbendazim 1.0 g/l.

A tonic with nutrients and growth regulators for sugarcane spray the TNAU sugarcane Booster at 45, 60 and 75 days after planting spray booster at 1kg 1.5kg and 2kg respectively and add required quantity of sticking agent.

Coconut:

Installation of yellow sticky traps on the palm trunk to trap the adult whiteflies.

Encourage buildup of parasitoids (*Encarsia* sp.) and reintroduce parasitized pupae of emerging zones of whitefly outbreak.

In sever case, spray neem oil 10 ml/lit of water or spray azadirachtin 2.0 ml/lit of water with 1.0 ml of sticking agent.

Apply coconut husks with convex surface facing upwards (100 Nos) or dried coconut leaves (15 Nos) or coir pith upto a height of 10 cm in the basin of 1.8 meter radius around the palms as much for soil moisture conservation particularly during summer season.

Banana:

To control wilt in banana, apply *Bacillus subtilis* and *Trichoderma viride* @ 2.5 kg/ha. Or drenching with *Bacillus subtilis* 5 ml/litre. To overcome micro nutrient deficiency and to improve quality of banana fruit micronutrient plays a major role. Micro nutrient can be sprayed at 3,5 and 7 months after planting of banana.

COIMBATORE

Weather Warning on Agriculture

Heavy rain, Thunderstorms & lightning, squally wind is expected in some places and hence farmers are requested to stake banana crop which are more than five month old.

Agro Advisory

Crop	Stage	Advisory
General		Moderate to Heavy rainfall expected in Coimbatore, Erode and Tiruppur this week. Wind speed of 4 -18 kmph is expected in Western Zone district Rainfall may be accompanied by thunderstorm, lightning and squally wind speed. Hence suitable precautionary measures may be followed to protect the crops and animals. By utilizing expected rainfall, summer ploughing is suggested across the slope to utilize the rain effectively.
Sugarcane	Vegetative	Since rainfall is expected post pone the irrigation or based soil moisture condition irrigation may be done. If crown mealy bug noticed, spray imidacloprid 17.8 SL @ 30ml / acre
Vegetables	Vegetative	Prevailing weather is favourable for the mite attack in vegetables. Hence, monitor the crop and if needed spray propargite 57% EC @ 400 – 600 ml / acre.
Banana		Since rainfall along with thunder, lightning and heavy rain with squally wind is expected. Hence provide stacking especially for the more than 7 months old banana. Apply <i>Trichoderma viride</i> @ 2.5 kg/ha + 300 kg of FYM to manage wilt disease.
Coconut		Since heavy rain expected an Inward basin may be formed around the coconut trees to conserve the summer rains in the root zone. Due to high temperature Rugose spiraling whitefly incidence is increasing. Root Wilt (<i>Candidatus Phytoplasma</i>) Recommended NPK + FYM-50kg + Neem Cake 5 kg / palm/yr

		<p>+ Green Manuring Azospirillum – 200g + Phosphobacteria - 200g + VAM – 100g + Bacillus subtilis – 200g + Trichoderma asperellum 200 g / palm / yr TNAU Coconut tonic root feeding @ 40 ml / palm at 6 months interval Soil application of ‘TNAU COCOCON’ @ 5 litre / acre at 3 months interval Need based Root feeding of hexaconazole (3ml + 200 ml water) for leaf Rot</p>
--	--	---

Animal Husbandry		<p>Animals should not be send for grazing between 10 am to 4 pm, considering high temperature and low relative humidity. Make the cattle at cool and provide with drinking water @ 80-100 liters per day per animal in four splits Prevailing high temperature & low humidity affects the feed in take reduces milk yield in dairy cow. Increase dietary minerals in food supplements to counteract the harsh environment. In some places Thullumari disease is observed in sheep and goats hence farmers are requested to allow the sheep and goats only during sunshine hours further they are requested to vaccinate in nearby veterinary hospital. Prevailing high temperature leads to temperature stress in poultry and hence wetted gunny bags may be hanged in the side walls of the shed.</p>
------------------	--	---

SMS advisory:

Provide proper ventilation in sheds to ensure proper airflow to avoid increased humidity following showers, which can cause heat stress.

ERODE

Weather Warning on Agriculture

Heavy rain, Thunderstorms & lightning, squally wind is expected in some places and hence farmers are requested to stake banana crop which are more than five month old.

Agro Advisory

Crop	Stage	Advisory
General		<p>Moderate to Heavy rainfall expected in Coimbatore, Erode and Tiruppur this week. Wind speed of 4 -18 kmph is expected in Western Zone district Rainfall may be accompanied by thunderstorm, lightning and squally wind speed. Hence suitable precautionary measures may be followed to protect the crops and animals. By utilizing expected rainfall, summer ploughing is suggested across the slope to utilize the rain effectively.</p>
Sugarcane	Vegetative	<p>Since rainfall is expected post pone the irrigation or based soil moisture condition irrigation may be done. If crown mealy bug noticed, spray imidacloprid 17.8 SL @ 30ml / acre</p>

Vegetables	Vegetative	Prevailing weather is favourable for the mite attack in vegetables. Hence, monitor the crop and if needed spray propargite 57% EC @ 400 – 600 ml / acre.
Banana		Since rainfall along with thunder, lightning and heavy rain with squally wind is expected. Hence provide stacking especially for the more than 7 months old banana. Apply Trichoderma viride @ 2.5 kg/ha + 300 kg of FYM to manage wilt disease.
Coconut		Since heavy rain expected an Inward basin may be formed around the coconut trees to conserve the summer rains in the root zone. Due to high temperature Rugose spiraling whitefly incidence is increasing. Root Wilt (Candidatus Phytoplasma) Recommended NPK + FYM-50kg + Neem Cake 5 kg / palm/yr + Green Manuring Azospirillum – 200g + Phosphobacteria - 200g + VAM – 100g + Bacillus subtilis – 200g + Trichoderma asperellum 200 g / palm / yr TNAU Coconut tonic root feeding @ 40 ml / palm at 6 months interval Soil application of 'TNAU COCOCON' @ 5 litre / acre at 3 months interval Need based Root feeding of hexaconazole (3ml + 200 ml water) for leaf Rot
Animal Husbandry		Animals should not be send for grazing between 10 am to 4 pm, considering high temperature and low relative humidity. Make the cattle at cool and provide with drinking water @ 80-100 liters per day per animal in four splits Prevailing high temperature & low humidity affects the feed in take reduces milk yield in dairy cow. Increase dietary minerals in food supplements to counteract the harsh environment. In some places Thullumari disease is observed in sheep and goats hence farmers are requested to allow the sheep and goats only during sunshine hours further they are requested to vaccinate in nearby veterinary hospital. Prevailing high temperature leads to temperature stress in poultry and hence wetted gunny bags may be hanged in the side walls of the shed.

SMS advisory:

Provide proper ventilation in sheds to ensure proper airflow to avoid increased humidity following showers, which can cause heat stress.

TIRUPPUR

Weather Warning on Agriculture

Heavy rain, Thunderstorms & lightning, squally wind is expected in some places and hence farmers are requested to stake banana crop which are more than five month old.

Agro Advisory

Crop	Stage	Advisory
General		Moderate to Heavy rainfall expected in Coimbatore, Erode and Tiruppur this week. Wind speed of 4 -18 kmph is expected in Western Zone district Rainfall may be accompanied by thunderstorm, lightning and squally wind speed. Hence suitable precautionary measures may be followed to protect the crops and animals. By utilizing expected rainfall, summer ploughing is suggested across the slope to utilize the rain effectively.
Sugarcane	Vegetative	Since rainfall is expected post pone the irrigation or based soil moisture condition irrigation may be done. If crown mealy bug noticed, spray imidacloprid 17.8 SL @ 30ml / acre
Vegetables	Vegetative	Prevailing weather is favourable for the mite attack in vegetables. Hence, monitor the crop and if needed spray propargite 57% EC @ 400 – 600 ml / acre.
Banana		Since rainfall along with thunder, lightning and heavy rain with squally wind is expected. Hence provide stacking especially for the more than 7 months old banana. Apply Trichoderma viride @ 2.5 kg/ha + 300 kg of FYM to manage wilt disease.
Coconut		Since heavy rain expected an Inward basin may be formed around the coconut trees to conserve the summer rains in the root zone. Due to high temperature Rugose spiraling whitefly incidence is increasing. Root Wilt (Candidatus Phytoplasma) Recommended NPK + FYM-50kg + Neem Cake 5 kg / palm/yr + Green Manuring Azospirillum – 200g + Phosphobacteria - 200g + VAM – 100g + Bacillus subtilis – 200g + Trichoderma asperellum 200 g / palm / yr TNAU Coconut tonic root feeding @ 40 ml / palm at 6 months interval Soil application of 'TNAU COCOCON' @ 5 litre / acre at 3 months interval Need based Root feeding of hexaconazole (3ml + 200 ml water) for leaf Rot
Animal Husbandry		Animals should not be send for grazing between 10 am to 4 pm, considering high temperature and low relative humidity. Make the cattle at cool and provide with drinking water @ 80-100 liters per day per animal in four splits Prevailing high temperature & low humidity affects the feed in

		<p>take reduces milk yield in dairy cow. Increase dietary minerals in food supplements to counteract the harsh environment.</p> <p>In some places Thullumari disease is observed in sheep and goats hence farmers are requested to allow the sheep and goats only during sunshine hours further they are requested to vaccinate in nearby veterinary hospital.</p> <p>Prevailing high temperature leads to temperature stress in poultry and hence wetted gunny bags may be hanged in the side walls of the shed.</p>
--	--	---

SMS advisory:

Provide proper ventilation in sheds to ensure proper airflow to avoid increased humidity following showers, which can cause heat stress.

**ARIYALUR, KARUR, NAGAPATTINAM, PERAMBALUR,
THANJAVUR, TIRUVARUR AND TIRUCHIRAPALLI,
MAYILADUTHURAI**

Agro Advisory

General		Postpone/avoid irrigation to all crops depending upon rain condition.
Rice	Nursery	For summer nursery, an area of 20 cents can be allotted for planting 1 ha with the application of FYM or compost @ 1 ton. For clay soil, 4 kg gypsum and 1 kg DAP/cent can be applied 10 days prior to pulling out of seedlings to avoid root snapping.
Rice	Main field	Application of 10 kg zinc sulphate and 200 kg gypsum per acre is recommended at the time of transplanting, in addition to the recommended dose of nitrogen, Phosphorus and Potassium.
Banana		Provide support with wooden poles to avoid lodging of tree due to wind.
Cattle	Cow and Goat	Calves above 3 months and sheep and goat above 2 months should be vaccinated against Foot and Mouth disease

		When the air temperature is above 35 degrees Celsius the dairy cattle tend to drink more water thereby decreasing the Dry Matter Intake (DMI). Hence to overcome this succulent protein rich green fodder mixed with green fodder may be provided to livestock.
--	--	---

SMS advisory: Postpone/avoid irrigation to all crops depending upon rain condition.

RAMANATHAPURAM

SMS Advisory	Chances of Moderate rain on 9 & 10 May. Maximum temperature would be in the range of 31°C to 35°C and the minimum temperature would be between 26°C-29°C. Cloudy sky will appear for the next five days. Wind is expected with the speed of 12-14 Kmph from, Southeast direction.
--------------	---

General Advisory:

Heavy rain is likely to occur at isolated places over the district on 9 & 10 May 2026. Farmers are advised to act swiftly to repair the water channels, farm ponds and other water bodies to harvest the rain water. During these days farmers are advised to avoid irrigation, intercultural operations and application of plant protection measures to the standing crops, provide adequate drainage facilities in the fields to avoid water stagnation due to heavy rainfall, do not allow the animals to grazing outside, standing near the trees or electric poles & provide safe shelter.

Wind Warning:

Banana, Papaya, Moringa & Sugarcane

Thunderstorm with Lightning with Strong Gusty Wind with a speed of 40 to 50 kmph is expected during 8, 9 & 10 May 2026.

Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya and Moringa, to prevent from lodging.

Gourds:

For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support.

Weather Based Agro-Advisories: (Maize, Horticulture Crops, Coconut, Horticulture crop & Animal Husbandry)

Maize:

Postpone the sowing / Planting Activities.

Drain the excess rain water from the field.

Postpone Plant protection activities such as Fertilizer application and Pesticides spraying for next 5 days.

Horticulture Crops:

Avoid irrigating the fields.

Drain the excess rain water from the field.

Postpone Plant protection activities such as Fertilizer application and Pesticides spraying for next 5 days

Coconut:

Heavy rainfall is expected in the coming days, coconut growers are advised to safely remove the old and decayed fronds. Provide adequate drainage facilities in the field.

Horticulture crop:

Since heavy rainfall expected, farmers are advised to provide adequate drainage facilities in the fields to avoid water stagnation and farmers are advised to repair the water channels.

Avoid spraying of fertilizer, fungicide and pesticide during the rainy days.

Weather based Agro Advisory:

Heavy rainfall is expected on 9 & 10 May 2026. During these days farmers are advised to avoid irrigation, inter-cultural operations and application of plant protection measures to the standing crops.

Provide adequate drainage facilities in the field and renovate the rain water harvesting structures to avoid water stagnation due to heavy rainfall. Try to harvest the rainwater in the farm ponds / water bodies.

Provide support to crops likes Banana, Papaya and sugarcane.

Animal Husbandry:

To protect the livestock from thunderstorms and lightning Farmers are requested to avoid grazing the animals during rains, particularly grazing sheep and goat in the flock which are more prone to lightning shock.

Farmers should avoid tethering animals near barbed wire fencing posts and trees. Trees near the animal sheds should be pruned to avoid accidental falls over during heavy wind.

Further trees like rain trees and vathanarayanan are more prone to uprooting. More chance of lightning near the hilly area and undulated landscapes than in plains.

Fence with wooden poles and metal wires, it is advisable to check the condition of the buried parts at most once in 5 years and do the necessary replacement before the thunderstorm season begins.

MADURAI

SMS Advisory	Chances of Moderate rain during 8, 9 & 10 May. Maximum temperature would be in the range of 33°C to 37°C and the minimum temperature would be between 23°C-27°C. Cloudy sky will appear for the next five days. Wind is expected with the speed of 8-12 Kmph from, Southeast & South directions.
--------------	--

General Advisory:

Heavy rain is likely to occur at isolated places over the district during 8, 9 & 10 May 2026. Farmers are advised to act swiftly to repair the water channels, farm ponds and other water bodies to harvest the rain water. During these days farmers are advised to avoid irrigation, intercultural operations and application of plant protection measures to the standing crops, provide adequate drainage facilities in the fields to avoid water stagnation due to heavy rainfall, do not allow the animals to grazing outside, standing near the trees or electric poles & provide safe shelter.

Wind Warning:

Banana, Papaya, Moringa & Sugarcane

Thunderstorm with Lightning with Strong Gusty Wind with a speed of 40 to 50 kmph is expected during 8, 9 & 10 May 2026.

Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya and Moringa, to prevent from lodging.

Gourds:

For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support.

Weather Based Agro-Advisories: (Maize, Horticulture Crops, Coconut, Horticulture crop & Animal Husbandry)

Maize:

Postpone the sowing / Planting Activities.

Drain the excess rain water from the field.

Postpone Plant protection activities such as Fertilizer application and Pesticides spraying for next 5 days.

Horticulture Crops:

Avoid irrigating the fields.

Drain the excess rain water from the field.

Postpone Plant protection activities such as Fertilizer application and Pesticides spraying for next 5 days.

Coconut:

Heavy rainfall is expected in the coming days, coconut growers are advised to safely remove the old and decayed fronds. Provide adequate drainage facilities in the field.

Horticulture crop:

Since heavy rainfall expected, farmers are advised to provide adequate drainage facilities in the fields to avoid water stagnation and farmers are advised to repair the water channels.

Avoid spraying of fertilizer, fungicide and pesticide during the rainy days.

Weather based Agro Advisory:

Heavy rainfall is expected during 8, 9 & 10 May 2026. During these days farmers are advised to avoid irrigation, intercultural operations and application of plant protection measures to the standing crops.

Provide adequate drainage facilities in the field and renovate the rain water harvesting structures to avoid water stagnation due to heavy rainfall. Try to harvest the rainwater in the farm ponds / water bodies.

Provide support to crops likes Banana, Papaya and sugarcane.

Animal Husbandry:

To protect the livestock from thunderstorms and lightning Farmers are requested to avoid grazing the animals during rains, particularly grazing sheep and goat in the flock which are more prone to lightning shock.

Farmers should avoid tethering animals near barbed wire fencing posts and trees. Trees near the animal sheds should be pruned to avoid accidental falls over during heavy wind.

Further trees like rain trees and vathanarayanan are more prone to uprooting. More chance of lightning near the hilly area and undulated landscapes than in plains.

Fence with wooden poles and metal wires, it is advisable to check the condition of the buried parts at most once in 5 years and do the necessary replacement before the thunderstorm season begins.

PUDUKKOTTAI

SMS Advisory	Chances of Moderate rain on 9 & 10 May. Maximum temperature would be in the range of 35°C to 39°C and the minimum temperature would be between 24°C-28°C. Generally cloudy sky will appear for the next five days. Wind is expected with the speed of 10-12 Kmph from, Southeast & East directions.
-----------------	---

General Advisory:

THUNDERSTORM WARNING:

Heavy rain is likely to occur at isolated places over the district on 9 & 10 May 2026. Farmers are advised to act swiftly to repair the water channels, farm ponds and other water bodies to harvest the rain water. During these days farmers are advised to avoid irrigation, intercultural

operations and application of plant protection measures to the standing crops, provide adequate drainage facilities in the fields to avoid water stagnation due to heavy rainfall, do not allow the animals to grazing outside, standing near the trees or electric poles & provide safe shelter.

Wind Warning:

Banana, Papaya, Moringa & Sugarcane

Thunderstorm with Lightning with Strong Gusty Wind with a speed of 40 to 50 kmph is expected during 6, 7, 8, 9 & 10 May 2026.

Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya and Moringa, to prevent from lodging.

Gourds:

For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support.

Weather Based Agro-Advisories: (Maize, Horticulture Crops, Tomato, Banana & Animal Husbandry)

Maize:

Postpone the sowing / Planting Activities.

Drain the excess rain water from the field.

Postpone Plant protection activities such as Fertilizer application and Pesticides spraying for next 5 days.

Horticulture Crops:

Avoid irrigating the fields.

Drain the excess rain water from the field.

Postpone Plant protection activities such as Fertilizer application and Pesticides spraying for next 5 days.

Tomato:

Due to heavy rainfall, there are chances of water logging. Provide adequate drainage facility. Postpone the inter-cultural / plant protection activities like fertilizer and pesticide application, until the rain completely stops. Harvest the matured crops immediately on getting heavy rainfall forecast and keep the harvested produce in a safe place.

Banana:

Since Heavy rainfall is expected to occur at one or two places, there are chances of water logging and stacking of banana crops. Provide adequate drainage facility and avoid fertilizer and pesticide application. Remove and destruct the decayed and rotten leaves. Provide support to 5 months old trees using wooden poles.

Weather based Agro Advisory:

Heavy rainfall is expected on 9 & 10 May 2026. During these days farmers are advised to avoid irrigation, intercultural operations and application of plant protection measures to the standing crops.

Provide adequate drainage facilities in the field and renovate the rain water harvesting structures to avoid water stagnation due to heavy rainfall. Try to harvest the rainwater in the farm ponds / water bodies.

Provide support to crops likes Banana, Papaya and sugarcane.

Animal Husbandry:

To protect the livestock from thunderstorms and lightning Farmers are requested to avoid grazing the animals during rains, particularly grazing sheep and goat in the flock which are more prone to lightning shock.

Farmers should avoid tethering animals near barbed wire fencing posts and trees. Trees near the animal sheds should be pruned to avoid accidental falls over during heavy wind.

Further trees like rain trees and vathanarayanan are more prone to uprooting. More chance of lightning near the hilly area and undulated landscapes than in plains.

Fence with wooden poles and metal wires, it is advisable to check the condition of the buried parts at most once in 5 years and do the necessary replacement before the thunderstorm season begins.

DINDIGUL

SMS Advisory	Chances of Moderate rain during 6, 9 & 10 May. Maximum temperature would be in the range of 33°C to 37°C and the minimum temperature would be between 23°C-27°C. Cloudy sky will appear for the next five days. Wind is expected with the speed of 8-12 Kmph from, Southeast & South directions.
--------------	--

General Advisory:

Heavy rain is likely to occur at isolated places over the district during 6, 9 & 10 May 2026. Farmers are advised to act swiftly to repair the water channels, farm ponds and other water bodies to harvest the rain water. During these days farmers are advised to avoid irrigation, intercultural operations and application of plant protection measures to the standing crops, provide adequate drainage facilities in the fields to avoid water stagnation due to heavy rainfall, do not allow the animals to grazing outside, standing near the trees or electric poles & provide safe shelter.

Wind Warning:

Banana, Papaya, Moringa & Sugarcane

Thunderstorm with Lightning with Strong Gusty Wind with a speed of 40 to 50 kmph is expected during 6, 9 & 10 May 2026.

Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya and Moringa, to prevent from lodging.

Gourds:

For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support.

Weather Based Agro-Advisories: (Maize, Horticulture Crops, Coconut, Horticulture crop & Animal Husbandry)

Maize:

Postpone the sowing / Planting Activities.

Drain the excess rain water from the field.

Postpone Plant protection activities such as Fertilizer application and Pesticides spraying for next 5 days.

Horticulture Crops:

Avoid irrigating the fields.

Drain the excess rain water from the field.

Postpone Plant protection activities such as Fertilizer application and Pesticides spraying for next 5 days.

Coconut:

Heavy rainfall is expected in the coming days, coconut growers are advised to safely remove the old and decayed fronds. Provide adequate drainage facilities in the field.

Horticulture crop:

Since heavy rainfall expected, farmers are advised to provide adequate drainage facilities in the fields to avoid water stagnation and farmers are advised to repair the water channels.

Avoid spraying of fertilizer, fungicide and pesticide during the rainy days.

Weather based Agro Advisory:

Heavy rainfall is expected during 6, 9 & 10 May 2026. During these days farmers are advised to avoid irrigation, inter-cultural operations and application of plant protection measures to the standing crops.

Provide adequate drainage facilities in the field and renovate the rain water harvesting structures to avoid water stagnation due to heavy rainfall. Try to harvest the rainwater in the farm ponds / water bodies.

Provide support to crops likes Banana, Papaya and sugarcane.

Animal Husbandry:

To protect the livestock from thunderstorms and lightning Farmers are requested to avoid grazing the animals during rains, particularly grazing sheep and goat in the flock which are more prone to lightning shock.

Farmers should avoid tethering animals near barbed wire fencing posts and trees. Trees near the animal sheds should be pruned to avoid accidental falls over during heavy wind.

Further trees like rain trees and vathanarayanan are more prone to uprooting. More chance of lightning near the hilly area and undulated landscapes than in plains.

Fence with wooden poles and metal wires, it is advisable to check the condition of the buried parts at most once in 5 years and do the necessary replacement before the thunderstorm season begins.

SIVAGANGA

SMS Advisory	Chances of Moderate rain on 9 & 10 May. Maximum temperature would be in the range of 32°C to 36°C and the minimum temperature would be between 23°C-27°C. Generally cloudy sky will appear for the next five days. Wind is expected with the speed of 10-12 Kmph from, Southeast & South directions.
--------------	--

General Advisory:

Heavy rain is likely to occur at isolated places over the district on 9 & 10 May 2026. Farmers are advised to act swiftly to repair the water channels, farm ponds and other water bodies to harvest the rain water. During these days farmers are advised to avoid irrigation, intercultural operations and application of plant protection measures to the standing crops, provide adequate drainage facilities in the fields to avoid water stagnation due to heavy rainfall, do not allow the animals to grazing outside, standing near the trees or electric poles & provide safe shelter.

Wind Warning:

Banana, Papaya, Moringa & Sugarcane

Thunderstorm with Lightning with Strong Gusty Wind with a speed of 40 to 50 kmph is expected on 9 & 10 May 2026.

Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya and Moringa, to prevent from lodging.

Gourds:

For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate supportt.

Weather Based Agro-Advisories: (Maize, Horticulture Crops, Tomato, Banana & Animal Husbandry)

Maize:

Postpone the sowing / Planting Activities.

Drain the excess rain water from the field.

Postpone Plant protection activities such as Fertilizer application and Pesticides spraying for next 5 days.

Horticulture Crops:

Avoid irrigating the fields.

Drain the excess rain water from the field.

Postpone Plant protection activities such as Fertilizer application and Pesticides spraying for next 5 days

Tomato:

Due to heavy rainfall, there are chances of water logging. Provide adequate drainage facility. Postpone the intercultural / plant protection activities like fertilizer and pesticide application, until the rain completely stops. Harvest the matured crops immediately on getting heavy rainfall forecast and keep the harvested produce in a safe place.

Banana:

Since Heavy rainfall is expected to occur at one or two places, there are chances of water logging and stacking of banana crops. Provide adequate drainage facility and avoid fertilizer and pesticide application. Remove and destruct the decayed and rotten leaves. Provide support to 5 months old trees using wooden poles.

Weather based Agro Advisory:

Heavy rainfall is expected on 9 & 10 May 2026. During these days farmers are advised to avoid irrigation, intercultural operations and application of plant protection measures to the standing crops.

Provide adequate drainage facilities in the field and renovate the rain water harvesting structures to avoid water stagnation due to heavy rainfall. Try to harvest the rainwater in the farm ponds / water bodies.

Provide support to crops likes Banana, Papaya and sugarcane.

Animal Husbandry:

To protect the livestock from thunderstorms and lightning Farmers are requested to avoid grazing the animals during rains, particularly grazing sheep and goat in the flock which are more prone to lightning shock.

Farmers should avoid tethering animals near barbed wire fencing posts and trees. Trees near the animal sheds should be pruned to avoid accidental falls over during heavy wind.

Further trees like rain trees and vathanarayanan are more prone to uprooting. More chance of lightning near the hilly area and undulated landscapes than in plains.

Fence with wooden poles and metal wires, it is advisable to check the condition of the buried parts at most once in 5 years and do the necessary replacement before the thunderstorm season begins.

THENI

SMS Advisory	Chances of Moderate rain during 6, 7, 8, 9 & 10 May. Maximum temperature would be in the range of 31°C to 35°C and the minimum temperature would be between 20°C-24°C. Cloudy sky will appear for the next five days. Wind is expected with the speed of 8-12 Kmph from, Southeast & South directions.
-----------------	--

General Advisory:

Heavy rain is likely to occur at isolated places over the district during 6, 7, 8, 9 & 10 May 2026. Farmers are advised to act swiftly to repair the water channels, farm ponds and other water bodies to harvest the rain water. During these days farmers are advised to avoid irrigation, intercultural operations and application of plant protection measures to the standing crops, provide adequate drainage facilities in the fields to avoid water stagnation due to heavy rainfall, do not allow the animals to grazing outside, standing near the trees or electric poles & provide safe shelter.

Wind Warning:

Banana, Papaya, Moringa & Sugarcane

Thunderstorm with Lightning with Strong Gusty Wind with a speed of 40 to 50 kmph is expected during 6, 7, 8, 9 & 10 May 2026.

Hence farmers are advised to provide propping in sugarcane and to provide adequate support for crops like Banana, Papaya and Moringa, to prevent from lodging.

Gourds:

For Gourds check the structures /pandhals for damages in the strings and poles and provide adequate support.

Weather Based Agro-Advisories: (Maize, Horticulture Crops, Tomato, Banana & Animal Husbandry)

Maize:

Postpone the sowing / Planting Activities.

Drain the excess rain water from the field.

Postpone Plant protection activities such as Fertilizer application and Pesticides spraying for next 5 days.

Horticulture Crops:

Avoid irrigating the fields.

Drain the excess rain water from the field.

Postpone Plant protection activities such as Fertilizer application and Pesticides spraying for next 5 days.

Tomato:

Due to heavy rainfall, there are chances of water logging. Provide adequate drainage facility. Postpone the intercultural / plant protection activities like fertilizer and pesticide application, until the rain completely stops. Harvest the matured crops immediately on getting heavy rainfall forecast and keep the harvested produce in a safe place.

Banana:

Since Heavy rainfall is expected to occur at one or two places, there are chances of water logging and stacking of banana crops. Provide adequate drainage facility and avoid fertilizer and pesticide application. Remove and destruct the decayed and rotten leaves. Provide support to 5 months old trees using wooden poles.

Weather based Agro Advisory:

Heavy rainfall is expected during 6, 7, 8, 9 & 10 May 2026. During these days farmers are advised to avoid irrigation, intercultural operations and application of plant protection measures to the standing crops.

Provide adequate drainage facilities in the field and renovate the rain water harvesting structures to avoid water stagnation due to heavy rainfall. Try to harvest the rainwater in the farm ponds / water bodies.

Provide support to crops likes Banana, Papaya and sugarcane.

Animal Husbandry:

To protect the livestock from thunderstorms and lightning Farmers are requested to avoid grazing the animals during rains, particularly grazing sheep and goat in the flock which are more prone to lightning shock.

Farmers should avoid tethering animals near barbed wire fencing posts and trees. Trees near the animal sheds should be pruned to avoid accidental falls over during heavy wind.

Further trees like rain trees and vathanarayanan are more prone to uprooting. More chance of lightning near the hilly area and undulated landscapes than in plains.

Fence with wooden poles and metal wires, it is advisable to check the condition of the buried parts at most once in 5 years and do the necessary replacement before the thunderstorm season begins.

KANYAKUMARI

AGRO ADVISORIES

S. N.	CROP	STAGES	ADVISORIES
1.	Horticultural crops.	Planting stage	As 12 to 18 km / hr wind speed is anticipated, it is advised to provide staking to newly planted horticultural plants.
2.	Agricultural and Horticultural crops.	All stages	Spraying of insecticides, weedicides and application of fertilizer should be postponed due to rain.

3.	Pepper	All stages	The runner shoots may be protected from spreading on the ground by coiling it on supports.
4.	Banana	Harvesting stage	Harvested banana bunches should be stored in dry places to prevent disease incidence due to rainfall.
5.	Coconut	All stages	Basins can be formed and green manure sowing may be taken up in the basins.
6.	Rice	Nursery stage	Care must be taken to avoid stagnation of water in the seed bed.
7.	Cinnamon	Planting stage	One year old seedlings may be transplanted under partial shade
8.	Rubber	Tapping Stage	To protect the quality of rubber latex shade provision may be given.

SMS Advisory:

Ginger: Provide proper drainage channels to avoid water logging during the rainy season.

KARAIKAL

Agro Advisory

Crop/livestock	Stage/Particulars	Agro Advisory
Cotton	Vegetative growth Sucking pest	To manage weeds in cotton, apply post emergence herbicide pyriithobac sodium @ 62.5g a.i./ha + quizalofop ethyl @ 50 g a.i./ha at 20-25 DAS.
		To manage sucking pest in cotton, spray azadirachtin 0.03 (300 ppm) @ 2500-5000 ml/ha or imidacloprid 17.8 SL @100-125 ml/ha or thiamethoxam 25 WG @100 g/ha.
Rice fallow pulses	Flowering and pod development	To control powdery mildew, spray NSKE @ 5% or neem oil @ 3% twice at 10 days interval or spray Eucalyptus leaf extract @ 10% at initiation of disease and 10 days later or Spray any one of the following fungicides at initiation of disease and second spray at 15 days later. Carbendazim 50 WP @ 500g/ha Wettable sulphur 80 WP @ 1500g/ha Propiconazole 25% EC @ 500ml/ha
		To control leaf crinkle, rogue out the infected plants up to 45 days. Foliar spray of notchi leaf extract 10% at 30 DAS or neem formulation 3 ml/lit or Methyl demeton 25 EC @500 ml/ha or Dimethoate 30 EC @ 500 ml/ha repeat after 15 days, if necessary.
		To manage yellow mosaic virus disease in black gram and green gram, spray Methyldematton 25%EC 1ml/l or Dimethoate 30%EC 1ml/l to manage vectors like

		whitefly and aphids. Two percent of DAP spray (Dissolve 4.5 kg of DAP in 10 litres of water previous day night and at morning the supernatant liquid shall be mixed with 190 litres of water) may be done at the time of flowering and another dose at fifteen days after first spray for rice fallow pulses.
Sesame	Vegetative stage	In Sesame, to manage Jassids, apply Acetamiprid 20 SP @ 0.3g/l.
Snake gourd and bitter gourd	Flowering stage	Spray ethrel @ 100 ppm (1 ml/10 litres of water) in snake gourd and bitter gourd for four times commencing from 15 th day after sowing at weekly intervals to increase female flowers.
Ribbed gourd	Flowering stage	Spray ethrel @ 250 ppm (2.5 ml/10 litres of water) in ribbed gourd for four times commencing from 15 th day after sowing at weekly intervals to increase female flowers.
Brinjal	Phomopsis blight	To manage Phomopsis blight in brinjal, spray <i>Pseudomonas fluorescens</i> or <i>Bacillus subtilis</i> @ 5g or 5ml/ l.
	Shoot borer	To manage Brinjal shoot borer, spray Emamectin Benzoate 4% SG @ 0.4g/l.
Coconut	Bearing stage	Root feeding of coconut tonic (200 ml / tree) in 3 months interval to avoid button shedding in coconut.
Livestock	Heat stress	Heat stress is likely to occur in cattle due to prevailing hot weather. Therefore, following steps can be followed to protect livestock from summer heat. Cattle can be protected from heat by spreading straw, dry grass, palm fronds, coconut leaves or coir waste on the surface of cattle sheds and sprinkling water over it. Cattle can be sent for grazing between 06 am to 10 am and 04 pm to 06 pm. Roofs should be erected over the drinking troughs for livestock to provide cold drinking water to the livestock. Cattle should be bathed twice daily. If cattle show signs of heat stress such as excessive drinking of water, rapid breathing, excessive salivation, loss of appetite, shivering and falling down, then the affected animals should be treated at the nearest veterinary hospital.

SMS: Light drizzle to moderate rain is expected from 06th to 10th May.

PUDUCHERRY


Agro Advisory

Crop/livestock	Stage/Particulars	Agro Advisory
Sesame	Vegetative stage	In Sesame, to manage Jassids, apply Acetamiprid 20 SP @ 0.3g/l.
Sugarcane	Vegetative stage	To control shoot borer in sugarcane, soil application of Cartaphydrochloride 4G 10kg/hectare may be done.
Cassava	Vegetative stage	To manage red mite in Cassava, apply Pongamia oil 1 ml/l or spray Propargite 1.5 ml and Sulphur 2.5 g/l.
Brinjal	Flowering to Bearing stage	To manage Brinjal shoot borer, spray Emamectin Benzoate 4% SG @ 0.4g/l.
Snake gourd and bitter gourd	Flowering stage	Spray ethrel @ 100 ppm (1 ml/10 litres of water) in snake gourd and bitter gourd for four times commencing from 15th day after sowing at weekly intervals to increase female flowers.
Ribbed gourd	Flowering stage	Spray ethrel @ 250 ppm (2.5 ml/10 litres of water) in ribbed gourd for four times commencing from 15th day after sowing at weekly intervals to increase female flowers.
Coconut	Bearing stage	Root feeding of coconut tonic (200 ml / tree) in 3 months interval to avoid button shedding in coconut.
Livestock	Heat stress	<p>Heat stress is likely to occur in cattle due to prevailing hot weather. Therefore, following steps can be followed to protect livestock from summer heat.</p> <p>Cattle can be protected from heat by spreading straw, dry grass, palm fronds, coconut leaves or coir waste on the surface of cattle sheds and sprinkling water over it.</p> <p>Cattle can be sent for grazing between 06 am to 10 am and 04 pm to 06 pm.</p> <p>Roofs should be erected over the drinking troughs for livestock to provide cold drinking water to the livestock.</p> <p>Cattle should be bathed twice daily.</p> <p>If cattle show signs of heat stress such as excessive drinking of water, rapid breathing, excessive salivation, loss of appetite, shivering and falling down, then the affected animals should be treated at the nearest veterinary hospital.</p>


SMS: Light drizzle to light rain is expected from 06th to 10th May.

District level Heavy Rainfall Warning for next 5 days


Day 1 (05.05.2026)




Day 2 (06.05.2026)



Day 3 (07.05.2026)




Day 4 (08.05.2026)



Meteorological Subdivision Level Rainfall Weather Forecast for next 5 days of RMC

Chennai



District Forecast & Warning for Next 5 Days issued on: 05.05.2026

NAME OF THE DISTRICT	RAINFALL INTENSITY	06.05.2026	07.05.2026	08.05.2026	09.05.2026	10.05.2026
		WED	THU	FRI	SAT	SUN
TIRUVALLUR	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
CHENNAI	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light
	Probability	Likely	Likely	Likely	Likely	Likely
CHENGALPATTU	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light
	Probability	Likely	Likely	Likely	Likely	Likely
KANCHEEPURAM	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
VILLUPURAM	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
PUDUCHERRY	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light	Light-Moderate
	Probability	Likely	Likely	Very Likely	Likely	Likely
CUDDALORE	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
MAYILADUTHURAI	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
NAGAPATTINAM	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
KARAIKAL	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely

TIRUVARUR	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
THANJAVUR	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
PUDUKKOTTAI	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
KALLAKURUCHI	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
VELLORE	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
RANIPET	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
TIRUPPATTUR	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
TIRUVANNAMALAI	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
ARIYALUR	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
PERAMBALUR	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
TIRUCHIRAPALLI	Intensity	ISOL HEAVY	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
KARUR	Intensity	ISOL HEAVY	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
NAMAKKAL	Intensity	ISOL	Light-	Light-	Light-	Light-

		HEAVY	Moderate	Moderate	Moderate	Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
SALEM	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
DHARMAPURI	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
KRISHNAGIRI	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate	Light-Moderate
	Probability	Likely	Likely	Likely	Likely	Likely
ERODE	Intensity	ISOL HEAVY	Light-Moderate	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
NILGIRIS	Intensity	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
COIMBATORE	Intensity	Light-Moderate	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
TIRUPPUR	Intensity	Light-Moderate	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
THENI	Intensity	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
DINDIGUL	Intensity	ISOL HEAVY	Light-Moderate	Light-Moderate	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
MADURAI	Intensity	Light-Moderate	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
VIRUDHUNAGAR	Intensity	Light-Moderate	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
SIVAGANGA	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely

TENKASI	Intensity	Light-Moderate	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
RAMANATHAPURAM	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
THOOTHUKUDI	Intensity	Light-Moderate	Light-Moderate	Light-Moderate	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
TIRUNELVELI	Intensity	Light-Moderate	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely
KANYAKUMARI	Intensity	Light-Moderate	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY	ISOL HEAVY
	Probability	Likely	Likely	Likely	Likely	Likely

Warning Colour Codes	हिन्दी	मराठी
WARNING (TAKE ACTION)	चेतावनी	चेतावणी
ALERT (BE PREPARED)	सतर्क रहें	सतर्क असावे
WATCH (BE UPDATED)	निगरानी रखें	लक्ष्य असावे
NO WARNING (NO ACTION)	कोई चेतावनी नहीं	धोक्याची सूचना नाही



Heavy Rain



Heavy Snow



Thunderstorm



Dust Storm



Strong Winds



Visibility



Cyclone



Squall/ Hail



Frost



Cold Wave



Heat Wave



Sea State